







Z.COM HOME	MOLECULAR DIVERSITY	DIVERSITY INFORMATION PAGES		
Articles thru 2000	Articles 2001	Reviews	SPOS Papers	SPPS Papers
Robotics/Automation	Classics	Patents	Books	Procedures
Symposia	Companies	Suppliers	Scientists	Graffiti Wall
Links	<u>Jobs</u>	Hot News	Site News	Fun

## Selection of papers on solid phase peptide synthesis published after 1996

Solid phase peptide synthesis was invented by Bruce Merrifield in 1963 and became very quickly the routine tool for preparation of peptides and later small proteins. Surprisingly, solid phase synthesis of small organic molecules was accepted only after another 30 years. The compilation of papers dealing with solid phase peptide synthesis would be extremely long. Luckily, Gregg Fields edited the book Solid Phase Peptide Synthesis in the series Methods in Enzymology (Fields, G.B. (1997) Solid-Phase Peptide Synthesis. Academic Press, San Diego. #9830), which covers all aspects of this methodology. We decided to cover here only papers published after publication of Fields' book. This compilation will probably never be complete, but we will try to cover the most important methodological papers. This list is arranged by the year and the name of the first author. Any omissions, mistakes and inconsistencies please communicate to the authors of compilation (michal@5z.com). (If you are pointing to any reference, please use its ID# -- the last number in the reference.) You are welcome to use this compilation in your work. Proper way to quote it in your paper is: Lebl M., Leblova Z.: Dynamic database of references in solid phase synthesis. Internet http://www.5z.com.

## Recently added entries are marked in bold

- Balaspiri, L., & Langel, U. (2001) Synthesis of b-amyloid precursor peptide and presenilin segments. J. Peptide Sci., 7, 58-60. #16430
- 2. Bullesbach, E.E., & Schwabe, C. (2001) Synthesis and conformational analysis of the insulin-like 4 gene product. J. Pept. Res., 57, 77-83. #16258
- 3. Burgess, K., Han, I., Zhang, A., Zheng, W.H., Shanmugam, K., Featherstone, M.S., & Saragovi, H.U. (2001) DiSSiMiL: Diverse Small Size Mini-Libraries applied to simple and rapid epitope mapping of a monoclonal antibody. J. Pept. Res., 57, 68-76. #16259
- 4. Dekker, N., Cox, R.C., Kramer, R.A., & Egmond, M.R. (2001) Substrate specificity of the integral membrane protease OmpT determined by spatially addressed peptide libraries. Biochemistry, 40, 1694-1701. #16271
- Demarcus, M., Ganadu, M.L., Mura, G.M., Porcheddu, A., Quaranta, L., Reginato, G., & Taddei, M. (2001) Small ring constrained peptidomimetics. Synthesis of epoxy peptidomimetics, inhibitors of cysteine proteases. J. Org. Chem., 66, 697-706. #16273
- 6. Dolence, E.K., Dolence, J.M., & Poulter, C.D. (2001) Solid-phase synthesis of a radiolabeled, biotinylated, and farnesylated Ca(1)a(2)X peptide substrate for Ras- and a-mating factor converting enzyme. Bioconjug. Chem., 12, 35-43. #16275
- 7. Futaki, S., Youjun, Z., & Sugiura, Y. (2001) Detecting a tag on a channel opening: Blockage of the biotinylated channels by strepravidin. Tetrahedron Lett., 42, 1563-1565. #16423
- 8. Gatos, D., & Tzavara, C. (2001) Comparison of the solid-phase fragment condensation and phase-change approaches in the synthesis of salmon I calcitonin. J. Pept. Res., 57, 168-174. #16427





- 9. Geginat, G., Schenk, S., Skoberne, M., Goebel, W., & Hof, H. (2001) A novel approach of direct ex vivo epitope mapping identifies dominant and subdominant CD4 and CD8 T cell epitopes from Listeria monocytogenes. Journal of Immunology, 166, 1877-1884. #16288
- 10. Henlin, J.M., Boutin, J.A., Kucharczyk, N., Desmet-Beaufort, C., Loynel, A., Bertrand, M., Genton, A., Tucker, G.C., Atassi, G., & Fauchere, J.L. (2001) From peptide libraries to optimized nonpeptide ligands in the search for Sfarnesyltransferase inhibitors. J. Pept. Res., 57, 85-96. #16425
- 11. Kitagawa, K., Aida, C., Fujiwara, H., Yagami, T., Futaki, S., Kogire, M., Ida, J., & Inoue, K. (2001) Facile solid-phase synthesis of sulfated tyrosine-containing peptides: Total synthesis of human big gastrin-II and cholecystokinin (CCK)-39 (1,2). J. Org. Chem., 66, 1-10. #16313
- 12. Kumar, K.S., Roice, M., Sasikumar, P.G., Poduri, C.D., Sugunan, V.S., Pillai, V.N.R., & Das, M.R. (2001) Syntheses of immunodominant peptide regions of hepatitis C viral pathogens using PS-BDODMA resin: A single peptide derived from the conserved domain (E2/NS1) was highly effective in detecting anti-HCV antibodies. J. Pept. Res., 57, 140-150. #16426
- 13. Lelievre, D., Bure, C., Laot, F., & Delmas, A. (2001) Synthesis of peptide di-aldehyde precursor for stepwise chemoselective ligations via oxime bonds. Tetrahedron Lett., 42, 235-238. #16326
- 14. Licha, K., Hessenius, C., Becker, A., Henklein, P., Bauer, M., Wisniewski, S., Wiedenmann, B., & Semmler, W. (2001) Synthesis, characterization, and biological properties of cyanine-labeled somatostatin analogues as receptor-targeted fluorescent probes. Bioconjug. Chem., 12, 44-50. #16329
- 15. Marchetto, R., Nicolas, E., Castillo, N., Bacardit, J., Navia, M., Vila, J., & Giralt, E. (2001) Two short peptides including segments of subunit A of Escherichia coli DNA gyrase as potential probes to evaluate the antibacterial activity of quinolones. J. Peptide Sci., 7, 27-40. #16441
- 16. McNamara, L.M.A., Andrews, M.J.I., Mitzel, F., Siligardi, G., & Tabor, A.B. (2001) Unexpected conformational properties of a peptide constrained by an aliphatic link between the i and i+4 positions. Tetrahedron Lett., 42, 1591-1593. #16424
- 17. Nilsson, M.R., Nguyen, L.L., & Raleigh, D.P. (2001) Synthesis and purification of amyloidogenic peptides. Anal. Biochem., 288, 76-82. #16355
- 18. Qiu, W., Gu, X., Soloshonok, V.A., Carducci, M.D., & Hruby, V.J. (2001) Stereoselective synthesis of conformationally constrained reverse turn dipeptide mimetics. Tetrahedron Lett., 42, 145-148. #16181
- 19. Royo, M., Jimenez, J.C., Lopez-Macia, A., Giralt, E., & Albericio, F. (2001) Solid-phase synthesis of peptides containing alpha, beta-didehydroamino acids. Eur. J. Org. Chem., 45-48. #16373
- 20. Shuker, S.B., Esterbrook, J., & Gonzalez, J. (2001) Solid-phase synthesis of a novel peptide substituted calix[4]arene. Synlett, 210-213. #16382
- 21. Aimoto, S., Hasegawa, K., Li, X. & Kawakami, T. (2000) Synthesis of phosphorylated CRE BP1(19-106) amide using a phosphopeptide thioester prepared directly by an Fmoc solid-phase method. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 82-83). Kluwer Academic Publisher, Dordrecht. #15701
- 22. Ajikumar, P.K., & Devaky, K.S. (2000) EGDMA-cross-linked polystyrene resin: An efficient support for gel phase peptide synthesis. Proc. Indian Acad. Sci. -Chem. Sci., 112, 465-474. #15819
- 23. Alam, M.R., Maeda, M., & Sasaki, S. (2000) DNA-binding peptides searched from the solid-phase combinatorial library with the use of the magnetic beads attaching the target duplex DNA. Bioorg, Med. Chem., 8, 465-473. #14913
- 24. Albericio, F. (2000) Orthogonal protecting groups for Na-amino and C-terminal carboxyl functions in solid-phase peptide synthesis. Biopolymers (Pept. Sci.), 55, 123-139. #16434
- 25. Albericio, F. (2000) Orthogonal protecting groups for N alpha-amino and C-terminal carboxyl functions in solid-phase peptide synthesis. Biopolymers, 55, 123-139. #16237
- Alsina, J., Yokum, T.S., Albericio, F., & Barany, G. (2000) A modified Backbone Amide Linker (BAL) solid-phase peptide synthesis strategy accommodating prolyl, N-alkylamino acyl, or histidyl derivatives at the C-terminus. Tetrahedron Lett., 41, 7277-7280. #15629

- Alsina, J., Yokum, T.S., Albericio, F. & Barany, G. (2000) Backbone Amide Linker (BAL) methodology to accommodate C-terminal hindered, unreactive, and/or sensitive modification. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 102-103). Kluwer Academic Publisher, Dordrecht. #15707
- 28. Andersson, L., Blomberg, L., Flegel, M., Lepsa, L., Nilsson, B., & Verlander, M. (2000) Large-scale synthesis of peptides. Biopolymers (Pept. Sci.), 55, 227-250. #16440
- 29. Andersson, L., Blomberg, L., Flegel, M., Lepsa, L., Nilsson, B., & Verlander, M. (2000) Large-scale synthesis of peptides. Biopolymers, 55, 227-250. #16238
- 30. Ando, S., Nakahara, Y., Ito, Y., & Ogawa, T. (2000) Solid-phase synthesis of the glycopeptide of human glycophorin AM bearing the consecutive sialyl-T antigen. Carbohyd. Res., 329, 773-780. #16239
- 31. Anichini, B., Ricci, R., Fabbri, G., Balacco, G., Mauro, S., Triolo, A., Altamura, M., Maggi, C.A., & Quartara, L. (2000) Solid-phase synthesis of MEN 11270, a new cyclic peptide kinin B2 receptor antagonist. J. Peptide Sci., 6, 612-620. #16152
- 32. Annis, I. & Barany, G. (2000) Alternative solid-phase reagents for formation of intramolecular sulfur-sulfur bridges in peptides under mild conditions. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 96-97). Kluwer Academic Publisher, Dordrecht. #15704
- 33. Antopolsky, M., & Azhayev, A. (2000) Stepwise solid-phase synthesis of peptide-oligonucleotide phosphorothioate conjugates employing Fmoc peptide chemistry. Tetrahedron Lett., 41, 9113-9117. #15823
- 34. Arnon, R., Tarrab-Hazdai, R., & Steward, M. (2000) A mimotope peptide-based vaccine against Schistosoma mansoni: Synthesis and characterization. Immunology, 101, 555-562. #16245
- 35. Arunan, C., & Pillai, V.N.R. (2000) Synthesis of acyl carrier protein fragment 65-74 on a flexible cross-linked polystyrene support: Comparison with Merrifield resin. Tetrahedron, 56, 3005-3011. #14623
- 36. Arunan, C., Naaraj, R., & Pillai, V.N.R. (2000) Solid-phase synthesis of a modified 13-residue seminalplasmin fragment on 1,6-hexanediol diacrylate-crosslinked polystyrene support. Peptides, 21, 773-777. #15758
- 37. Balaspiri, L., Tegyei, Z., Dux, M., Jancso, G., Jozsa, R., Csernus, V., & Mess, B. (2000) Synthesis of chicken galanin and its N-and C-terminal segments, and preparation of their antisera. Lett. Peptide Sci., 7, 23-26. #15431
- 38. Balse, P., Han, G., Hruby, V.J. & Sun, L. (2000) Diisopropylethylamine salts for protected amino acids: Direct synthesis of peptides on solid-phase. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 80-81). Kluwer Academic Publisher, Dordrecht. #15700
- 39. Balse, P.M., Kim, H.J., Han, G., & Hruby, V.J. (2000) Evaluation of new base-labile 2-(4-nitrophenylsulfonyl)ethoxycarbonyl (Nsc)-amino acids for solid-phase peptide synthesis. J. Pept. Res., 56, 70-79. #15444
- Baumann, M., Dietrich, U., Koenigs, C. & Griesinger, C. (2000) Combinatorial peptide synthesis, NMR-aided screening and testing of low molecular weight RNA-ligands. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 170-171). Kluwer Academic Publisher, Dordrecht. #15717
- 41. Bechinger, B. (2000) Biophysical investigations of membrane perturbations by polypeptides using solid-state NMR spectroscopy. Molecular Membrane Biology, 17, 135-142. #15832
- 42. Beebe, K.D., Wang, P., Arabaci, G., & Pei, D.H. (2000) Determination of the binding specificity of the SH2 domains of protein tyrosine phosphatase SHP-1 through the screening of a combinatorial phosphotyrosyl peptide library. Biochemistry, 39, 13251-13260. #15833
- 43. Beligere, G.S., & Dawson, P.E. (2000) Design, synthesis, and characterization of 4-ester CI2, a model for backbone hydrogen bonding in protein alpha-helices. J. Amer. Chem. Soc., 122, 12079-12082. #16250
- 44. Besser, D., Muller, B., Agricola, I., & Reissmann, S. (2000) Synthesis of differentially protected N-acylated reduced pseudodipeptides as building units for backbone cyclic peptides. J. Peptide Sci., 6, 130-138. #14943
- 45. Besser, D., Muller, B., Kleinwachter, P., Greiner, G., Seyfarth, L., Steinmetzer, T., Arad, O., & Reissmann, S. (2000) Synthesis

- and characterization of octapeptide somatostatin analogues with backbone cyclization: Comparison of different strategies, biological activities and enzymatic stabilities. Journal fur Praktische Chemie-Practical Applications and Applied Chemistry, 342, 537-545. #15837
- 46. Bibbs, L., Ambulos, N.P., Kates, S.A., Khatri, A., Medzihradszky, K.F., Osapay, G. & Weintraub, S.T. (2000) Strategies for the synthesis of labeled peptides. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 113-114). Kluwer Academic Publisher, Dordrecht. #15710
- 47. Bishop, B.M., McCafferty, D.G., & Erickson, B.W. (2000) 4'-Aminomethyl-2,2'-bipyridyl-4-carboxylic acid (Abc) and related derivatives: Novel bipyridine amino acids for the solid-phase incorporation of a metal coordination site within a peptide backbone. Tetrahedron, 56, 4629-4638. #14946
- 48. Blaskovich, M.A., Boatman, P.D., Cao, B., Eguchi, M., Kim, H.O., Lee, M., Little, T., Mathew, F., McCann, I., Mehlin, C., Nakanishi, H., Nelson, S., Nguyen, M., Ogbu, C., Qabar, M.N., Ruan, F., Shea, J.P., Stasiak, M., Urban, J. & Kahn, M. (2000) Highly efficient and versatile construction of secondary structure peptide mimetic libraries: Application to biology and drug development. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 191-193). Kluwer Academic Publisher, Dordrecht. #15726
- 49. Blondelle, S.E., Crooks, E., Aligue, R., Agell, N., Bachs, O., Esteve, V., Tejero, R., Celda, B., Pastor, M.T., & Perez-Paya, E. (2000) Novel, potent calmodulin antagonists derived from an all-D hexapeptide combinatorial library that inhibit in vivo cell proliferation. Activity and structural characterization. J. Pept. Res., 55, 148-162. #14266
- 50. Blondelle, S.E., Esteve, V., Celda, B., Pastor, M.T., & Perez-Paya, E. (2000) Influence of the hydrophylic face on the folding ability and stability of a-helix bundles: Relevance to the peptide catalytic activity. J. Pept. Res., 56, 121-131. #15789
- 51. Bonner, G.G., Davis, P., Stropova, D., Edsall, S., Yamamura, H.I., Porreca, F., & Hruby, V.J. (2000) Opiate aromatic pharmacophore structure-activity relationships in CTAP analogues determined by topographical bias, two-dimensional NMR, and biological activity assays. J. Med. Chem., 43, 569-580. #14952
- 52. Bordusa, F. (2000) Nonconventional amide bond formation catalysis: Programming enzyme specificity with substrate mimetics. Braz. J. Med. Biol. Res., 33, 469-485. #14953
- 53. Borgia, J.A., Oegema, T.R.Jr. & Fields, G.B. (2000) Synthesis and application of a glycoprotein derived from the proteoglycan linkage structure. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 773-774). Kluwer Academic Publisher, Dordrecht. #15751
- 54. Bourel, L., Carion, O., Gras-Masse, H., & Melnyk, O. (2000) The deprotection of Lys(Mtt) revisited. J. Peptide Sci., 6, 264-270. #14956
- 55. Bourne, G.T., McGeary, R.P., Golding, S.W., Meutermans, W.D.F., Alewood, P.F. & Smythe, M.L. (2000) An evaluation of a novel safety catch linker for development of cyclic peptide libraries. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 98-99). Kluwer Academic Publisher, Dordrecht. #15705
- 56. Boykins, R.A., Joshi, M., Syin, C., Dhawan, S., & Nakhasi, H. (2000) Synthesis and construction of a novel multiple peptide conjugate system: strategy for a subunit vaccine design. Peptides, 21, 9-17. #14958
- 57. Boysen, R.I., & Hearn, M.T.W. (2000) Direct characterisation by electrospray ionisation mass spectroscopy of mercuropolypeptide complexes after deprotection of acetamidomethyl groups from protected cysteine residues of synthetic polypeptides. J. Biochem. Biophys. Meth., 45, 157-168. #15846
- 58. Bozo, B., Farkas, J., Toth, G., Wollemann, M., Szucs, M., & Benyhe, S. (2000) Receptor binding and G-protein activation by new Met(5)-enkephalin-Arg(6)-Phe(7) derived peptides. Life Sci., 66, 1241-1251. #14959
- 59. Brask, J., & Jensen, K.J. (2000) Carbopeptides: Chemoselective ligation of peptide aldehydes to an aminooxy-functionalized D-galactose template. J. Peptide Sci., 6, 290-299. #15448
- 60. Braslau, R., Axon, J.R., & Lee, B. (2000) Synthesis of N-hydroxy peptides: Chemical ligation of O-acyl hydroxamic acids. Organic Letters, 2, 1399-1401. #14964
- 61. Buchardt, J., & Meldal, M. (2000) Novel methodology for the solid-phase synthesis of phosphinic peptides. Journal of the Chemical Society-Perkin Transaction 1, 3306-3310. #15852

- 62. Bui, C.T., Ercole, F., Pham, Y., Campbell, R., Rasoul, F.A., Maeji, N.J., & Ede, N.J. (2000) Improving the performance of an acidlabile 4-hydroxymethyl phenoxyacetic acid (HMO) linker on resin and SynPhaseTM grafted solid-supports. J. Peptide Sci., 6, 534-538. #15756
- 63. Bui, C.T., Bray, A.M., Nguyen, T., Ercole, F., Rasoul, F., Sampson, W., & Maeji, N.J. (2000) Acetophenone-based linker for solid-phase peptide synthesis. J. Peptide Sci., 6, 49-56. #14310
- 64. Bui, C.T., Bray, A.M., Nguyen, T., Ercole, F., & Maeji, N.J. (2000) Solid phase synthesis of C-terminal peptide amides: Development of a new aminoethyl-polystyrene linker on the MultipinTM solid support. J. Peptide Sci., 6, 243-250. #14408
- 65. Burgess, K. (2000) Solid-phase organic synthesis. John Wiley & Sons, New York. ISBN 0471318256 #14304
- 66. Calcagni, A., Gavuzzo, E., Lucente, G., Mazza, F., Morera, E., Paradisi, M.P., & Rossi, D. (2000) Peptides containing the sulfonamide junction. 2. Structure and conformation of Z-Tau-Pro-D-Phe-NHiPr. Biopolymers, 54, 379-387. #15855
- 67. Camarero, J.A., Adeva, A., & Muir, T.W. (2000) 3-Thiopropionic acid as a highly versatile multidetachable thioester resin linker. Lett. Peptide Sci., 7, 17-21. #15430
- 68. Cammish, L.E., & Kates, S.A. (2000) Instrumentation for automated solid phase peptide synthesis. Fmoc Solid Phase Peptide Synthesis: A Practical ApproachÉ, 222, 277-302. #15857
- 69. Carlson, C.B., & Beal, P.A. (2000) Solid-phase synthesis of acridine-based threading intercalator peptides. Bioorg. Med. Chem. Lett., 10, 1979-1982. #15764
- 70. Carreno, C., Mendez, M.E., Kim, Y.D., Kim, H.J., Kates, S.A., Andreu, D., & Albericio, F. (2000) Nsc and Fmoc Na-amino protection for solid-phase peptide synthesis: A parallel study. J. Pept. Res., 56, 63-69. #15443
- 71. Cavero, M., Hobbs, A., Madge, D., Motherwell, W.B., Selwood, D., & Potier, P. (2000) Synthesis and biological evaluation of enantiopure thionitrites: The solid-phase synthesis and nitrosation of D-glutathione as a molecular probe. Bioorg. Med. Chem. Lett., 10, 641-644. #14632
- 72. Cerovsky, V., & Bordusa, F. (2000) Protease-catalyzed fragment condensation via substrate mimetic strategy: A useful combination of solid-phase peptide synthesis with enzymatic methods. J. Pept. Res., 55, 325-329. #14363
- 73. Chetyrkina, S., Estieu-Gionnet, K., Lain, G., Bayle, M., & Deleris, G. (2000) Synthesis of N-Fmoc-4-[(diethylphosphono)-2',2'-difluoro-1'-hydroxyethyl]phenylalanine, a novel phosphotyrosyl mimic for the preparation of signal transduction inhibitory peptides. Tetrahedron Lett., 41, 1923-1926. #14347
- 74. Cho, J.K., Park, B.D., & Lee, Y.S. (2000) A novel core-shell type polymer support for solid-phase peptide synthesis. Tetrahedron Lett., 41, 7481-7485. #15861
- 75. Cilli, E.M., Jubilut, G.N., Ribeiro, S.C.F., Oliveira, E., & Nakaie, C.R. (2000) Importance of the solvation degree of peptide-resin beads for amine groups determination by the picric acid method. J. Braz. Chem. Soc., 11, 474-478. #15862
- 76. Clippingdale, A.B., Barrow, C.J., & Wade, J.D. (2000) Peptide thioester preparation by Fmoc solid phase peptide synthesis for use in native chemical ligation. J. Peptide Sci., 6, 225-234. #14405
- 77. Conde-Frieboes, K., Andersen, S., & Breinholt, J. (2000) Synthesis of symmetrical dimeric N,N'-linked peptides on solid support by olefin metathesis. Tetrahedron Lett., 41, 9153-9156. #16159
- 78. Cotton, G.J., & Muir, T.W. (2000) Generation of a dual-labeled fluorescence biosensor for Crk-II phosphorylation using solid-phase expressed protein ligation. Chem. Biol., 7, 253-261. #15864
- 79. Cruz, L.J., Quintana, D., Iglesias, E., Garcia, Y., Huerta, V., Garay, H.E., Duarte, C., & Reyes, O. (2000) Immunogenicity comparison of a multi-antigenic peptide bearing V3 sequences of the human immunodeficiency virus type 1 with TAB9 protein in mice. J. Peptide Sci., 6, 217-224. #14402
- 80. Cudic, M., Wade, J.D., & Otvos, L.Jr. (2000) Convenient synthesis of a head-to-tail cyclic peptide containing an expanded ring. Tetrahedron Lett., 41, 4527-4531. #14594

- 81. Dai, X., Su, Z., & Liu, J.O. (2000) An improved synthesis of a selective aVb3-integrin antagonist cyclo(-RGDfK-). Tetrahedron Lett., 41, 6295-6298. #15531
- 82. Dal, X.D., Su, Z., & Liu, J.O. (2000) An improved synthesis of a selective alpha(v)beta(3)-integrin antagonist cyclo(-RGDfK-). Tetrahedron Lett., 41, 6295-6298. #15868
- 83. Dankwardt, S.M., Billedeau, R.J., Lawley, L.K., Abbot, S.C., Martin, R.L., Chan, C.S., Van Wart, H.E., & Walker, K.A.M. (2000) Solid-phase synthesis of di- and tripeptidic hydroxamic acids as inhibitors of procollagen C-proteinase. Bioorg. Med. Chem. Lett., 10, 2513-2516. #15870
- 84. Denis, B., & Trifilieff, E. (2000) Synthesis of palmitoyl-thioester T-cell epitopes of myelin proteolipid protein (PLP). Comparison of two thiol protecting groups (StBu and Mmt) for on-resin-acylation. J. Peptide Sci., 6, 372-377. #15450
- 85. Dionyssopoulou, H., Mouzaki, A., Slootstra, J., Puijk, W., Meloen, R., Cordopatis, P., & Sotiropoulou, G. (2000) Synthetic peptides as putative therapeutic agents in transplantation medicine: Application of PEPSCAN to the identification of functional sequences in the extracellular domain of the interleukin-2 receptor beta chain (IL-2R beta). J. Immunol. Method, 241, 83-95. #15882
- 86. Dolence, E.K., Dolence, J.M., & Poulter, C.D. (2000) Solid-phase synthesis of a farnesylated CaaX peptide library: Inhibitors of the Ras CaaX endoprotease. Journal of Combinatorial Chemistry, 2, 522-536. #15605
- 87. Drakopoulou, E., Uray, K., Mezo, G., Price, M.R., Vita, C., & Hudecz, F. (2000) Synthesis and antibody recognition of mucin 1 (MUC1)-a-conotoxin chimera. J. Peptide Sci., 6, 175-185. #14361
- 88. Dufau, I., & Mazarguil, H. (2000) Design of a fluorescent amino acid derivative usable in peptide synthesis. Tetrahedron Lett., 41, 6063-6066. #15517
- 89. Dutta, A.S., Gormley, J.J., Coath, M., Hassall, L., Hayward, C.F., Gellert, P.R., Kittlety, R.S., Alcock, P.J., Ferguson, R., Halterman, T., Jamieson, A., Moors, J.A., Moores, J.M., Rees, A., Wood, L.J., Reilly, C.F., & Haworth, D. (2000) Potent cyclic peptide inhibitors of VLA-4 (a4b1 integrin)-mediated cell adhesion. Discovery of compounds like cyclo(MePhe-Leu-Asp-Val-D-Arg-D-Arg) (ZD7349) compatible with depot formulation. J. Peptide Sci., 6, 398-412. #15453
- Dutta, A.S., Crowther, M., Gormley, J.J., Hassall, L., Hayward, C.F., Gellert, P.R., Kittlety, R.S., Alcock, P.J., Jamieson, A., Moores, J.M., Rees, A., Wood, L.J., Reilly, C.F., & Haworth, D. (2000) Potent cyclic monomeric and dimeric peptide inhibitors of VLA-4 (alpha(4)beta(1) integrin)-mediated cell adhesion based on the Ile-Leu-Asp-Val tetrapeptide. J. Peptide Sci., 6, 321-341. #15892
- 91. Ede, N.J., Eagle, S.N., Wickham, G., Bray, A.M., Warne, B., Shoemaker, K., & Rosenberg, S. (2000) Solid phase synthesis of peptide aldehyde protease inhibitors. Probing the proteolytic sites of hepatitis C virus polyprotein. J. Peptide Sci., 6, 11-18. #13956
- 92. Egleton, R.D., Mitchell, S.A., Huber, J.D., Janders, J., Stropova, D., Polt, R., Yamamura, H.I., Hruby, V.J., & Davis, T.P. (2000) Improved bioavailability to the brain of glycosylated Met-enkephalin analogs. Brain Res., 881, 37-46. #15894
- 93. El-Agnaf, O.M.A., Mahil, D.S., Patel, B.P., & Austen, B.M. (2000) Oligomerization and toxicity of beta-amyloid-42 implicated in Alzheimer's disease. Biochem. Biophys. Res. Commun., 273, 1003-1007. #15895
- 94. Falb, E., Yechezkel, T., Salitra, Y., Gellerman, G., Muller, D. & Gilon, C. (2000) In situ generation of Fmoc amino acid chlorides for extremely difficult couplings to sterically hindered secondary amines in solid-phase peptide synthesis. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 55-57). Kluwer Academic Publisher, Dordrecht. #15694
- 95. Falchi, A., Giacomelli, G., Porcheddu, A., & Taddei, M. (2000) 4-(4,6-Dimethoxy[1,3,5]triazin-2-yl)-4-methyl-morpholinium chloride (DMTMM): A valuable alternative to PyBOP for solid phase peptide synthesis. Synlett, 275-277. #15026
- 96. Fields, G.B., Tam, J.P., & Barany, G. (2000) Peptides for the New Millennium. Kluwer Academic Publisher, Dordrecht. ISBN 0792364457 #15688
- 97. Fiertek, D., Gromowska, M., Andersen, A.R., Hansen, P.H., Majewski, T., & Izdebski, J. (2000) Primary structure of fox (Vulpes vulpes) proinsulin based on sequence studies of pancreatic peptides and cDNA. J. Peptide Sci., 6, 413-419. #15454
- 98. Freund, E., Vitali, F., Linden, A., & Robinson, J.A. (2000) Solid-phase synthesis using (allyloxy)carbonyl(Alloc) chemistry of a putative heptapeptide intermediate in vancomycin biosynthesis containing m-chloro-3-hydroxytyrosine. Helv. Chim. Acta, 83,

2572-2579. #15902

- Fruchart, J.S., Grandjean, C., Bonnet, D., Rommens, C., Gras-Masse, H. & Melnyk, O. (2000) A new tartaric acid-based linker for the synthesis of C-terminal peptide a-oxo-aldehydes. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 104-106). Kluwer Academic Publisher, Dordrecht. #15708
- 100. Fruchart, J.S., Lippens, G., Warrass, R., Seetharaman, C., Dhalluin, C., & Boutillon, C. (2000) The Chemical Shift Index method applied to resin-bound peptides. J. Pept. Res., 56, 346-351. #16286
- 101. Ghosh, A.K., & Fan, E. (2000) A novel method for sequence independent incorporation of activated/protected cysteine in Fmoc solid phase peptide synthesis. Tetrahedron Lett., 41, 165-168. #13884
- 102. Goetz, M., Rusconi, F., Belghazi, M., Schmitter, J.M., & Dufourc, E.J. (2000) Purification of the c-erbB2/neu membrane-spanning segment: A hydrophobic challenge. Journal of Chromatography B, 737, 55-61. #13985
- 103. Goldstein, A.S., & Gelb, M.H. (2000) An alternate preparation of thioester resin linkers for solid-phase synthesis of peptide C-terminal thioacids. Tetrahedron Lett., 41, 2797-2800. #14539
- 104. Gordon, L.M., Lee, K.Y.C., Lipp, M.M., Zasadzinski, J.A., Walther, F.J., Sherman, M.A., & Waring, A.J. (2000) Conformational mapping of the N-terminal segment of surfactant protein B in lipid using 13C-enhanced Fourier transform infrared spectroscopy. J. Pept. Res., 55, 330-347. #14364
- 105. Gottschling, D., Seliger, H., Tarrason, G., Piulats, J., Wiersma, M., & Eritja, R. (2000) Synthesis of peptide nucleic acid-peptide chimeras carrying the c-myc tag-sequence. Lett. Peptide Sci., 7, 35-39. #15433
- 106. Groth, T., Meldal, M. & Bock, K. (2000) Backbone amide protection in solid-phase synthesis of peptide isosters derived from N-terminal g-aldehydes. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 146-147). Kluwer Academic Publisher, Dordrecht. #15712
- 107. Gu, C.G., Tsaprailis, G., Breci, L., & Wysocki, V.H. (2000) Selective gas-phase cleavage at the peptide bond terminal to aspartic acid in fixed-charge derivatives of asp-containing peptides. Anal. Chem., 72, 5804-5813. #16294
- 108. Guichard, G., Semetey, V., Didierjean, C., Aubry, A., Rodriguez, M. & Briand, J.P. (2000) An efficient preparation of Osuccinimidyl carbamate derivatives from N-protected b-amino acids: Application to the synthesis of urea containing pseudopeptides and oligoureas. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 148-149). Kluwer Academic Publisher, Dordrecht. #15713
- 109. Guillaumie, F., Kelly, N.M., Kappel, J.C., Barany, G. & Jensen, K.J. (2000) Solid-phase synthesis of peptide aldehydes by a Backbone Amide Linker (BAL) strategy. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 100-101). Kluwer Academic Publisher, Dordrecht. #15706
- 110. Guillaumie, F., Kappel, J.C., Kelly, N.M., Barany, G., & Jensen, K.J. (2000) Solid-phase synthesis of C-terminal peptide aldehydes from amino acetals anchored to a backbone amide linker (BAL) handle. Tetrahedron Lett., 41, 6131-6135. #15518
- 111. Gururaja, T.L., Narasimhamurthy, S., Payan, D.G., & Anderson, D.C. (2000) A novel artificial loop scaffold for the noncovalent constraint of peptides. Chem. Biol., 7, 515-527. #15921
- 112. Hargittai, B., Annis, I., & Barany, G. (2000) Application of solid-phase Ellman's reagent for preparation of disulfide-paired isomers of a-conotoxin SI. Lett. Peptide Sci., 7, 47-52. #15434
- 113. Hargittai, B., Annis, I. & Barany, G. (2000) Application of solid-phase Ellman's reagent for preparation of disulfide-paired isomers of a-conotoxin SI. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 94-95). Kluwer Academic Publisher, Dordrecht. #15703
- 114. Harris, J.L., Backes, B.J., Leonetti, F., Mahrus, S., Ellman, J.A., & Craik, C.S. (2000) Rapid and general profiling of protease specificity by using combinatorial fluorogenic substrate libraries. Proc. Natl. Acad. Sci. USA, 97, 7754-7759. #15073
- Heinonen, P., Rosenberg, J., & Lonnberg, H. (2000) Synthesis of achiral alpha, alpha-bis(aminomethyl)-beta-alanines and their use in the preparation of branched beta-peptide conjugates of N-2-alkyl-1,2,3,4-tetrahydroisoquinolines on solid support. Eur. J. Org. Chem., 3647-3652. #15926

- Hernandez, J.F., Gagnon, J., Chiche, L., Nguyen, T.M., Andrieu, J.P., Heitz, A., Hong, T.T., Pham, T.T.C., & Nguyen, D.L. (2000)
   Squash trypsin inhibitors from Momordica cochinchinensis exhibit an atypical macrocyclic structure. Biochemistry, 39, 5722-5730. #15083
- 117. Hernandez, M., Pozo, L., Gomez, I., & Melchor, A. (2000) Chimeric synthetic peptide as antigen for immunodiagnosis of HIV-1 infection. Biochem. Biophys. Res. Commun., 272, 259-262. #15084
- 118. Hilpert, K., Hansen, G., Wessner, H., Schneider-Mergener, J., & Hohne, W. (2000) Characterizing and optimizing protease/peptide inhibitor interactions, a new application for spot synthesis. J. Biochem. Tokyo, 128, 1051-1057. #16299
- 119. Hlavacek, J., Marcova, R., Budesinsky, M., & Slaninova, J. (2000) 1-Deamino-1(15)-carba and -dicarba analogues of endothelin-1. Collect. Czech. Chem. Commun., 65, 407-424. #15088
- 120. Holm, A., Jorgensen, R.M., Ostergaard, S., & Theisen, M. (2000) Ligand-presenting assembly: A method for C- and N-terminal antigen presentation. J. Pept. Res., 56, 105-113. #15447
- 121. Howe, J., Quibell, M., & Johnson, T. (2000) A new generation of reversible backbone-amide protection for the solid phase synthesis of difficult sequences. Tetrahedron Lett., 41, 3997-4001. #14563
- 122. Hu, B.H., & Messersmith, P.B. (2000) Protection of 3,4-dihydroxyphenylalanine (DOPA) for Fmoc solid-phase peptide synthesis. Tetrahedron Lett., 41, 5795-5798. #15423
- 123. Ishii, A., Hojo, H., Kobayashi, A., Nakamura, K., Nakahara, Y., & Ito, Y. (2000) A facile silyl linker strategy for the solid-phase synthesis of protected glycopeptide: Synthesis of an N-terminal fragment of IL-2 (1-10). Tetrahedron, 56, 6235-6243. #15773
- 124. Ius, A., Bacigalupo, M.A., Longhi, R., & Meroni, G. (2000) Selectively conjugated melittins for liposome time-resolved fluoroimmunoassay of theophylline in serum. Fresenius J. Anal. Chem., 366, 869-872. #15112
- 125. Jensen, K.J., & Barany, G. (2000) Carbopeptides: Carbohydrates as potential templates for de novo design of protein models. J. Pept. Res., 56, 3-11. #15119
- 126. Jiaang, W.T., Tseng, P.H., & Chen, S.T. (2000) Facile solid phase synthesis of YEE(ah-GalNAc)(3), a ligand with known high affinity for the asialoglycoprotein receptor. Synlett, 797-800. #15123
- 127. Jiaang, W.T., Chang, M.Y., Tseng, P.H., & Chen, S.T. (2000) A concise synthesis of the O-glycosylated amino acid building block; using phenyl selenoglycoside as a glycosyl donor. Tetrahedron Lett., 41, 3127-3130. #15122
- 128. Johansson, A., Akerblom, E., Ersmark, K., Lindeberg, G., & Hallberg, A. (2000) An improved procedure for N- to C-directed (inverse) solid-phase peptide synthesis. Journal of Combinatorial Chemistry, 2, 496-507. #15602
- 129. Jones, R.C.F., & Dickson, J. (2000) An amidazoline pseudodipeptide suitable for solid phase peptide synthesis. J. Peptide Sci., 6, 621-624. #16153
- 130. Karlstrom, A., Rosenthal, K., & Unden, A. (2000) Study of the alkylation propensity of cations generated by acidolytic cleavage of protecting groups in Boc chemistry. J. Pept. Res., 55, 36-40. #14033
- 131. Kempe, M. (2000) Oxytocin receptor mimetics prepared by molecular imprinting. Lett. Peptide Sci., 7, 27-33. #15432
- 132. Kennedy, K.J., Lundquist, J.T.IV., Simandan, T.L., Kokko, K.P., Beeson, C.C., & Dix, T.A. (2000) Design rationale, synthesis, and chararacterization of non-natural analogs of the cationic amino acids arginine and lysine. J. Pept. Res., 55, 348-358. #14365
- 133. Kim, H.J. & Samukov, V.V. (2000) Na-2-(4-Nitrophenylsulfonyl)ethoxycarbonyl (Nsc) as an amino-protecting group and its application for peptide synthesis. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 50-52). Kluwer Academic Publisher, Dordrecht. #15692
- 134. Klugmann, K., Karawajczyk, B., Kunikowska, D., Glosnicka, R., & Mackiewicz, Z. (2000) Synthesis of hepatitis C virus protein fragments and evaluation of their immunogenicity. Part II. Polish J. Chem., 74, 1085-1090. #15955
- 135. Koeller, K.M., Smith, M.E.B., & Wong, C.H. (2000) Tyrosine sulfation on a PSGL-1 glycopeptide influences the reactivity of glycosyltransferases responsible for synthesis of the attached O-glycan. J. Amer. Chem. Soc., 122, 742-743. #15146

- 136. Koeller, K.M., Smith, M.E.B., & Wong, C.H. (2000) Chemoenzymatic synthesis of PSGL-1 glycopeptides: Sulfation on tyrosine affects glycosyltransferase-catalyzed synthesis of the O-glycan. Bioorg. Med. Chem., 8, 1017-1025. #15145
- 137. Komba, S., Werdelin, O., Jensen, T., & Meldal, M. (2000) Synthesis of tumor associated sialyl-T-glycopeptides and their immunogenicity. J. Peptide Sci., 6, 585-593. #16150
- 138. Korol'kov, V.I., Platonova, G.A., Azanova, V.V., Tennikova, T.B., & Vlasov, G.P. (2000) In situ preparation of peptidylated polymers as ready-to-use adsorbents for rapid immunoaffinity chromatography. Lett. Peptide Sci., 7, 53-61. #15435
- 139. Kudryavtseva, E.V., Sidorova, M.V., Ovchinnikov, M.V., & Bespalova, Z.D. (2000) Hydrogen peroxide for disulfide bridge formation in methionine-containing peptides. J. Peptide Sci., 6, 208-216. #14403
- 140. Kumar, K.S., Pillai, V.N.R., & Das, M.R. (2000) Syntheses of four peptides from the immunodominant region of hepatitis C viral pathogens using PS-TTEGDA support for the investigation of HCV infection in human blood. J. Pept. Res., 56, 88-96. #15446
- 141. Laczko, I., Vas, E., Toth, G.K., Marczinovits, I., Kiss, M., Husz, S., & Molnar, J. (2000) Conformational consequences of coupling bullous pemphigoid antigenic peptides to glutathione-S-transferase and their diagnostic significance. J. Peptide Sci., 6, 378-386. #15451
- 142. Lauer-Fields, J.L., Tuzinski, K.A., Shimokawa, K., Nagase, H., & Fields, G.B. (2000) Hydrolysis of triple-helical collagen peptide models by matrix metalloproteinases. Journal of Biological Chemistry, 275, 13282-13290. #15157
- 143. Lawrence, S.A. (2000) Hydrazine carboxylate t-butylester (BOC-hydrazide): Its properties and applications. Chim. Oggi, 18, 20-24. #14573
- 144. Lebl, M., Ma, J., Pires, J., Dooley, C. & Houghten, R.A. (2000) Rapid parallel synthesis of 584 betides, peptides composed largely of beta-amino acids with side-chains not found in natural peptides. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 174-175). Kluwer Academic Publisher, Dordrecht. #15719
- 145. Lebl, M. (2000) New technique for high-throughput synthesis of peptides, peptidomimetics and nonpeptide small organic molecule arrrays. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 164-166). Kluwer Academic Publisher, Dordrecht. #15715
- 146. Lee, Y.S., Lee, H.J., Lee, D. & Kim, H.J. (2000) Nsc- and Fmoc-amino acids for automated solid-phase peptide synthesis: Comparative study of the stability in various conditions. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 53-54). Kluwer Academic Publisher, Dordrecht. #15693
- 147. Leelasvatanakij, L., & Aldrich, J.V. (2000) A solid-phase synthetic strategy for the preparation of peptide-based affinity labels: Synthesis of dynorphin A analogs. J. Pept. Res., 56, 80-87. #15445
- 148. Li, P., & Xu, J.C. (2000) HOBt and HOAt-derived immonium salts: New and highly efficient coupling reagents for peptide synthesis. Tetrahedron Lett., 41, 721-724. #14229
- 149. Li, P., & Xu, J.C. (2000) (1H-Benzotriazol-1-yloxy)-N,N-dimethylmethaniminium hexachloroantimonate (BOMI), a novel coupling reagent for solution and solid-phase peptide synthesis. J. Pept. Res., 55, 110-119. #14265
- 150. Li, P., & Xu, J.C. (2000) Total synthesis of cyclosporin O both in solution and in the solid phase using novel thiazolium-, immonium-, and pyridinium-type coupling reagents: BEMT, BDMP, and BEP. J. Org. Chem., 65, 2951-2958. #15777
- 151. Li, Q., Moutiez, M., Charbonnier, J.B., Vaudry, K., Menez, A., Quemeneur, E., & Dugave, C. (2000) Design of a Gag pentapeptide analogue that binds human cyclophilin A more efficiently than the entire capsid protein: New insights for the development of novel anti-HIV-1 drugs. J. Med. Chem., 43, 1770-1779. #15171
- 152. Li, W.R., Lin, S.T., & Yang, J.H. (2000) An expedient N-terminal attachment methodology for the solid phase peptide synthesis. Synlett, 1608-1612. #15972
- 153. Li, X., Zhang, L., Hall, S.E., & Tam, J.P. (2000) A new ligation method for N-terminal tryptophan-containing peptides using the Pictet-Spengler reaction. Tetrahedron Lett., 41, 4069-4073. #14591
- 154. Licha, K., Bhargava, S., Rheinlander, C., Becker, A., Schneider-Mergener, J., & Volkmer-Engert, R. (2000) Highly parallel nano-

- synthesis of cleavable peptide-dye conjugates on cellulose membranes. Tetrahedron Lett., 41, 1711-1715. #14337
- 155. Liu, C.F., Rao, C. & Tam, J.P. (2000) Use of orthogonal ligation methods for the synthesis of a hetero peptide dendrimer. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 118-119). Kluwer Academic Publisher, Dordrecht. #15711
- 156. Liu, G., Fan, Y., Zhao, D., Zhao, Z.G. & Lam, K.S. (2000) Development of novel peptide antibiotics for vancomycin resistant infection using the "one-bead, one-compound" combinatorial library method. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 736-737). Kluwer Academic Publisher, Dordrecht. #15748
- 157. Loidl, G., Musiol, H.J., Groll, M., Huber, R., & Moroder, L. (2000) Synthesis of bivalent inhibitors of eucaryotic proteasomes. J. Peptide Sci., 6, 36-46. #14062
- 158. Malkar, N.B., Lauer-Fields, J.L., & Fields, G.B. (2000) Convenient synthesis of glycosylated hydroxylysine derivatives for use in solid-phase peptide synthesis. Tetrahedron Lett., 41, 1137-1140. #14267
- 159. Malkinson, J.P., Falconer, R.A., & Toth, I. (2000) Synthesis of C-terminal glycopeptides from resin-bound glycosyl azides via a modified Staudinger reaction. J. Org. Chem., 65, 5249-5252. #15779
- 160. Marastoni, M., Bazzaro, M., Gavioli, R., Micheletti, F., Traniello, S., & Tomatis, R. (2000) Design of dimeric peptides obtained from a subdominant Epstein-Barr virus LMP2-derived epitope. Eur. J. Med. Chem., 35, 593-598. #15780
- 161. Mathieu, M.N., Wade, J.D., Tan, Y.Y., Summers, R.J., & Tregear, G.W. (2000) Novel strategy for the synthesis of template-assembled analogues of rat relaxin. J. Peptide Sci., 6, 235-242. #14407
- 162. Matsumura, S., Sakamoto, S., Ueno, A., & Mihara, H. (2000) Construction of alpha-helix peptides with beta-cyclodextrin and dansyl units and their conformational and molecular sensing properties. Chem. Eur. J., 6, 1781-1788. #15200
- 163. Mayo, K.H. (2000) Recent advances in the design and construction of synthetic peptides: For the love of basics or just for the technology of it. Trends Biotech., 18, 212-217. #15203
- 164. McBride, J.D., Freeman, H.N., & Leatherbarrow, R.J. (2000) Identification of chymotrypsin inhibitors from a second-generation template assisted combinatorial peptide library. J. Peptide Sci., 6, 446-452. #15611
- McMurray, J.S., Khabashesku, O., Birtwistle, J.S., & Wang, W. (2000) Convenient preparation of 4-(tetrazol-5-yl)-phenylalanine for use in Fmoc-based solid-phase peptide synthesis. Tetrahedron Lett., 41, 6555-6558. #15524
- 166. McNamara, J.F., Lombardo, H., Pillai, S.K., Jensen, I., Albericio, F., & Kates, S.A. (2000) An efficient solid-phase strategy for the construction of chemokines. J. Peptide Sci., 6, 512-518. #15755
- 167. Melnyk, O., Fehrentz, J.A., Martinez, J., & Gras-Masse, H. (2000) Functionalization of peptides and proteins by aldehyde or keto groups. Biopolymers (Pept. Sci. ), 55, 165-186. #16436
- 168. Mergler, M., Dick, F., Gosteli, J., & Nyfeler, R. (2000) Protected peptide p-nitroanilides by solid-phase synthesis. Lett. Peptide Sci., 7, 1-7. #15429
- Merrifield, B., Barany, G., Deber, C.M., Goodman, M., Hodges, R.S., Hruby, V.J., Muir, T.W., Offord, R., Spatola, A.F., Veber, D.F. & Fields, G.B. (2000) Perspectives for the new peptide millennium. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 797-804). Kluwer Academic Publisher, Dordrecht. #15754
- 170. Meutermans, W.D.F., Golding, S.W., Bourne, G.T., Miranda, L.P., Dooley, M.J., Alewood, P.F. & Smythe, M.L. (2000) Synthesis of small cyclic peptides: An auxiliary approach to address the "difficult cyclization" problem. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 183-185). Kluwer Academic Publisher, Dordrecht. #15723
- 171. Miao, Z.W., & Tam, J.P. (2000) Dehydropeptides from orthogonal ligation of unprotected peptides. Organic Letters, 2, 3711-3713.
- 172. Micklatcher, C.L., Lutgring, R.A. & Chmielewski, J. (2000) Synthetic steps toward de novo designed catalytic five helix bundle proteins. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 496-497). Kluwer Academic Publisher, Dordrecht. #15742

- 173. Miranda, L.P., & Alewood, P.F. (2000) Challenges for protein chemical synthesis in the 21st century: Bringing genomics and proteomics. Biopolymers (Pept. Sci.), 55, 217-226. #16439
- 174. Miranda, L.P., Meutermans, W.D.F., Smythe, M.L., & Alewood, P.F. (2000) Activated O -> N acyl transfer auxiliary: Efficient amide-backbone substitution of hindered "difficult" peptides. J. Org. Chem., 65, 5460-5468. #16006
- 175. Monnee, M.C.F., Marijne, M.F., Brouwer, A.J., & Liskamp, R.M.J. (2000) A practical solid phase synthesis of oligopeptidosulfonamide foldamers. Tetrahedron Lett., 41, 7991-7995. #15640
- 176. Moraes, C.M., Bemquerer, M.P., & Miranda, M.T.M. (2000) Solvolysis and aminolysis on peptidyl-Kaiser oxime resin assisted by Ca2+ and Eu3+: A mild procedure to prepare a-methyl and -ethyl esters of protected peptides. J. Pept. Res., 55, 279-288. #14648
- 177. Nishiuchi, Y., Nishio, H., Inui, T., Bodi, J., & Kimura, T. (2000) Combined solid-phase and solution approach for the synthesis of large peptides or proteins. J. Peptide Sci., 6, 84-93. #14311
- 178. Nishiyama, Y., Shikama, S., Morita, K., & Kurita, K. (2000) Cyclohexyl ether as a new hydroxy-protecting group for serine and threonine in peptide synthesis. Journal of the Chemical Society-Perkin Transaction 1, 1949-1954. #779
- 179. Noda, M. (2000) 2-Methoxy-10,11-dihydro-5H-dibenzo[a,d]cycloheplen-5-one: Application for the peptide amide linkers and guanidino protection of arginine during Fmoc solid-phase peptide synthesis. Amer. Lab., 32, 44. #15222
- 180. Nowick, J.S., Chung, D.M., Maitra, K., Maitra, S., Stigers, K.D., & Sun, Y. (2000) An unnatural amino acid that mimics a tripeptide beta-strand and forms beta-sheetlike hydrogen-bonded dimers. J. Amer. Chem. Soc., 122, 7654-7661. #16028
- 181. O'Donnell, M.J., Drew, M.D., Pottorf, R.S., & Scott, W.L. (2000) UPS on Weinreb resin: A facile solid-phase route to aldehyde and ketone derivatives of "unnatural" amino acids and peptides. Journal of Combinatorial Chemistry, 2, 172-181. #14381
- 182. Okayama, T., Burritt, A., & Hruby, V.J. (2000) 4-Alkoxy-2-hydroxybenzaldehyde (AHB): A versatile aldehyde linker for solid-phase synthesis of C-terminal modified peptides and peptidomimetics. Organic Letters, 2, 1787-1790. #15228
- 183. Osapay, K., Tran, D., Ladokhin, A.S., White, S.H., Henschen, A.H., & Selsted, M.E. (2000) Formation and characterization of a single Trp-Trp cross-link in indolicidin that confers protease stability without altering antimicrobial activity. Journal of Biological Chemistry, 275, 12017-12022. #15232
- 184. Otaka, A. (2000) Development of new deprotecting methodologies for peptides and application to studies on signaling mechanism. Yakugaku Zasshi Journal of the Pharmaceutical Society of Japan, 120, 54-67. #14096
- Otaka, A., Mitsuyama, E., Kinoshita, T., Tamamura, H., & Fujii, N. (2000) Stereoselective synthesis of CF2-substituted phosphothreonine mimetics and their incorporation into peptides using newly developed deprotection procedures. J. Org. Chem., 65, 4888-4899. #16032
- 186. Otvos, L.Jr., Bokonyi, K., Pease, A.M., Giles-Davis, W., Rogers, M.E., Hintz, P.A., Hoffmann, R. & Ertl, H.C.J. (2000) In situ identification of T helper cell epitopes from a cellulose-bound peptide array. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 685-686). Kluwer Academic Publisher, Dordrecht. #15746
- 187. Pellois, J.P., Wang, W., & Gao, X. (2000) Peptide synthesis based on t-Boc chemistry and solution photogenerated acids. Journal of Combinatorial Chemistry, 2, 355-360. #14688
- 188. Percot, A., Zhu, X.X., & Lafleur, M. (2000) Immobilization of lipid vesicles on polymer support via an amphiphilic peptidic anchor: Application to a membrane enzyme. Bioconjug. Chem., 11, 674-678. #16043
- Perodin, J., Bouley, R., Escher, E., Assimomytis, N., Magafa, V., Manessi-Zoupa, E., Theodoropoulos, D., & Cordopatis, P. (2000)
   Angiotensin II analogues with sulphur-containing side-chains in position 5. A structure-activity relationship study. Arzneim. Forsch., 50, 526-529. #15247
- 190. Pfeifer, M.E., Moehle, K., Linden, A., & Robinson, J.A. (2000) Synthesis and solution conformation of beta-hairpin mimetics utilizing a template derived from (2S,3R,4R)-diaminoproline. Helv. Chim. Acta, 83, 444-464. #15251
- 191. Planas, M., Bardaji, E., & Barany, G. (2000) Synthesis of cyclic peptide hybrids with amino acid and nucleobase side-chains. Tetrahedron Lett., 41, 4097-4100. #14592

- 192. Planas, M., Bardaji, E. & Barany, G. (2000) Chemical synthesis of cyclic peptide nucleic acid-peptide hybrids. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 786-787). Kluwer Academic Publisher, Dordrecht. #15753
- 193. Pons, M., Albericio, F., Royo, M., & Giralt, E. (2000) Disulfide bonded cyclic peptide dimers and trimers: An easy entry to high symmetry peptide frameworks. Synlett, 172-181. #15255
- 194. Priem, G., Rocheblave, L., De Michelis, C., Courcambeck, J., & Kraus, J.L. (2000) Synthesis and chemical reactivity of thiophenoxyphenylalanine bioisosteres, suitable synthons for the design of HIV protease inhibitors. J. Chem. Soc. Perkin Trans. 1, 819-824. #15258
- 195. Rabanal, F., Pastor, J.J., Nicolas, E., Albericio, F., & Giralt, E. (2000) Synthesis of aspartimide-free protected peptides on base-labile functionalized resins. Tetrahedron Lett., 41, 8093-8096. #15625
- 196. Rana, S., Dubuc, J., Bradley, M., & White, P. (2000) The synthesis of the cyclic antibacterial peptide Kawaguchipeptin B on 11 different DVB cross-linked PS resins. Tetrahedron Lett., 41, 5135-5139. #14606
- 197. Rasoul, F., Ercole, F., Pham, Y., Bui, C.T., Wu, Z., James, S.N., Trainor, R.W., Wickham, G., & Maeji, N.J. (2000) Grafted supports in solid-phase synthesis. Biopolymers (Pept. Sci.), 55, 207-216. #16438
- 198. Rinnova, M., Novakova, M., Kasicka, V., & Jiracek, J. (2000) Side reactions during photochemical cleavage of an amethyl-6-nitroveratryl-based photolabile linker. J. Peptide Sci., 6, 355-365. #15449
- 199. Rivier, J.E., Jiang, G.C., Struthers, R.S., Koerber, S.C., Porter, J., Cervini, L.A., Kirby, D.A., Craig, A.G., & Rivier, C.L. (2000)
  Design of potent dicyclic (1-5/4-10) gonadotropin releasing hormone (GnRH) antagonists. J. Med. Chem., 43, 807-818. #15272
- Rivier, J.E., Struthers, R.S., Porter, J., Lahrichi, S.L., Jiang, G.C., Cervini, L.A., Ibea, M., Kirby, D.A., Koerber, S.C., & Rivier, C.L. (2000) Design of potent dicyclic (4-10/5-8) gonadotropin releasing hormone (GnRH) antagonists. J. Med. Chem., 43, 784-796. #15270
- Rivier, J.E., Porter, J., Cervini, L.A., Lahrichi, S.L., Kirby, D.A., Struthers, R.S., Koerber, S.C., & Rivier, C.L. (2000) Design of monocyclic (1-3) and dicyclic (1-3/4-10) gonadotropin releasing hormone (GnRH) antagonists. J. Med. Chem., 43, 797-806. #15271
- 202. Roice, M., Kumar, K.S., & Pillai, V.N.R. (2000) Optimization in peptide synthetic conditions of 1,4-butanediol dimethacrylate cross-linked polystyrene resin and its efficiency in solid phase peptide synthesis. Tetrahedron, 56, 3725-3734. #15277
- 203. Rubina, A.Y., Bespalova, Z.D., & Bushuev, V.N. (2000) The solid phase synthesis of peptides containing an arginine residue with an unprotected guanidine group [Russian]. Bioorganicheskaia Khimiia, 26, 263-272. #15284
- 204. Sabirov, A.N., Samukov, V.V., Pozdnyakov, P.I. & Kim, H.J. (2000) Nin-4-Nitrobenzenesulfonyl (Nbs): A new protecting group for the indole moiety of tryptophan. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 60-61). Kluwer Academic Publisher, Dordrecht. #15696
- Sakura, N., Kurosawa, K., & Hashimoto, T. (2000) Structure-activity relationships of neuromedin U. IV. Absolute requirement of the arginine residue at position 7 of dog neuromedin U-8 for contractile activity. Chem. Pharm. Bull. Tokyo, 48, 1166-1170. #15782
- 206. Sanderson, J.M., Singh, P., Fishwick, C.W.G., & Findlay, J.B.C. (2000) The synthesis and reactivity of optically pure amino acids bearing side-chain thioamides. Journal of the Chemical Society-Perkin Transaction 1, 3227-3231. #16065
- Sanz-Nebot, V., Benavente, F., & Barbosa, J. (2000) Separation and characterization of multicomponent peptide mixtures by liquid chromatography-electrospray ionization mass spectrometry. Application to crude products of the synthesis of leuprolide. J. Chromatogr., 870, 315-334. #15293
- Sanz-Nebot, V., Benavente, F., Castillo, A., & Barbosa, J. (2000) Liquid chromatography-electrospray mass spectrometry of multicomponent peptide mixtures - Characterization of a mixture from the synthesis of the hormone goserelin. J. Chromatogr., 889, 119-133. #16066
- 209. Satyanarayana, J., Situ, H., Narasimhamurthy, S., Bhayani, N., Bobek, L.A., & Levine, M.J. (2000) Divergent solid-phase synthesis and candidacidal activity of MUC7 D1, a 51-residue histidine-rich N-terminal domain of human salivary mucin MUC7. J. Pept. Res., 56, 275-282. #15790

- 210. Schreiber, J.V., & Seebach, D. (2000) Solid-phase synthesis of a beta-dodecapeptide with seven functionalized side chains and CD-spectroscopic evidence for a dramatic structural switch when going from water to methanol solution. Helv. Chim. Acta, 83, 3139-3152. #16377
- 211. Schumann, C., Seyfarth, L., Greiner, G., & Reissmann, S. (2000) Synthesis of different types of dipeptide building units containing N- or C-terminal arginine for the assembly of backbone cyclic peptides. J. Pept. Res., 55, 428-435. #15303
- Seebach, D., Schreiber, J.V., Abele, S., Daura, X., & van Gunsteren, W.F. (2000) Structure and conformation of beta-oligopeptide derivatives with simple proteinogenic side chains: Circular dichroism and molecular dynamics investigations. Helv. Chim. Acta, 83, 34-57. #15307
- 213. Seitz, O. (2000) Solid-phase synthesis of doubly labeled peptide nucleic acids as probes for the real-time detection of hybridization. Angew. Chem. Int. Ed., 39, 3249. #16070
- 214. Shan, D.X., Zheng, A.L., Ballard, C.E., Wang, W., Borchardt, R.T., & Wang, B.H. (2000) A facilitated cyclic ether formation and its potential application in solid-phase peptide and organic synthesis. Chem. Pharm. Bull. Tokyo, 48, 238-244. #14325
- 215. Sheppeck, J.E., Kar, H., Gosink, L., Wheatley, J.B., Gjerstad, E., Loftus, S.M., Zubiria, A.R., & Janc, J.W. (2000) Synthesis of a statistically exhaustive fluorescent peptide substrate library for profiling protease specificity. Bioorg. Med. Chem. Lett., 10, 2639-2642. #16380
- Shin, I., Lee, M.R., Lee, J., Jung, M., Lee, W., & Yoon, J. (2000) Synthesis of optically active phthaloyl D-aminooxy acids from L-amino acids or L-hydroxy acids as building blocks for the preparation of aminooxy peptides. J. Org. Chem., 65, 7667-7675.
   #16072
- 217. Shogren-Knaak, M.A., McDonnell, K.A., & Imperiali, B. (2000) a-Chloroacetyl capping of peptides: An N-terminal capping strategy suitable for Edman sequencing. Tetrahedron Lett., 41, 827-829. #14260
- 218. Shrimpton, C.N., Abbenante, G., Lew, R.A., & Smith, A.I. (2000) Development and characterization of novel potent and stable inhibitors of endopeptidase EC 3.4.24.15. Biochem. J., 345, 351-356. #14143
- 219. Singh, L., Nakahara, Y., & Ito, Y. (2000) An efficient access to protected disialylated glycohexaosyl threonine present on the leukosialin of activated T-lymphocytes. Carbohyd. Res., 325, 132-142. #15315
- 220. Sjolin, P., & Kihlberg, J. (2000) Deacetylation of N-alpha-methylated glycopeptides reveals that aza-enolates provide protection against beta-elimination of carbohydrates O-linked to serine. Tetrahedron Lett., 41, 4435-4439. #15318
- 221. Sklyarov, L.Y., Sbitneva, I.N., Kopina, N.A., & Sidorovich, I.G. (2000) Amino acid pyridoxyl eaters in peptide synthesis [Russian]. Bioorganicheskaia Khimiia, 26, 273-284. #15319
- 222. Soll, R., & Beck-Sickinger, A.G. (2000) On the synthesis of orexin A: A novel one-step procedure to obtain peptides with two intramolecular disulphide bonds. J. Peptide Sci., 6, 387-397. #15452
- Songster, M.F., Biancalana, S., Cook, R.M., Dick, D.J. & Hudson, D. (2000) Orthogonal methods for the synthesis of multiply labeled peptide probes and substrates. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 111-112). Kluwer Academic Publisher, Dordrecht. #15709
- 224. Spector, S., Wang, M.H., Carp, S.A., Robblee, J., Hendsch, Z.S., Fairman, R., Tidor, B., & Raleigh, D.P. (2000) Rational modification of protein stability by the mutation of charged surface residues. Biochemistry, 39, 872-879. #15322
- 225. Srebalus, C.A., Li, J.W., Marshall, W.S., & Clemmer, D.E. (2000) Determining synthetic failures in combinatorial libraries by hybrid gas-phase separation methods. J. Amer. Soc. Mass Spectrom., 11, 352-355. #15325
- 226. Stetsenko, D.A., & Gait, M.J. (2000) Efficient conjugation of peptides to oligonucleotides by "native ligation". J. Org. Chem., 65, 4900-4908. #16076
- 227. Sutcliffe-Goulden, J.L., O'Doherty, M.J., & Bansal, S.S. (2000) Solid phase synthesis of [F-18]labelled peptides for positron emission tomography. Bioorg. Med. Chem. Lett., 10, 1501-1503. #15330
- 228. Swinnen, D., & Hilvert, D. (2000) Facile, Fmoc-compatible solid-phase synthesis of peptide C-terminal thioesters. Organic Letters,

- 2, 2439-2442. #16085
- 229. Thieriet, N., Alsina, J., Guibe, F. & Albericio, F. (2000) Solid-phase peptide synthesis in the N--->C direction. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 78-79). Kluwer Academic Publisher, Dordrecht. #15699
- 230. Thieriet, N., Guibe, F., & Albericio, F. (2000) Solid-phase peptide synthesis in the reverse (N -> C) direction. Organic Letters, 2, 1815-1817. #15335
- 231. Tong, Y.S., Fobian, Y.M., Wu, M.Y., Boyd, N.D., & Moeller, K.D. (2000) Conformationally constrained substance P analogues: The total synthesis of a constrained peptidomimetic for the Phe(7)-Phe(8) region. J. Org. Chem., 65, 2484-2493. #15341
- 232. Tornoe, C.W., Davis, P., Porreca, F., & Meldal, M. (2000) a-Azido acids for direct use in solid-phase peptide synthesis. J. Peptide Sci., 6, 594-602. #16151
- 233. Tornoe, C.W., Sonke, T., Maes, I., Schoemaker, H.E., & Meldal, M. (2000) Enzymatic and chiral HPLC resolution of alpha-azido acids and amides. Tetrahedron-Asymmetry, 11, 1239-1248. #15342
- 234. Tornoe, C.W., Sengelov, H., & Meldal, M. (2000) Solid-phase synthesis of chemotactic peptides using alpha-azido acids (vol 6, pg 314, 2000). J. Peptide Sci., 6, 539. #16095
- 235. Tornoe, C.W., Sengelov, H., & Meldal, M. (2000) Solid-phase synthesis of chemotactic peptides using a-azido acids. J. Peptide Sci., 6, 314-320. #15554
- 236. Uray, K., Kajtar, J., Vass, E., Price, M.R., Hollosi, M., & Hudecz, F. (2000) Effect of D-amino acid substitution in a mucin 2 epitope on mucin-specific monoclonal antibody recognition. Arch. Biochem. Biophys., 378, 25-32. #15347
- Varady, L., Rajur, S.B., Nicewonger, R.B., Guo, M.J., & Ditto, L. (2000) Fast and quantitative high-performance liquid chromatography method for the determination of 9-fluorenylmethoxycarbonyl release from solid-phase synthesis resins. J. Chromatogr., 869, 171-179. #15350
- 238. Vichier-Guerre, S., Lo-Man, R., Bay, S., Deriaud, E., Nakada, H., Leclerc, C., & Cantacuzene, D. (2000) Short synthetic glycopeptides successfully induce antibody responses to carcinoma-associated Tn antigen. J. Pept. Res., 55, 173-180. #15353
- 239. Volonterio, A., Bravo, P., Moussier, N., & Zanda, M. (2000) Solid-phase synthesis of partially-modified retro and retro-inverso y [NHCH(CF3)]-peptides. Tetrahedron Lett., 41, 6517-6521. #15537
- 240. Wang, H., Miao, Z.W., Lai, L.H., & Xu, X.J. (2000) An efficient procedure for cleavage of T-butyl protected cysteine in solid phase peptide synthesis. Syn. Commun., 30, 727-735. #14675
- 241. Wang, H.S., Su, W., Liu, D.X., Yang, X.W., Da, C.S., & Wang, R. (2000) Preparation of a polymer supported BINOL-Ti catalyst and its application in asymmetric addition of diethylzinc to aldehydes [Chinese]. Chem. J. Chin. Univ., 21, 1524-1526. #16109
- 242. Warrass, R., Wieruszeski, J.M., Boutillon, C., & Lippens, G. (2000) High-resolution magic angle spinning NMR study of resinbound polyalanine peptides. J. Amer. Chem. Soc., 122, 1789-1795. #14676
- 243. Wenschuh, H., Volkmer-Engert, R., Schmidt, M., Schulz, M., Schneider-Mergener, J., & Reineke, U. (2000) Coherent membrane supports for parallel microsynthesis and screening of bioactive peptides. Biopolymers (Pept. Sci. ), 55, 188-206. #16437
- 244. Wiemann, A., Frank, R., & Tegge, W. (2000) Synthesis of suitably protected hydroxymethylene phosphonate- and 'phosphote phosphonate'-analogues of phosphoserine and their incorporation into synthetic peptides. Tetrahedron, 56, 1331-1337. #15370
- 245. Wilhelm, R.R., Srinivasan, A. & Schmidt, M.A. (2000) Evaluation of ivDde as a quasi-orthogonal protecting group for Fmoc solid-phase peptide synthesis. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 58-59). Kluwer Academic Publisher, Dordrecht. #15695
- 246. Wirkus-Romanowska, I., Miecznikowska, H., Zablocka, A., Rybka, K., Fortuna, W., Miedzybrodzki, R., Szymaniec, S., Janusz, M., Lisowski, J., & Kupryszewski, G. (2000) Fragments of a proline-rich polypeptide settled on a hexapeptide carcass constructed of glycine and L-lysine residues. Synthesis and biological properties. Polish J. Chem., 74, 219-225. #15375

- Wirkus-Romanowska, I., Miecznikowska, H., Janusz, M., Szymaniec, S., Fortuna, W., Miedzybrodzki, R., Zablocka, A., Lisowski, J., & Kupryszewski, G. (2000) New analogues of proline-rich protein fragments. Synthesis and their effect on resistance of murine thymocytes to hydrocortisone. Polish J. Chem., 74, 979-984. #15374
- 248. Xie, H., Becker, J.M., Gibbs, R.A., & Naider, F. (2000) Structure, biological activity and membrane partitioning of analogs of the isoprenylated a-factor mating peptide of Saccharomyces cerevisiae. J. Pept. Res., 55, 372-383. #14400
- 249. Yokum, T.S., & Barany, G. (2000) Strategy in solid-phase peptide synthesis. Solid-Phase Synthesis: A Practical Guide, 79-102. #16411
- 250. Yu, Q., Lehrer, R.I. & Tam, J.P. (2000) Design and synthesis of salt-insensitive cyclic a-defensins. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 521-522). Kluwer Academic Publisher, Dordrecht. #15743
- 251. Zakhariev, S., Szekely, Z., Guarnaccia, C., Antcheva, N. & Pongor, S. (2000) A highly effective method for synthesis of Nw-substituted arginines. In G.B. Fields, J.P. Tam & G. Barany (Eds.), Peptides for the New Millennium. (pp. 74-75). Kluwer Academic Publisher, Dordrecht. #15697
- 252. Zamborelli, T.J., Dodson, W.S., Harding, B.J., Zhang, J., Bennett, B.D., Lenz, D.M., Young, Y., Haniu, M., Liu, C.F., Jones, T., & Jarosinski, M.A. (2000) A comparison of folding techniques in the chemical synthesis of the epidermal growth factor-like domain in neu differentiation factor a/b. J. Pept. Res., 55, 359-371. #14399
- 253. Zaoral, M., Blaha, I., Budesinsky, M., Machova, A., & Slaninova, J. (2000) Non-sequential vasopressin peptides. Stereochemistry and biological activity. J. Peptide Sci., 6, 123-129. #14443
- 254. Zhang, S., & Lee, J.P. (2000) Selectively H-2-labeled Glu/Asp: Application to pK(a) measurements in A beta amyloid peptides. J. Pept. Res., 55, 1-6. #14202
- Ahn, I.A., Kim, S.W., & Ro, S. (1999) Solid phase synthesis of azapeptides using an automatic synthesizer. Mol. Diversity, 4, 23-24. #12922
- Akaji, K., Hayashi, Y., Kiso, Y., & Kuriyama, N. (1999) Convergent synthesis of dolastatin 15 by solid phase coupling of an N-methylamino acid. J. Org. Chem., 64, 405-411. #12622
- 257. Al-Obeidi, F., & Ostrem, J.A. (1999) Factor Xa inhibitors. Exp. Opin. Ther. Patents, 9, 931-953. #13469
- 258. Albericio, F., Frieden, A., del Fresno, M., Royo, M., Alsina, J., Jensen, K.J., Kates, S.A. & Barany, G. (1999) Solid-phase synthesis of "head-to-tail" cyclic and C-terminal modified peptides. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 7-10). Mayflower Scientific Limited, Birmingham. #13450
- Angell, Y.M., Han, Y., Albericio, F. & Barany, G. (1999) Minimization of cysteine racemization during stepwise solid-phase peptide synthesis. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 247-248). Mayflower Scientific Limited, Birmingham. #13687
- Annis, I., Chen, L. & Barany, G. (1999) Mild solid-phase reagents for facile formation of intramolecular disulfide bonds in peptides and proteins. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 249-252). Mayflower Scientific Limited, Birmingham. #13688
- Antopolsky, M., & Azhayev, A. (1999) Stepwise solid-phase synthesis of peptide-oligonucleotide conjugates on new solid supports. Helv. Chim. Acta, 82, 2130-2140. #13905
- 262. Appel, J.R., Johnson, J., Narayanan, V.L., & Houghten, R.A. (1999) Identification of novel antitumor agents from mixture-based synthetic combinatorial libraries using cell-based assays. Mol. Diversity, 4, 91-102. #12926
- 263. Appendini, P., & Hotchkiss, J.H. (1999) Antimicrobial activity of a 14-residue peptide against Escherichia coli O157: H7. Journal of Applied Microbiology, 87, 750-756. #13906
- Arrowsmith, J., Missailidis, S., & Stevens, M.F.G. (1999) Antitumour imidazotetrazines. Part 37. Conjugation of the DNA majorgroove alkylating imidazotetrazine mitozolomide to peptide motifs recognizing the minor groove. Anti-Cancer Drug. Des., 14, 205-217. #13908

- Ast, T., Heine, N., Germeroth, L., Schneider-Mergener, J., & Wenschuh, H. (1999) Efficient assembly of peptomers on continuous surfaces. Tetrahedron Lett., 40, 4317-4318. #12928
- Aubagnac, J.L., Enjalbal, C., Drouot, C., Combarieu, R., & Martinez, J. (1999) Imaging time-of-flight secondary ion mass spectrometry of solid-phase peptide syntheses. J. Mass Spectrom., 34, 749-754. #13475
- 267. Ayers, B., Blaschke, U.K., Camarero, J.A., Cotton, G.J., Holford, M., & Muir, T.W. (1999) Introduction of unnatural amino acids into proteins using expressed protein ligation. Biopolymers (Pept. Sci.), 51, 343-354. #14270
- 268. Babu, V.V.S., Gayathri, K., & Gopi, H.N. (1999) Fmoc-peptide acid chlorides in fragment coupling: Synthesis of beta-casomorphin by 3+2 divergent approach. Syn. Commun., 29, 79-91. #12707
- 269. Bajusz, S., & Hudecz, F. (1999) Peptides 1998. Akademiai Kiado, Budapest. #14406
- 270. Balaspiri, L., Janaky, T., Tegyei, Z., Krajcsovits, F., Blazso, G., Mak, M., Takßcs, T. & Kasa, P. (1999) Solid phase synthesis of new galanins and galanin segments: Structure-activity relationships. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 55-58). Mayflower Scientific Limited, Birmingham. #13459
- 271. Barany, G., Angell, Y.M., Annis, I., Chen, L., Gross, C.M. & Hargittai, B. (1999) Current strategies for disulfide bridge formation in synthetic peptides and proteins. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 85-88). Mayflower Scientific Limited, Birmingham. #13465
- 272. Bark, S.J., & Kent, S.B.H. (1999) Engineering an unnatural N-alpha-anchored disulfide into BPTI by total chemical synthesis: Structural and functional consequences. FEBS Lett., 460, 67-76. #13912
- 273. Barlos, K., & Gatos, D. (1999) 9-Fluorenylmethyloxycarbonyl/tbutyl-based convergent protein synthesis. Biopolymers (Pept. Sci.), 51, 266-278. #13894
- 274. Baru, M.B., Mustaeva, L.G., Gorbunova, E.Y., Vagenina, I.V., Kitaeva, M.A., & Cherskii, V.V. (1999) Spectrophotometric monitoring in continuous-flow Boc-based solid-phase peptide synthesis. J. Pept. Res., 54, 263-269. #13751
- 275. Bechinger, B. (1999) The structure, dynamics and orientation of antimicrobial peptides in membranes by multidimensional solid-state NMR spectroscopy. Biochimica et Biophysica Acta Biomembranes, 1462, 157-183. #14935
- 276. Bednarek, M.A., Silva, M.V., Arison, B., MacNeil, T., Kalyani, R.N., Huang, R.R.C., & Weinberg, D.H. (1999) Structure-function studies on the cyclic peptide MT-II, lactam derivative of alpha-melanotropin. Peptides, 20, 401-409. #13478
- 277. Behrendt, R., Renner, C., Schenk, M., Wang, F.Q., Wachtveitl, J., Oesterhelt, D., & Moroder, L. (1999) Photomodulation of the conformation of cyclic peptides with azobenzene moieties in the peptide backbone. Angew. Chem. Int. Ed., 38, 2771-2774. #13916
- 278. Beligere, G.S., & Dawson, P.E. (1999) Synthesis of a three zinc finger protein, Zif268, by native chemical ligation. Biopolymers (Pept. Sci.), 51, 363-369. #14272
- 279. Belvisi, L., Gennari, C., Mielgo, A., Potenza, D., & Scolastico, C. (1999) Conformational preferences of peptides containing reverse-turn mimetic bicyclic lactams: Inverse gamma-turns versus type-II ' beta-turns Insights into beta-hairpin stability. Eur. J. Org. Chem., 389-400. #12709
- 280. Berezhkovskiy, L., Pham, S., Reich, E.P., & Deshpande, S. (1999) Synthesis and kinetics of cyclization of MHC class II-derived cyclic peptide vaccine for diabetes. J. Pept. Res., 54, 112-119. #13480
- 281. Bergeron, R.J., Bussenius, J., Muller, R., McCosar, B.H., & McManis, J.S. (1999) (R)- And (S)-N-(benzyloxycarbonyl)-3,4-epoxybutylamine. New 4-amino-2-bydroxybutyl synthons for the synthesis of hypusine reagents and (R)-6-and (S)-7-hydroxyspermidine. Tetrahedron: Asymmetry, 10, 4285-4294. #14941
- 282. Biederman, K.J., Lee, H., Haney, C.A., Kaczmarek, M., & Buettner, J.A. (1999) Combinatorial peptide on-resin analysis: Optimization of static nanoelectrospray ionization technique for sequence determination. J. Pept. Res., 53, 234-243. #12859
- 283. Birr, C. (1999) Organic synthetic methodology extensions forgotten in peptide science. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 75-84). Mayflower Scientific Limited, Birmingham. #13464

- 284. Bisland, S.K., Singh, D., & Gariepy, J. (1999) Potentiation of chlorin e6 photodynamic activity in vitro with peptide-based intracellular vehicles. Bioconjug. Chem., 10, 982-992. #13922
- 285. Blackburn, C. & Kates, S.A. (1999) Homodetic cyclic peptides via solid phase synthesis. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 129-134). Mayflower Scientific Limited, Birmingham. #13663
- 286. Bofill, J.M., & Albericio, F. (1999) Understanding the structure/reactivity of aminium/uronium salts as coupling reagents in peptide synthesis. Tetrahedron Lett., 40, 2641-2644. #12627
- 287. Bogdanowich-Knipp, S.J., Jois, S.D.S., & Siahaan, T.J. (1999) Effect of conformation on the conversion of cyclo-(1,7)-Gly-Arg-Gly-Asp-Ser-Pro-Asp-Gly-OH to its cyclic imide degradation product. J. Pept. Res., 54, 43-53. #13102
- 288. Boguslavsky, J. (1999) Need protein? Organic chemistry holds the key. Drug. Dis. Develop., 30-33. #12914
- 289. Bonnet, D., Samson, F., Rommens, C., Gras-Masse, H., & Melnyk, O. (1999) Synthesis of hydrazinopeptides using solid-phase N-electrophilic amination: Exctension to the Fmoc/tert-butyl strategy and chemistry of the N-N bond in strong acid media. J. Pept. Res., 54, 270-278. #13752
- Bosze, S., Koczan, G., Csik, G., Falus, A. & Hudecz, F. (1999) A new fluorophore for peptide labelling: Synthesis and analysis of its interaction with interleukin-6 receptor. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 189-192). Mayflower Scientific Limited, Birmingham. #13672
- 291. Boudjabi, S., Dewynter, G., Voyer, N., Toupet, L., & Montero, J.L. (1999) Sulfahydantoins as tripeptide constraints: Synthesis and structure of chiral substituted 3-oxo-1,2,5-thiadiazolidine 1,1-dioxides. Eur. J. Org. Chem., 2275-2283. #14955
- Bourne, G.T., Meutermans, W.D.F., Alewood, P.F., McGeary, R.P., Scanlon, M., Watson, A.A., & Smythe, M.L. (1999) A
  backbone linker for BOC-based peptide synthesis and on-resin cyclization: Synthesis of stylostatin 1. J. Org. Chem., 64, 30953101. #12861
- 293. Bourne, G.T., Meutermans, W.D.F., & Smythe, M.L. (1999) The development of solid phase protocols for a backbone amide linker and its application to the Boc-based assembly of linear peptides. Tetrahedron Lett., 40, 7271-7274. #13774
- 294. Brickmann, K., Yuan, Z.Q., Sethson, I., Somfai, P., & Kihlberg, J. (1999) Synthesis of conformationally restricted mimetics of gamma-turns and incorporation into desmopressin, an analogue of the peptide hormone vasopressin. Chem. Eur. J., 5, 2241-2253. #13485
- 295. Broadbridge, R.J., Sharma, R.P. & Akhtar, M. (1999) Design and synthesis of novel Src homology-2 domain inhibitors. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 211-216). Mayflower Scientific Limited, Birmingham. #13678
- 296. Broddefalk, J., Forsgren, M., Sethson, I., & Kihlberg, J. (1999) Preparation of a diglycosylated hydroxylysine building block used in solid-phase synthesis of a glycopeptide from type II collagen. J. Org. Chem., 64, 8948-8953. #13924
- Buku, A. (1999) Mast cell degranulating (MCD) peptide: A prototypic peptide in allergy and inflammation. Peptides, 20, 415-420.
   #13489
- 298. Cabezas, E., & Satterthwait, A.C. (1999) The hydrogen bond mimic approach: Solid-phase synthesis of a peptide stabilized as an a-helix with a hydrazone link. J. Amer. Chem. Soc., 121, 3862-3875. #12862
- Caciagli, V., Cardinali, F., Sisto, A. & Lombardi, P. (1999) Scaling-up procedure for continuous flow SPPS. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 63-66). Mayflower Scientific Limited, Birmingham. #13461
- 300. Canne, L.E., Botti, P., Simon, R.J., Chen, Y.J., Dennis, E.A., & Kent, S.B.H. (1999) Chemical protein synthesis by solid phase ligation of unprotected peptide segments. J. Amer. Chem. Soc., 121, 8720-8727. #14232
- Ceruso, M.A., McComsey, D.F., Leo, G.C., Andrade-Gordon, P., Addo, M.F., Scarborough, R.M., Oksenberg, D., & Maryanoff, B.E. (1999) Thrombin receptor-activating peptides (TRAPs): Investigation of bioactive conformations via structure-activity, spectroscopic, and computational studies. Bioorg. Med. Chem., 7, 2353-2371. #13926

- 302. Cervini, L., Theobald, P., Corrigan, A., Craig, A.G., Rivier, C., Vale, W., & Rivier, J. (1999) Corticotropin releasing factor (CRF) agonists with reduced amide bonds and Ser(7) substitutions. J. Med. Chem., 42, 761-768. #12720
- 303. Chaloin, L., Mery, J., Van Mau, N., Divita, G., & Heitz, F. (1999) Synthesis of a template-associated peptide designed as a transmembrane ion channel former. J. Peptide Sci., 5, 381-391. #13928
  - 304. Chen, H., Cong, L.N., Li, Y.H., Yao, Z.J., Wu, L., Zhang, Z.Y., Burke, T.R., & Quon, M.J. (1999) Phosphotyrosyl mimetic peptide reverses impairment of insulin-stimulated translocation of GLUT4 caused by overexpression of PTP1B in rat adipose cells. Biochemistry, 38, 384-389. #12722
  - Chen, L., Golser, R., Machova, A., Slaninova, J., & Barany, G. (1999) Chemical syntheses and biological studies on dimeric chimeras of oxytocin and the V-2-antagonist, d(CH2)(5)[D-Ile(2),Ile(4)]arginine vasopressin. J. Med. Chem., 42, 5002-5009. #13932
  - 306. Chitnumsub, P., Fiori, W.R., Lashuel, H.A., Diaz, H., & Kelly, J.W. (1999) The nucleation of monomeric parallel beta-sheet-like structures and their self-assembly in aqueous solution. Bioorg. Med. Chem., 7, 39-59. #12724
  - 307. Chiva, C., Vilaseca, M., Giralt, E., & Albericio, F. (1999) An HPLC-ESMS study on the solid-phase assembly of C-terminal proline peptides. J. Peptide Sci., 5, 131-140. #12725
  - 308. Cilli, E.M., Marchetto, R., Schreier, S., & Nakaie, C.R. (1999) Correlation between the mobility of spin-labeled peptide chains and resin solvation: An approach to optimize the synthesis of aggregating sequences. J. Org. Chem., 64, 9118-9123. #13937
  - Clippingdale, A.B., Macris, M., Wade, J.D., & Barrow, C.J. (1999) Synthesis and secondary structural studies of penta(acetyl-Hmb)A beta(1-40). J. Pept. Res., 53, 665-672. #12945
  - 310. Cornille, F., Wecker, K., Loffet, A., Genet, R., & Roques, B. (1999) Efficient solid-phase synthesis of Vpr from HIV-1 using low quantities of uniformly 13C-, 15N-labeled amino acids for NMR structural studies. J. Pept. Res., 54, 427-435. #13787
  - 311. Cotton, G.J., & Muir, T.W. (1999) Peptide ligation and its application to protein engineering. Chem. Biol., 6, R247-R256. #13939
  - 312. Coupal, M., De Lean, A., McNicoll, N., & Fournier, A. (1999) Development of p-benzoylbenzoylated [N,C,rANP(1-28)]pBNP32 (pBNP1) derivatives and affinity photolabeling of the bovine NPR-A receptor. Biochem. Biophys. Res. Commun., 258, 81-86. #12948
  - 313. Cudic, M., Bulet, P., Hoffmann, R., Craik, D.J., & Otvos, L.Jr. (1999) Chemical synthesis, antibacterial activity and conformation of diptericin, an 82-mer peptide originally isolated from insects. Eur. J. Biochem., 266, 549-558. #14234
  - 314. Daquinag, A.C., Sato, T., Koda, H., Takao, T., Fukuda, M., Shimonishi, Y., & Tsukamoto, T. (1999) A novel endogenous inhibitor of phenoloxidase from Musca domestica has a cystine motif commonly found in snail and spider toxins. Biochemistry, 38, 2179-2188. #12728
  - 315. Davies, M., & Bradley, M. (1999) Internal resin capture: A self purification method for the synthesis of C-terminally modified peptides. Tetrahedron, 55, 4733-4746. #12730
  - 316. Davis, B.G. (1999) Recent developments in glycoconjugates. J. Chem. Soc. Perkin Trans. 1, 3215-3237. #13943
  - 317. Dawson, N.F., Tan, Y.Y., Macris, M., Otvos, L., Summers, R.J., Tregear, G.W., & Wade, J.D. (1999) Solid-phase synthesis of ovine Leydig cell insulin-like peptide: A putative ovine relaxin? J. Pept. Res., 53, 542-547. #12953
  - 318. Dawson, N.F., Craik, D.J., McManus, A.M., Dashper, S.G., Reynolds, E.C., Tregear, G.W., Otvos, L., & Wade, J.D. (1999) Chemical synthesis, characterization and activity of RK-1, a novel alpha-defensin-related peptide. J. Peptide Sci., 6, 19-25. #13945
  - 319. de Bont, D.B.A., Sliedregt-Bol, K.M., Hofmeyer, L.J.F., & Liskamp, R.M.J. (1999) Increased stability of peptidesulfonamide peptidomimetics towards protease catalyzed degradation. Bioorg. Med. Chem., 7, 1043-1047. #13499
  - 320. Debenham, S.D., Cossrow, J., & Toone, E.J. (1999) Synthesis of alpha- and beta-carbon-linked serine analogues of the P-k trisaccharide. J. Org. Chem., 64, 9153-9163. #13950
  - 321. DeGrado, W.F., Schneider, J.P., & Hamuro, Y. (1999) The twists and turns of beta-peptides. J. Pept. Res., 54, 206-217. #13951

- 322. Delarue, S., & Sergheraert, C. (1999) PyBroP: A convenient activator for the synthesis of formamidines. Tetrahedron Lett., 40, 5487-5490. #13122
- 323. Doleckova, L., Pavlik, M., Mares, M. & Kluh, I. (1999) Inhibitory specificity spectrum of peptide a-amylase inhibitors designed by limited combinatorial libraries. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 277-278). Mayflower Scientific Limited, Birmingham. #13693
- 324. Dong, S.L., Wang, T., Chen, Q., & Wang, R. (1999) Synthesis, structure-activity relationship of nociceptin and its fragments. Chin. Sci. Bull., 44, 1655-1659. #15014
- Dragovich, P.S., Zhou, R., Skalitzky, D.J., Fuhrman, S.A., Patick, A.K., Ford, C.E., Meador, J.W., & Worland, S.T. (1999) Solid-phase synthesis of irreversible human rhinovirus 3C protease inhibitors. Part 1: Optimization of tripeptides incorporating N-terminal amides. Bioorg. Med. Chem., 7, 589-598. #12957
- 326. Drozdz, R., Hintermann, E., Tanner, H., Zumsteg, U., & Eberle, A.N. (1999) (D-(p-Benzoylphenylalanine)13,tyrosine19)-melanin-concentrating hormone, a potent analogue for MCH receptor crosslinking. J. Peptide Sci., 5, 234-242. #14430
- 327. Echner, H. & Voelter, W. (1999) The 5-phenyl-5H-dibenzo[a,d]cyclohepten-5-yl (Pdbs) and the 5-anisyl-5H-dibenzo[a,d] cyclohepten-5-yl (Adbs) residues: Two new protecting groups for the Nim-function of histidine. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 279-282). Mayflower Scientific Limited, Birmingham. #13694
- 328. Eggenweiler, H.M., Clausen, N. & Bayer, E. (1999) a-Hydroxymethylacrylic acid in solid phase synthesis: An extremely nucleophile sensitive linker. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 283-286). Mayflower Scientific Limited, Birmingham. #13695
- Eggleston, I.M., Peggion, C., Svendsen, J.S. & Mutter, M. (1999) Practical synthesis and application of a b-turn mimetic for SPPS.
   In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 287-290). Mayflower Scientific Limited, Birmingham. #13696
- 330. Eichler, J. & Houghten, R.A. (1999) Scaffolded peptide combinatorial libraries. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 155-158). Mayflower Scientific Limited, Birmingham. #13667
- 331. Eisele, F., Owen, D.J., & Waldmann, H. (1999) Peptide conjugates as tools for the study of biological signal transduction. Bioorg. Med. Chem., 7, 193-224. #12736
- 332. Elbayed, K., Bourdonneau, M., Furrer, J., Richert, T., Raya, J., Hirschinger, J., & Piotto, M. (1999) Origin of the residual NMR linewidth of a peptide bound to a resin under magic angle spinning. J. Magn. Resonance, 136, 127-129. #12531
- 333. Eleftheriou, S., Gatos, D., Panagopoulos, A., Stathopoulos, S., & Barlos, K. (1999) Attachment of histidine, histamine and urocanic acid to resins of the trityl-type. Tetrahedron Lett., 40, 2825-2828. #12636
- 334. Englebretsen, D.R., & Robillard, G.T. (1999) An N-terminal method for peptide solubilisation. Tetrahedron, 55, 6623-6634. #12959
- Epton, R. (1999) Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. Mayflower Scientific Limited, Birmingham. ISBN 0952701138 #13448
- 336. Evstigneeva, R.P., Zheltukhina, G.A., Khalil', V., & Efimova, E.I. (1999) The synthesis of Arg-containing peptides and their conjugates with protohemin IX and tetraphenylporphyrin. Bioorganicheskaia Khimiia, 25, 572-580. #13960
- 337. Falb, E., Yechezkel, T., Salitra, Y., & Gilon, C. (1999) In situ generation of Fmoc-amino acid chlorides using bis(trichloromethyl) carbonate and its utilization for difficult couplings in solid-phase peptide synthesis. J. Pept. Res., 53, 507-517. #12960
- 338. Fischer, M., & Tran, C.D. (1999) Investigation of solid phase peptide synthesis by the near-infrared multispectral imaging technique: A detection method for combinatorial chemistry. Anal. Chem., 71, 2255-2261. #13402
- 339. Fotsch, C., Kumaravel, G., Sharma, S.K., Wu, A.D., Gounarides, J.S., Nirmala, N.R., & Petter, R.C. (1999) On-resin macrocyclization of peptides via intramolecular SnAr reactions. Bioorg. Med. Chem. Lett., 9, 2125-2130. #13514

- 340. Francke, C., Loyal, R., Ohad, I., & Haehnel, W. (1999) In vitro assembly of a beta(2) cytochrome b(559)-like complex from the chemically synthesised beta-subunit encoded by the Synechocystis sp 6803 psbF gene. FEBS Lett., 442, 75-78. #12354
- 341. Freund, E., & Robinson, J.A. (1999) Solid-phase synthesis of a putative heptapeptide intermediate in vancomycin biosynthesis. Chem. Commun., 2509-2510. #14236
- 342. Fruchart, J.S., Gras-Masse, H., & Melnyk, O. (1999) A new linker for the synthesis of C-terminal peptide alpha-oxo-aldehydes. Tetrahedron Lett., 40, 6225-6228. #13517
- 343. Fukuda, H., Shimizu, T., Nakajima, M., Mori, H., & Shirasawa, T. (1999) Synthesis, aggregation, and neurotoxicity of the Alzheimer's Ab1-42 amyloid peptide and its isoaspartyl isomers. Bioorg. Med. Chem. Lett., 9, 953-956. #12867
- 344. Fukuda, H., Irie, K., Nakahara, A., Ohigashi, H., & Wender, P.A. (1999) Solid-phase synthesis, mass spectrometric analysis of the zinc-folding, and phorbol ester-binding studies of the 116-mer peptide containing the tandem cysteine-rich C1 domains of protein kinase C gamma. Bioorg. Med. Chem., 7, 1213-1221. #13519
- 345. Gademann, K., Hintermann, T., & Schreiber, J.V. (1999) b-Peptides: Twisting and turning. Curr. Med. Chem., 6, 905-925. #13973
- 346. Ganguly, D., & Rodriguez, M.J. (1999) Cyclopeptamine lipopeptide antifungal agents. Exp. Opin. Ther. Patents, 9, 67-71. #12358
- 347. Gennari, C., Mielgo, A., Potenza, D., Scolastico, C., Piarulli, U., & Manzoni, L. (1999) Solid-phase synthesis of peptides containing reverse-turn mimetic bicyclic lactams. Eur. J. Org. Chem., 379-388. #12742
- 348. Gibson, C., Goodman, S.L., Hahn, D., Holzemann, G., & Kessler, H. (1999) Novel solid-phase synthesis of azapeptides and azapeptoides via Fmoc-strategy and its application in the synthesis of RGD-mimetics. J. Org. Chem., 64, 7388-7394. #13981
- 349. Giraud, M., Cavelier, F., & Martinez, J. (1999) A side-reaction in the SPPS of Trp-containing peptides. J. Peptide Sci., 5, 457-461. #13983
- 350. Goetz, M., Schmitter, J.M., Geoffre, S., & Dufourc, E.J. (1999) Chemical synthesis of yeast mitochondrial ATP synthase membranous subunit 8. J. Peptide Sci., 5, 245-250. #13106
- 351. Gomez-Martinez, P., Dessolin, M., Guibe, F., & Albericio, F. (1999) N-a-Alloc temporary protection in solid-phase peptide synthesis. The use of amine-borane complexes as allyl group scavengers. J. Chem. Soc. Perkin Trans. 1, 2871-2874. #13987
- 352. Gomme, P.T., Stanton, P.G., & Hearn, M.T.W. (1999) Evaluation of a pepscan approach to identify epitopes recognised by antihTSH monoclonal antibodies. J. Biochem. Biophys. Meth., 38, 53-70. #12594
- 353. Grahn, S., Kurth, T., Ullmann, D., & Jakubke, H.D. (1999) S' subsite mapping of serine proteases based on fluorescence resonance energy transfer. Biochimica et Biophysica Acta Protein Structure and Molecular Enzymology, 1431, 329-337. #12969
- 354. Gras-Masse, H., Georges, B., Estaquier, J., Tranchand-Bunel, D., Tartar, A., Druilhe, P., & Auriault, C. (1999) Convergent peptide libraries, or mixotopes, to elicit or to identify specific immune responses. Curr. Opin. Immunol., 11, 223-228. #12639
- 355. Guarna, A., Guidi, A., Machetti, F., Menchi, G., Occhiato, E.G., Scarpi, D., Sisi, S., & Trabocchi, A. (1999) Synthesis and reactivity of bicycles derived from tartaric acid and alpha-amino acids: A novel class of conformationally constrained dipeptide isosteres based upon enantiopure 3-aza-6,8-dioxabicyclo[3.2.1]octane-7-carboxylic acid. J. Org. Chem., 64, 7347-7364. #13995
- 356. Gung, B.W., Zou, D., Stalcup, A.M., & Cottrell, C.E. (1999) Characterization of a water-soluble, helical beta-peptide. J. Org. Chem., 64, 2176-2177. #12973
- Habermann, J. & Kunz, H. (1999) Pentafluorophenyluronium reagents and allylic anchoring groups in solid phase glycopeptide synthesis. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 217-220). Mayflower Scientific Limited, Birmingham. #13679
- 358. Hamuro, Y., Scialdone, M.A., & DeGrado, W.F. (1999) Resin-to-resin acyl- and aminoacyl-transfer reactions using oxime supports. J. Amer. Chem. Soc., 121, 1636-1644. #12641
- 359. Hanessian, S., Luo, X.H., & Schaum, R. (1999) Synthesis and folding preferences of gamma-amino acid oligopeptides: Stereochemical control in the formation of a reverse turn and a helix. Tetrahedron Lett., 40, 4925-4929. #13524

- 360. Haramura, M., Tsuzuki, K., Okamachi, A., Yogo, K., Ikuta, M., Kozono, T., Takanashi, H., & Murayama, E. (1999) Structure-activity study of intact porcine motilin. Chem. Pharm. Bull. Tokyo, 47, 1555-1559. #14003
- 361. Hargittai, B., & Barany, G. (1999) Controlled syntheses of natural and disulfide-mispaired regioisomers of alpha-conotoxin SI. J. Pept. Res., 54, 468-479. #14004
- 362. Heizmann, G., Hildebrand, P., Tanner, H., Ketterer, S., Pansky, A., Froidevaux, S., Beglinger, C., & Eberle, A.N. (1999) A combinatorial peptoid library for the identification of novel MSH and GRP/bombesin receptor ligands. Journal of Receptor and Signal Transduction Research, 19, 449-466. #12746
- 363. Henkel, W., Vogl, T., Echner, H., Voelter, W., Urbanke, C., Schleuder, D., & Rauterberg, J. (1999) Synthesis and folding of native collagen III model peptides. Biochemistry, 38, 13610-13622. #14012
- 364. Henklein, P., Ionescu, D., El-Faham, A., Wenschuh, H., Bienert, M., Carpino, L.A. & Beyermann, M. (1999) Coupling of highly hindered systems using protected amino acid chlorides. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 221-224). Mayflower Scientific Limited, Birmingham. #13680
- 365. Henklein, P., Mugge, C., Costisella, B., Wray, V., Domke, T., El-Faham, A., Lee, Y., Kates, S.A. & Carpino, L.A. (1999) Solution structure of uronium and phosphonium salts. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 309-310). Mayflower Scientific Limited, Birmingham. #13700
- 366. Higashimoto, Y., Kodama, H., Jelokhani-Niaraki, M., Kato, F., & Kondo, M. (1999) Structure-function relationship of model Aib-containing peptides as ion transfer intermembrane templates. J. Biochem. Tokyo, 125, 705-712. #12747
- Hintermann, E., Drozdz, R., Tanner, H. & Eberle, A.N. (1999) Synthesis and receptor binding activity of analogues and fragment of human melanin-concentrating hormone (MCH). In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 193-196). Mayflower Scientific Limited, Birmingham. #13673
- 368. Hioki, H., Yamada, T., Fujioka, C., & Kodama, M. (1999) Peptide library based on calix[4]arene. Tetrahedron Lett., 40, 6821-6825. #13761
- 369. Hoffmann, R., Craik, D.J., Bokonyi, K., Varga, I., & Otvos, L. (1999) High level of aspartic acid-bond isomerization during the synthesis of an N-linked tau glycopeptide. J. Peptide Sci., 5, 442-456. #14018
- Hoffmann, R., Ertl, H.C.J., Pease, A.M., Rovera, G. & Otvos, L.Jr. (1999) Optimization of synthesis and DNA hybridization of peptide nucleic acids. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 311-314). Mayflower Scientific Limited, Birmingham. #13701
- 371. Hoogerhout, P., Stittelaar, K.J., Brugghe, H.F., Timmermans, J.A.M., ten Hove, G.J., Jiskoot, W., Hoekman, J.H.G., & Roholl, P.J.M. (1999) Solid-phase synthesis and application of double-fluorescent-labeled lipopeptides, containing a CTL-epitope from the measles fusion protein. J. Pept. Res., 54, 436-443. #13788
- 372. Hsieh, H.P., Wu, Y.T., Chen, S.T., & Wang, K.T. (1999) Direct solid-phase synthesis of octreotide conjugates: Precursors for use as tumor-targeted radiopharmaceuticals. Bioorg. Med. Chem., 7, 1797-1803. #14020
- 373. Huang, H., & Rabenstein, D.L. (1999) A cleavage cocktail for methionine-containing peptides. J. Pept. Res., 53, 548-553. #12981
- 374. Huang, Y.D., Chen, H., Feng, Y.M., & Zhang, J.P. (1999) Preliminary crystallographic studies on two insulin analogues which retain high biological activities after alterations in A chain. Chin. Sci. Bull., 44, 1480-1483. #15101
- 375. Huang, Y.D., Yang, S.Z., Feng, Y.M., & Niu, C.I. (1999) An insulin analog with g-amino butyric acid substitution for A13Leu-A14Tyr. J. Pept. Res., 54, 18-22. #13100
- 376. Husbyn, M., Orning, L., Cuthbertson, A., & Fischer, P.M. (1999) Linear analogues derived from the first EGF-like domain of human blood coagulation factor VII: Enhanced inhibition of FVIIa/TF complex activity by backbone modification through aspartimide formation. J. Peptide Sci., 5, 323-329. #14023
- 377. Ingenito, R., Bianchi, E., Fattori, D., & Pessi, A. (1999) Solid phase synthesis of peptide C-terminal thioesters by Fmoc/t-Bu chemistry. J. Amer. Chem. Soc., 121, 11369-11374. #14025

- 378. Ishida, H., & Inoue, Y. (1999) Peptides that contain unnatural amino acids: Toward artificial proteins. Reviews on Heteroatom Chemistry, 19, 79-142. #12754
- 379. Jang, T.S., Shin, S.Y., Ha, J.M., & Kang, S.W. (1999) Structure-contractile activity relationships of neurokinin A on guinea pig tracheal smooth muscle. Bull. Kor. Chem. Soc., 20, 199-202. #12757
- Jenkins, K.E., Higson, A.P., Seeberger, P.H. & Caruthers, M.H. (1999) Solid phase synthesis of O-boranophosphopeptides and O-dithiophosphopeptides. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 225-228). Mayflower Scientific Limited, Birmingham. #13681
- 381. Jezek, J., Velek, J., Veprek, P., Velkova, V., Trnka, T., Pecka, J., Ledvina, M., Vondrasek, J., & Pisacka, M. (1999) Solid phase synthesis of glycopeptide dendrimers with Tn antigenic structure and their biological activities. Part I. J. Peptide Sci., 5, 46-55. #12759
- 382. Jubilut, G.N., Miranda, M.T., Tominaga, M., Okada, Y., Miranda, A., & Nakaie, C.R. (1999) Resin selection based on the lability of peptidyl-resin linkage towards HF and TFA steps: Dependence on the C-terminal amino acid and peptide length. Chem. Pharm. Bull. Tokyo, 47, 1560-1563. #14239
- 383. Juvvadi, P., Vunnam, S., Yoo, B., & Merrifield, R.B. (1999) Structure-activity studies of normal and retro pig cecropin-melittin hybrids. J. Pept. Res., 53, 244-251. #12874
- 384. Kaiser, T., Luppa, P., & Voelter, W. (1999) Solid-phase synthesis, isolation and analysis of a mouse protein, the macrophage migration inhibitory factor. J. Chromatogr., 852, 189-195. #13536
- 385. Kates, S.A., Cammish, L.E., & Albericio, F. (1999) Continuous-flow solid-phase peptide synthesis using polystyrene resins. J. Pept. Res., 53, 682-683. #12986
- 386. Kim, Y., Zeng, F.W., & Zimmerman, S.C. (1999) Peptide dendrimers from natural amino acids. Chem. Eur. J., 5, 2133-2138. #13539
- 387. Kim, Y.D., Cho, B.K., Park, J.I., Lee, S.H. & Kim, H.J. (1999) Solid phase peptide synthesis using 2-(4-nitrophenyl) sulfonylethoxycarbonyl (Nsc) amino acids for salmon calcitonin. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 315-318). Mayflower Scientific Limited, Birmingham. #13702
- 388. Kiso, Y., Matsumoto, H., Mizumoto, S., Kimura, T., Fujiwara, Y., & Akaji, K. (1999) Small dipeptide-based HIV protease inhibitors containing the hydroxymethylcarbonyl isostere as an ideal transition-state mimic. Biopolymers (Pept. Sci. ), 51, 59-68. #13095
- 389. Koeller, K.M., & Wong, C.H. (1999) Synthesis of biologically active glycopeptides. J. Chin. Chem. Soc., 46, 659-685. #15144
- Kohlbau, H.J., Meisenbach, M., Echner, H. & Voelter, W. (1999) Introduction of different methoxy-substituted 9-phenyl-9-fluoenylamine linkers: Mildly cleavable anchoring groups for the synthesis of peptide amides. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 319-322). Mayflower Scientific Limited, Birmingham. #13703
- 391. Kolodziejczyk, A.S., Sugajska, E., Falkiewicz, B., & Wisniewski, K. (1999) The application of the Pmc-protecting group to reduced peptide bond synthesis. Synlett, 1606-1608. #15150
- 392. Komba, S., Meldal, M., Werdelin, O., Jensen, T., & Bock, K. (1999) Convenient synthesis of Thr and Ser carrying the tumor associated sialyl-(2-->3)-T antigen as building blocks for solid-phase glycopeptide synthesis. J. Chem. Soc. Perkin Trans. 1, 415-419. #12647
- 393. Koppitz, M., Matha, B., & Kessler, H. (1999) Structure investigation of amphiphilic cyclopeptides in isotropic and anisotropic environments: A model study simulating peptide-membrane interactions. J. Peptide Sci., 5, 507-518. #14439
- 394. Kramer, A., Reineke, U., Dong, L., Hoffmann, B., Hoffmuller, U., Winkler, D., Volkmer-Engert, R., & Schneider-Mergener, J. (1999) Spot synthesis: Observations and optimizations. J. Pept. Res., 54, 319-327. #13753
- 395. Kudryavtseva, E.V., Sidorova, M.V., Molokoedov, A.S., Ovchinnikov, M.V., & Bespalova, Z.D. (1999) Directed and spontaneous disulfide bond closure by hydrogen peroxide during the synthesis of endothelin 1 and endothelin 3 [Russian]. Bioorganicheskaia Khimiia, 25, 107-116. #12991

- 396. Kuisle, O., Lolo, M., Quinoa, E., & Riguera, R. (1999) Monitoring the solid-phase synthesis of depsides and depsipeptides. A color test for hydroxyl groups linked to a resin. Tetrahedron, 55, 14807-14812. #14044
- 397. Kuisle, O., Quinoa, E., & Riguera, R. (1999) A general methodology for automated solid-phase synthesis of depsides and depsipeptides. Preparation of a valinomycin analogue. J. Org. Chem., 64, 8063-8075. #14045
- 398. Kutterer, K.M.K., Barnes, M.L., & Arya, P. (1999) Automated, solid-phase synthesis of C-neoglycopeptides: Coupling of glycosyl derivatives to resin-bound peptides. Journal of Combinatorial Chemistry, 1, 28-31. #12039 (Available on Internet)
- 399. Kuttner, G., Kramer, A., Schmidtke, G., Giessmann, E., Dong, L., Roggenbuck, D., Scholz, C., Seifert, M., Stigler, R.D., Schneider-Mergener, J., Porstmann, T., & Hohne, W. (1999) Characterization of neutralizing anti-pre-S1 and anti-pre-S2 (HBV) monoclonal antibodies and their fragments. Mol. Immunol., 36, 669-683. #13808
- 400. Kwak, J., Jefferson, E.A., Bhumralkar, M., & Goodman, M. (1999) Triple helical stabilities of guest-host collagen mimetic structures. Bioorg. Med. Chem., 7, 153-160. #12771
- Lanter, C.L., Guiles, J.W., & Rivero, R.A. (1999) The solid-phase synthesis of novel 14-membered macrocycles for high throughput screening. Mol. Diversity, 4, 149-153. #12996
- Lauer, J.L. & Fields, G.B. (1999) Protein-like molecular structure: Synthesis and application for inducing cellular receptor binding and signal transduction. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 185-188). Mayflower Scientific Limited, Birmingham. #13671
- 403. Lebl, M., Pires, J., Poncar, P., & Pokorny, V. (1999) Evaluation of gaseous hydrogen fluoride as a convenient reagent for parallel cleavage from the solid support. Journal of Combinatorial Chemistry, 1, 474-479. #13836
- 404. Lebl, M. (1999) Parallel personal comments on "classical" papers in combinatorial chemistry. Journal of Combinatorial Chemistry, 1, 3-24. #12028 (Available on Internet)
- Lee, H.B., & Balasubramanian, S. (1999) Studies on a dithiane-protected benzoin photolabile safety catch linker for solid-phase synthesis. J. Org. Chem., 64, 3454-3460. #12877
- 406. Lee, K., Hwang, S.Y., & Park, C.W. (1999) Thrombin inhibitors based on a propargylglycine template. Bioorg. Med. Chem. Lett., 9, 1013-1018. #12998
- 407. Lew, B.M., Mills, K.V., & Paulus, H. (1999) Characteristics of protein splicing in trans mediated by a semisynthetic split intein. Biopolymers (Pept. Sci.), 51, 355-362. #14271
- 408. Li, P., & Xu, J.C. (1999) BOMI: A novel peptide coupling reagent. Tetrahedron Lett., 40, 3605-3608. #12690
- 409. Li, S.H., & Dass, C. (1999) Iron(III)-immobilized metal ion affinity chromatography and mass spectrometry for the purification and characterization of synthetic phosphopeptides. Anal. Biochem., 270, 9-14. #13001
- 410. Li, X.X., Cheng, Y.Z., Zhang, L.G., & Zhang, L.H. (1999) Synthesis of spin labeled conjugate of peptide and peptide nucleic acid. Syn. Commun., 29, 1519-1525. #13002
- 411. Liehr, S., & Chenault, H.K. (1999) A comparison of the alpha-helix forming propensities and hydrogen bonding properties of serine phosphate and alpha-amino-gamma-phosphonobutyric acid. Bioorg. Med. Chem. Lett., 9, 2759-2762. #15174
- 412. Limal, D., Semetey, V., Dalbon, P., Jolivet, M., & Briand, J.P. (1999) Solid-phase synthesis of N,N'-unsymmetrically substituted ureas: Application to the synthesis of carbaza peptides. Tetrahedron Lett., 40, 2749-2752. #12650
- Lippens, G., Bourdonneau, M., Dhalluin, C., Warrass, R., Richert, T., Seetharaman, C., Boutillon, C., & Piotto, M. (1999) Study of compounds attached to solid supports using high resolution magic angle spinning NMR. Current Organic Chemistry, 3, 147-169. #12774
- 414. Lisowski, M., Stasiak, M., & Leplawy, M.T. (1999) The study on helix-inducing propensity of alpha-hydroxymethylserine based on the host-guest approach. Polish J. Chem., 73, 313-319. #12775
- 415. Litowski, J.R., Semchuk, P.D., Mant, C.T., & Hodges, R.S. (1999) Hydrophilic interaction/cation-exchange chromatography for

- the purification of synthetic peptides from closely related impurities: Serine side-chain acetylated peptides. J. Pept. Res., 54, 1-11. #13550
- 416. Lohse, A., Jensen, K.B., & Bols, M. (1999) The first combinatorial library of azasugar glycosidase inhibitors. Tetrahedron Lett., 40, 3033-3036. #12679
- 417. Lu, W.Y., Randal, M., Kossiakoff, A., & Kent, S.B.H. (1999) Probing intermolecular backbone H-bonding in serine proteinase-protein inhibitor complexes. Chem. Biol., 6, 419-427. #13554
- Luke, R.W.A. & Boyce, P.G.T. (1999) Design, synthesis and characterization of a beta-turn mimic library. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 335-338). Mayflower Scientific Limited, Birmingham. #13706
- 419. Lundquist, J.T., & Dix, T.A. (1999) Synthesis and human neurotensin receptor binding activities of neurotensin(8-13) analogues containing position 8 alpha-azido-N-alkylated derivatives of ornithine, lysine, and homolysine. J. Med. Chem., 42, 4914-4918. #14066
- 420. Madder, A., Farcy, N., Hosten, N.G.C., De Muynck, H., De Clercq, P.J., Barry, J., & Davis, A.P. (1999) A novel sensitive colorimetric assay for visual detection of solid-phase bound amines. Eur. J. Org. Chem., 2787-2791. #15987
- 421. Malik, Z., Amir, S., Pal, G., Buzas, Z., Varallyay, E., Antal, J., Szilagyi, Z., Vekey, K., Asboth, B., Patthy, A., & Graf, L. (1999) Proteinase inhibitors from desert locust, Schistocerca gregaria: Engineering of both P-1 and P-1' residues converts a potent chymotrypsin inhibitor to a potent trypsin. Biochimica et Biophysica Acta Protein Structure and Molecular Enzymology, 1434, 143-150. #14069
- 422. Manning, M., Stoev, S., Cheng, L.L., Wo, N.C., & Chan, W.Y. (1999) Synthesis and structure-activity investigation of novel vasopressin hypotensive peptide agonists. J. Peptide Sci., 5, 472-490. #14070
- 423. Marastoni, M., Guerrini, R., Balboni, G., Salvadori, S., Fantin, G., Fogagnolo, M., Lazarus, L.H., & Tomatis, R. (1999) Opioid deltorphin C analogues containing cis- or trans-2- or 3- or 4-aminocyclohexanecarboxylic acid residues. Arzneim. -Forsch., 49, 6-12. #12779
- 424. Marcaurelle, L.A., & Bertozzi, C.R. (1999) New directions in the synthesis of glycopeptide mimetics. Chem. Eur. J., 5, 1384-1390. #13010
- 425. Martin, F., Steinkuhler, C., Brunetti, M., Pessi, A., Cortese, R., De Francesco, R., & Sollazzo, M. (1999) A loop-mimetic inhibitor of the HCV-NS3 protease derived from a minibody. Protein Eng., 12, 1005-1011. #14074
- 426. Maruyama, T., Ikeo, T., & Ueki, M. (1999) A rapid and facile method for the preparation of peptide disulfides. Tetrahedron Lett., 40, 5031-5034. #13111
- Matusina, Z., Olbrimkova, R., Votavova, H., Neumann, J., Hradilek, M., Soucek, M., Malon, P., Kodicek, M., & Stibor, I. (1999)
   Linear heptapeptides containing DNA-intercalators. Synthesis and interaction with DNA. Collect. Czech. Chem. Commun., 64, 1419-1432. #15202
- Matysiak, S., Boldicke, T., Tegge, W. & Frank, R. (1999) Fmoc-Lys(Mmt)-OH: A convenient building block for the preparation of branched and cyclic peptides. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 339-340). Mayflower Scientific Limited, Birmingham. #13707
- 429. Mayfield, L.D., & Corey, D.R. (1999) Enhancing solid phase synthesis by a noncovalent protection strategy-efficient coupling of rhodamine to resin-bound peptide nucleic acids. Bioorg. Med. Chem. Lett., 9, 1419-1422. #12881
- McBride, J.D., Freeman, N. & Leatherbarrow, R.J. (1999) Selection of human elastase inhibitors from a conformationally-constrained combinatorial peptide library. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 167-172). Mayflower Scientific Limited, Birmingham. #13670
- 431. Meisenbach, M., Echner, H., Kohlbau, H.J. & Voelter, W. (1999) Methoxy substituted 9-phenylxanthenylamine linkers: Useful tools for the solid phase synthesis of protected peptide peptide amides. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 345-348). Mayflower Scientific Limited, Birmingham. #13708
- 432. Meldal, M., Tornoe, C., Tedebark, U., Jansson, A.M., Juliano, M.A., Panza, L. & Lay, L. (1999) Azido acids in a novel method of

- solid phase synthesis. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 19-22). Mayflower Scientific Limited, Birmingham. #13453
- 433. Melnyk, O., Rommens, C., & Gras-Masse, H. (1999) Chemistry of hydrazinopeptides: A new hydroperoxydeamination process. Tetrahedron Lett., 40, 1491-1494. #12601
- 434. Merette, S.A.M., Green, D., Patel, G., Scully, M.F., Kakkar, V.V. & Deadman, J.J. (1999) A general method for N to C terminal solid phase peptide synthesis applied to bioactive peptidic boronates. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 349-350). Mayflower Scientific Limited, Birmingham. #13709
- 435. Mergler, M., Dick, F., Gosteli, J., & Nyfeler, R. (1999) Solid phase synthesis of fully protected peptide alcohols. Tetrahedron Lett., 40, 4663-4664. #13015
- 436. Mergler, M. & Nyfeler, R. (1999) Polymeric diphenyldiazomethane: A useful resin for SPPS and solid phase organic chemistry. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 351-35). Mayflower Scientific Limited, Birmingham. #13710
- 437. Mergler, M., Gosteli, J., Grogg, P., Nyfeler, P., & Tanner, R. (1999) SASRIN(TM), a versatile tool in peptide synthesis and solid-phase organic chemistry. Chimia, 53, 29-34. #12782
- 438. Mezo, G., Majer, Z., Valero, M.L., Andreu, D., & Hudecz, F. (1999) Synthesis of cyclic Herpes simplex virus peptides containing 281-284 epitope of glycoprotein D-1 in endo- or exo-position. J. Peptide Sci., 5, 272-282. #13108
- 439. Mezo, G., Mihala, N., Koczan, G. & Hudecz, F. (1999) Application of cyclohexyloxycarbonyl protecting group in Boc chemistry for the synthesis of protected peptide fragment. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 355-356). Mayflower Scientific Limited, Birmingham. #13711
- 440. Miranda, L.P., & Alewood, P.F. (1999) Accelerated chemical synthesis of peptides and small proteins. Proc. Natl. Acad. Sci. USA, 96, 1181-1186. #12785
- Mizuno, M., Haneda, K., Iguchi, R., Muramoto, I., Kawakami, T., Aimoto, S., Yamamoto, K., & Inazu, T. (1999) Synthesis of a glycopeptide containing oligosaccharides: Chemoenzymatic synthesis of eel calcitonin analogues having natural N-linked oligosaccharides. J. Amer. Chem. Soc., 121, 284-290. #12786
- 442. Morris, M.C., Mery, J., Heitz, A., Heitz, F., & Divita, G. (1999) Design and synthesis of a peptide derived from positions 195-244 of human cdc25C phosphatase. J. Peptide Sci., 5, 263-271. #13107
- 443. Muller, B., Besser, D., Kleinwachter, P., Arad, O., & Reissmann, S. (1999) Synthesis of N-carboxyalkyl and N-aminoalkyl functionalized dipeptide building units for the assembly of backbone cyclic peptides. J. Pept. Res., 54, 383-393. #14086
- 444. Nagase, T. (1999) Development of peptidic endothelin receptor antagonists: Elucidating physiological roles of endothelin and medicinal chemistry efforts toward its receptor antagonists [Japanese]. J. Syn. Org. Chem. Jpn., 57, 888-897. #15213
- 445. Nagy, I.B., Mak, M., Varga, I., Kovacs, J., Fellinger, E. & Hudecz, F. (1999) Solid phase synthesis of KDEL peptides labelled with fluorophore and/or bifunctional chelating agent for receptor localisation. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 229-230). Mayflower Scientific Limited, Birmingham. #13682
- 446. Niidome, T., Anzai, S., Sonoda, J., Tokunaga, Y., Nakahara, M., Hatakeyama, T., & Aoyagi, H. (1999) Effect of amino acid substitution in amphiphilic a-helical peptides on peptide-phospholipid membrane interaction. J. Peptide Sci., 5, 298-305. #14432
- 447. Nishiyama, Y., & Kurita, K. (1999) Cyclohexyl ether as a new hydroxy-protecting group for serine in solid-phase peptide synthesis. Tetrahedron Lett., 40, 927-930. #12248
- 448. Nokihara, K. & Yasuhara, T. (1999) Highly efficient preparation of 7KDa peptides by single coupling assembly without capping followed by a single step of purification using reverse-phase HPLC. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 105-110). Mayflower Scientific Limited, Birmingham. #13658
- Offer, J., Quibell, M. & Johnson, T. (1999) A new generation of reversible backbone amide protection. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 357-360). Mayflower Scientific Limited, Birmingham. #13712

- 450. Oh, J.E., Hong, S.Y., & Lee, K.H. (1999) Design, synthesis and characterization of antimicrobial pseudopeptides corresponding to membrane-active peptide. J. Pept. Res., 54, 129-136. #13576
- 451. Ota, M., Sawa, A., Nio, N., & Ariyoshi, Y. (1999) Enzymatic ligation for synthesis of single-chain analogue of monellin by transglutaminase. Biopolymers, 50, 193-200. #13023
- 452. Ottl, J., Musiol, H.J., & Moroder, L. (1999) Heterotrimeric collagen peptides containing functional epitopes, synthesis of single-stranded collagen type I peptides related to the collagenase cleavage site. J. Peptide Sci., 5, 103-110. #12793
- 453. Ottl, J., & Moroder, L. (1999) Disulfide-bridged heterotrimeric collagen peptides containing the collagenase cleavage site of collagen type I. Synthesis and conformational properties. J. Amer. Chem. Soc., 121, 653-661. #12794
- 454. Overkleeft, H.S., Verhelst, S.H.L., Pieterman, E., Meeuwenoord, N.J., Overhand, M., Cohen, L.H., van der Marel, G.A., & van Boom, J.H. (1999) Design and synthesis of a protein: Farnesyltransferase inhibitor based on sugar amino acids. Tetrahedron Lett., 40, 4103-4106. #12898
- 455. Panek, Z., Pavlik, M. & Flegel, M. (1999) Asparagine as racemization sensitive site in analogs of neurohypophyseal hormones. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 365-366). Mayflower Scientific Limited, Birmingham. #13714
- 456. Papageorgopoulos, C., Caldwell, K., Shackleton, C., Schweingrubber, H., & Hellerstein, M.K. (1999) Measuring protein synthesis by mass isotopomer distribution analysis (MIDA). Anal. Biochem., 267, 1-16. #12799
- Paris, M., Pothion, C., Heitz, A., Rocheblave, L., Rouch, F., Fehrentz, J.A. & Martinez, J. (1999) Solid phase synthesis of peptide aldehydes. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 15-18).
   Mayflower Scientific Limited, Birmingham. #13452
- 458. Paris, M., Douat, C., Heitz, A., Gibbons, W., Martinez, J., & Fehrentz, J.A. (1999) Post-synthesis modification of aspartyl or glutamyl residue side-chains on solid support. Tetrahedron Lett., 40, 5179-5182. #13118
- 459. Patel, G., Husman, W., Jehanli, A.M., Deadman, J.J., Green, D., Kakkar, V.V., & Brennand, D.M. (1999) A cyclic peptide analogue of the loop III region of platelet-derived growth factor-BB is a synthetic antigen for the native protein. J. Pept. Res., 53, 68-74. #12654
- 460. Patterson, J.A., & Ramage, R. (1999) Solid phase synthesis of peptide C-terminal semicarbazones and aldehydes. Tetrahedron Lett., 40, 6121-6124. #13579
- 461. Pearce, A.J., Ramaya, S., Thorn, S.N., Bloomberg, G.B., Walter, D.S., & Gallagher, T. (1999) C-glycosyl tyrosines. Synthesis and incorporation into C-glycopeptides. J. Org. Chem., 64, 5453-5462. #13580
- 462. Pedersen, W.B., Almdal, K., Winther, L., & Berg, R.H. (1999) R.D. Forskniingscenter Riso. Grafted cross-linked polyolefin substrates for peptide synthesis and assays. US Patent. 5,886,104. 569,255, #13154
- 463. Pegoraro, S., Besse, D., Fiori, S., Rudolph-Bohner, S., Watanabe, T.X., Kimura, T. & Moroder, L. (1999) Synthesis and folding properties of selenocystine-containing peptides. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 89-92). Mayflower Scientific Limited, Birmingham. #13466
- Peluso, S., Dumy, P., Nkubana, C., Yokokawa, Y., & Mutter, M. (1999) Solid-phase strategies for the assembly of template-based protein mimetics. J. Org. Chem., 64, 7114-7120. #13816
- 465. Peng, Z.H. (1999) Solid phase synthesis and NMR conformational studies on cyclic decapeptide template molecule. Biopolymers, 49, 565-574. #12884
- 466. Peng, Z.H. (1999) NMR conformational analysis on cyclic decapeptide template molecule. Can. J. Chem., 77, 1394-1404. #13817
- 467. Pennington, M.W., Baur, P., Westling, J. & Dunn, B.M. (1999) Using 2-chlorotrityl resin tactics for the preparation of DABCYL-EDANS derivatized peptides: Application to a fluorogenic substrate for an aspartyl proteinase from human malaria parasite. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 367-370). Mayflower Scientific Limited, Birmingham. #13715

- 468. Peri, F., Grell, D., Dumy, P., Yokokawa, Y., Welzenbach, K., Weitz-Schmidt, G., & Mutter, M. (1999) Assembly of binding loops on aromatic templates as VCAM-1 mimetics. J. Peptide Sci., 5, 313-322. #14433
- 469. Perich, J.W., Black, C.L., Huq, N.L., & Reynolds, E.C. (1999) Epitope analysis of the multiphosphorylated peptide aS1-casein(59-79). J. Peptide Sci., 5, 221-233. #14429
- 470. Peterson, J.J., Pak, R.H., & Meares, C.F. (1999) Total solid-phase synthesis of 1,4,7,10-tetraazacyclododecane-N,N',N",N"-tetraacetic acid-functionalized peptides for radioimmunotherapy. Bioconjug. Chem., 10, 316-320. #12656
- 471. Pinilla, C., Martin, R., Gran, B., Appel, J.R., Boggiano, C., Wilson, D.B., & Houghten, R.A. (1999) Exploring immunological specificity using synthetic peptide combinatorial libraries. Curr. Opin. Immunol., 11, 193-202. #12800
- 472. Poupart, M.A., Fazal, G., Goulet, S., & Mar, L.T. (1999) Solid-phase synthesis of peptidyl trifluoromethyl ketones. J. Org. Chem., 64, 1356-1361. #12658
- 473. Ramage, R., Jiang, L., Kim, Y.D., Shaw, K., Park, J.L., & Kim, H.J. (1999) Comparative studies of Nsc and Fmoc as Naprotecting groups for SPPS. J. Peptide Sci., 5, 195-200. #12903
- 474. Reineke, U., Kramer, A., & Schneider-Mergener, J. (1999) Antigen sequence- and library-based mapping of linear and discontinuous protein-protein-interation sites by spot synthesis. Curr. Topics Microbiol. Immun., 243, 23-36. #13750
- 475. Renil, M., & Pillai, V.N.R. (1999) Solid-phase peptide synthesis using a new PS-TTEGDA resin: Synthesis of pardaxin (1-26). Ind. J. Chem. B-Org. Chem. Med. Chem., 38, 1030-1035. #15267
- 476. Rinnova, M., Lebl, M., & Soucek, M. (1999) Solid-phase peptide synthesis by fragment condensation: Coupling in swelling volume. Lett. Peptide Sci., 6, 15-22. #13834
- Rinnova, M., Soucek, M., Voburka, Z., Lebl, M. & Fusek, M. (1999) Sanger's reagent: An excellent end-capping and/or labeling agent for SPPS. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 385-386). Mayflower Scientific Limited, Birmingham. #13719
- 478. Rivier, J.E., Kirby, D.A., Lahrichi, S.L., Corrigan, A., Vale, W.W., & Rivier, C.L. (1999) Constrained corticotropin releasing factor antagonists (astressin analogues) with long duration of action in the rat. J. Med. Chem., 42, 3175-3182. #13591
- 479. Robertson, N., & Ramage, R. (1999) Total chemical synthesis of deglycosylated human erythropoietin. J. Chem. Soc., 1015-1021. #13038
- 480. Robertson, N., Jiang, L., & Ramage, R. (1999) Racemisation studies of a novel coupling reagent for solid phase peptide synthesis. Tetrahedron, 55, 2713-2720. #12660
- 481. Robinson, B.M., Tellam, D.J., Smart, D., Mohammad, Y.N., Brennand, J., Rivier, J.E., & Lovejoy, D.A. (1999) Cloning and characterization of corticotropin-releasing factor and urocortin in Syrian hamster (Mesocricetus auratus). Peptides, 20, 1177-1185. #14118
- 482. Roggero, M.A., Servis, C. & Corradin, G. (1999) Purification of synthetic polypeptides by immobilized metal affinity chromatography and methionine chemistry. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 111-114). Mayflower Scientific Limited, Birmingham. #13659
- 483. Roice, M., Kumar, K.S., & Pillai, V.N.R. (1999) Synthesis, characterization, and application of butanediol dimethacrylate cross-linked polystyrene: A flexible support for gel phase peptide synthesis. Macromolecules, 32, 8807-8815. #15276
- 484. Rose, K., & Vizzavona, J. (1999) Stepwise solid-phase synthesis of polyamides as linkers. J. Amer. Chem. Soc., 121, 7034-7038. #13592
- 485. Rosenthal, K., Erlandsson, M., & Und n, A. (1999) 4-(3-Hydroxy-4-methylpentyl)phenylacetic acid as a new linker for the solid phase synthesis of peptides with Boc chemistry. Tetrahedron Lett., 40, 377-380. #12106
- 486. Sampson, W.R., Patsiouras, H., & Ede, N.J. (1999) The synthesis of 'difficult' peptides using 2-hydroxy-4-methoxybenzyl or pseudoproline amino acid building blocks: A comparative study. J. Peptide Sci., 5, 403-409. #14125

- 487. Samson, F., Bonnet, D., Rommens, C., Gras-Masse, H., & Melnyk, O. (1999) Reactivity of Lys(NH2)-containing peptides toward endopeptidases. J. Peptide Sci., 5, 352-359. #14422
- 488. Sanz-Nebot, V., Garces, A., & Barbosa, J. (1999) Investigation of crudes of synthesis of carbetocin by liquid chromatography coupled to electrospray ionization mass spectrometry. J. Chromatogr., 833, 267-275. #13041
- 489. Sanz-Nebot, V., Toro, I., & Barbosa, J. (1999) Fractionation and characterization of a crude peptide mixture from the synthesis of eledoisin by liquid chromatography-electrospray ionization mass spectrometry. J. Chromatogr. A, 846, 25-38. #13418
- Sanz-Nebot, V., Toro, I., Garces, A., & Barbosa, J. (1999) Separation and identification of peptide mixtures in a synthesis crude of carbetocin by liquid chromatography/electrospray ionization mass spectrometry. Rapid Commun. Mass Spectrom., 13, 2341-2347. #15292
- 491. Sasaki, Y., Suto, T., Ambo, A., Ouchi, H., & Yamamoto, Y. (1999) Biological properties of opioid peptides replacing Tyr at position 1 by 2,6-dimethyl-Tyr. Chem. Pharm. Bull. Tokyo, 47, 1506-1509. #14128
- 492. Scheller, A., Oehlke, J., Wiesner, B., Dathe, M., Krause, E., Beyermann, M., Melzig, M., & Bienert, M. (1999) Structural requirements for cellular uptake of a-helical amphipathic peptides. J. Peptide Sci., 5, 185-194. #12902
- 493. Scott, R.H., Barnes, C., Gerhard, U., & Balasubramanian, S. (1999) Exploring a chemical encoding strategy for combinatorial synthesis using Friedel-Crafts alkylation. Chem. Commun., 1331-1332. #13821
- 494. Seebach, D., Abele, S., Gademann, K., & Jaun, B. (1999) Pleated sheets and turns of beta-peptides with proteinogenic side chains. Angew. Chem., 1595-1597. #13047
- 495. Seitz, O. (1999) Solid phase synthesis of protected peptide nucleic acids. Tetrahedron Lett., 40, 4161-4164. #12889
- 496. Sheridan, J.M., Hayes, G. & Austen, B.M. (1999) Convergent solid phase synthesis of immunoglobulin-like domains of macrophage specific Fc receptor (FcgRI). In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 391-394). Mayflower Scientific Limited, Birmingham. #13721
- 497. Sheridan, J.M., Hayes, G.M., & Austen, B.M. (1999) Solid-phase synthesis and cyclization of a large branched peptide from IgG Fc with affinity for Fc gamma RI. J. Peptide Sci., 5, 555-562. #14138
- 498. Shin, Y., Winans, K.A., Backes, B.J., Kent, S.B.H., Ellman, J.A., & Bertozzi, C.R. (1999) Fmoc-based synthesis of peptide-(alpha) thioesters: Application to the total chemical synthesis of a glycoprotein by native chemical ligation. J. Amer. Chem. Soc., 121, 11684-11689. #14141
- Sidorova, M.V., Molokoedov, A.S., Azmuko, A.A., Stambolskii, D.V., Kashirina, N.M., Bochkov, V.N., Tkachuk, V.A., & Bespalova, Z.D. (1999) Antibodies to synthetic peptide fragments of T-cadherin as inhibitors of T-cadherin binding to low density lipoproteins [Russian]. Bioorganicheskaia Khimiia, 25, 171-178. #13051
- 500. Siev, D.V., Gaudette, J.A., & Semple, J.E. (1999) Novel protocol for the solid-phase synthesis of peptidyl and peptidomimetic Plargininal derivatives. Tetrahedron Lett., 40, 5123-5127. #13117
- 501. Sithigorngul, P., Panchan, N., Vilaivan, T., Sithigorngul, W., & Petsom, A. (1999) Immunochemical analysis and immunocytochemical localization of crustacean hyperglycemic hormone from the eyestalk of Macrobrachium rosenbergii. Comparative Biochemistry and Physiology B-Biochemistry and Molecular Biology, 124, 73-80. #14146
- Sjolin, P., George, S.K., Bergquist, K.E., Roy, S., Svensson, A., & Kihlberg, J. (1999) Synthesis of deoxy and alanine-substituted derivatives of a T cell stimulating glycopeptide: An investigation of conditions for cleavage from the solid phase and deprotection.
   J. Chem. Soc. Perkin Trans. 1, 1731-1742. #13421
- 503. Slootstra, J.W., Schaaper, W.M.M., Puijk, W.C., Ligtvoet, G.J., Kuperus, D. & Meloen, R.H. (1999) Contribution of solid-phase in antibody binding activity of sythetic peptides. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 300-400). Mayflower Scientific Limited, Birmingham. #13722
- Smith, A.B., Benowitz, A.B., Sprengeler, P.A., Barbosa, J., Guzman, M.C., Hirschmann, R., Schweiger, E.J., Bolin, D.R., Nagy, Z., Campbell, R.M., Cox, D.C., & Olson, G.L. (1999) Design and synthesis of a competent pyrrolinone-peptide hybrid ligand for the class II major histocompatibility complex protein HLA-DR1. J. Amer. Chem. Soc., 121, 9286-9298. #14147

- 505. Smith, M.D., & Fleet, G.W.J. (1999) Designing secondary structures: 5-Azidomethyl tetrahydrofuran-2-carboxylates as carbohydrate-derived dipeptide isosteres. J. Peptide Sci., 5, 425-441. #14435
- 506. Soth, M.J., & Nowick, J.S. (1999) A peptide/oligourea/azapeptide hybrid that adopts a hairpin turn. J. Org. Chem., 64, 276-281. #12447
- 507. Spetzler, J.C., & Hoeg-Jensen, T. (1999) Preparation and application of O-amino-serine, Ams, a new building block in chemoselective ligation chemistry. J. Peptide Sci., 5, 582-592. #14150
- 508. St. Hilaire, P.M., Willert, M., Juliano, M.A., Juliano, L., & Meldal, M. (1999) Fluorescence-quenched solid phase combinatorial libraries in the characterization of cysteine protease substrate specificity. Journal of Combinatorial Chemistry, 1, 509-523. #14151
- St.Hilaire, P.M., Cipolla, L., Franco, A., Tedebark, U., Tilly, D.A., & Meldal, M. (1999) Synthesis of T-antigen-containing glycopeptides as potential cancer vaccines. J. Chem. Soc. Perkin Trans. 1, 3559-3564. #14152
- 510. Stoev, S., Cheng, L.L., Olma, A., Klis, W.A., Manning, M., Sawyer, W.H., Wo, N.C., & Chan, W.Y. (1999) An investigation of position 3 in arginine vasopressin with aliphatic, aromatic, conformationally-restricted, polar and charged amino acids. J. Peptide Sci., 5, 141-153. #12816
- 511. Suguro, T., & Yanai, M. (1999) Solid-phase synthesis of cyclooctadepsipeptide N-4909 using a cyclization-cleavage method with oxime resin. Journal of Antibiotics, 52, 835-838. #14157
- Szekely, Z., Zakhariev, S., Guarnaccia, C., Antcheva, N., & Pongor, S. (1999) A highly effective method for synthesis of Nomega-substituted arginines as building blocks for Boc/Fmoc peptide chemistry. Tetrahedron Lett., 40, 4439-4442. #13059
- 513. Tam, J.P., Lu, Y.A., & Yu, Q.T. (1999) Thia zip reaction for synthesis of large cyclic peptides: Mechanisms and applications. J. Amer. Chem. Soc., 121, 4316-4324. #13062
- 514. Tam, J.P., Yu, Q., & Miao, Z. (1999) Orthogonal ligation strategies for peptide and protein. Biopolymers (Pept. Sci. ), 51, 311-332. #14269
- Tatsu, Y., Shigeri, Y., Ishida, A., Kameshita, I., Fujisawa, H., & Yumoto, N. (1999) Synthesis of caged peptides using caged lysine: Application to the synthesis of caged AIP, a highly specific inhibitor of calmodulin-dependent protein kinase II. Bioorg. Med. Chem. Lett., 9, 1093-1096. #13064
- 516. Thieriet, N., Alsina, J., Dessolin, M., Guillerez, M.G., Giralt, E., Albericio, F., Loffet, A. & Guibe, F. (1999) PhSiH3/Paladium catalyst/carboxyactivated aminoacids: A ternary system for tandem deprotection coupling of Na --allocaminoacids and suppression of DKP formation in SPPS. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 127-128). Mayflower Scientific Limited, Birmingham. #13662
- 517. Tholey, A., Lindemann, A., Kinzel, V., & Reed, J. (1999) Direct effects of phosphorylation on the preferred backbone conformation of peptides: A nuclear magnetic resonance study. Biophys. J., 76, -87. #12454
- 518. Thorpe, D.S., Yeoman, H., Chan, A.W.E., Krchnak, V., Lebl, M., & Felder, S. (1999) Combinatorial chemistry reveals a new motif that binds the platelet fibrinogen receptor, gpIIbIIIa. Biochem. Biophys. Res. Commun., 256, 537-541. #12667
- 519. Tomboly, C., Spetea, M., Borsodi, A., & Toth, G. (1999) Synthesis of tritium labelled endomorphin II and its stability in the radioreceptor assay. Czechoslovak Journal of Physics, 49, 896. #13065
- 520. Tsai, J.H., Waldman, A.S., & Nowick, J.S. (1999) Two new beta-strand mimics. Bioorg. Med. Chem., 7, 29-38. #12824
- 521. Ueki, M., Ikeo, T., Iwadate, M., Asakura, T., Williamson, M.P., & Slaninova, J. (1999) Solid phase synthesis and biological activities of [Arg(8)]-vasopressin methylenedithioether. Bioorg. Med. Chem. Lett., 9, 1767-1772. #13068
- 522. Utkin, Y.N., Zhmak, M.N., Methfessel, C., & Tsetlin, V.I. (1999) Aromatic substitutions in alpha-conotoxin ImI. Synthesis of iodinated photoactivatable derivative. Toxicon, 37, 1683-1695. #14176
- 523. Valero, M.L., Giralt, E., & Andreu, D. (1999) A comparative study of cyclization strategies applied to the synthesis of head-to-tail cyclic analogs of a viral epitope. J. Pept. Res., 53, 56-67. #12668

- 524. Varkey, J.T., & Pillai, V.N.R. (1999) Merrifield resin and 1,6-hexanediol diacrylate-crosslinked polystyrene resin for solid-phase peptide synthesis: A comparative study. J. Appl. Polym. Sci., 71, 1933-1939. #12460
- 525. Varkey, J.T., & Pillai, V.N.R. (1999) Solid phase synthesis of hydrophobic peptides on 1,6-hexanediol diacrylate cross-linked polystyrene resin. J. Peptide Sci., 5, 577-581. #14179
- 526. Vazquez, E., Caamano, A.M., Castedo, L., Gramberg, D., & Mascarenas, J.L. (1999) A practical approach to orthogonally connected oligopyrrole-peptide conjugates. Tetrahedron Lett., 40, 3625-3628. #12692
- 527. Vazquez, E., Caamano, A.M., Castedo, L., & Mascarenas, J.L. (1999) An Fmoc solid-phase approach to linear polypyrrole-peptide conjugates. Tetrahedron Lett., 40, 3621-3624. #12691
- 528. Vella, F., Hernandez, J.F., Molla, A., Block, M.R., & Arlaud, G.J. (1999) Grafting an RGD motif onto an epidermal growth factor-like module: chemical synthesis and functional characterization of the chimeric molecule. J. Pept. Res., 54, 415-426. #14180
- 529. Veprek, P., & Jezek, J. (1999) Peptide and glycopeptide dendrimers. Part II. J. Peptide Sci., 5, 203-220. #13069
- 530. Veprek, P., & Jezek, J. (1999) Peptide and glycopeptide dendrimers. Part I. J. Peptide Sci., 5, 5-23. #12828
- 531. Vetter, S. (1999) High-affinity non-inhibitory ligands for alkaline phosphatase revealed from a restricted heptapeptide library. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 407-410). Mayflower Scientific Limited, Birmingham. #13723
- 532. Voelter, W., Kaiser, T., Hannappel, E., & Echner, H. (1999) Synthesis of Thymosin beta(Xen)(4) and its comparison with the natural tritetraconpeptide. Journal fur Praktische Chemie-Chemiker-Zeitung, 341, 47-51. #12830
- 533. Wang, P., Landon, M., Layfield, R., Mayer, R.J. & Ramage, R. (1999) Transfer active ester condensation: A novel technique used for peptide segment coupling. In R. Epton (Ed.), Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. (pp. 415-416). Mayflower Scientific Limited, Birmingham. #13724
- 534. Wang, W., & McMurray, J.S. (1999) A selective method for the preparation of primary amides: Synthesis of Fmoc-L-4-carboxamidophenylalanine and other compounds. Tetrahedron Lett., 40, 2501-2504. #12677
- 535. Wildemann, D., Drewello, M., Fischer, G., & Schutkowski, M. (1999) Extremely selective Mg(ClO4)2 mediated removal of Bpoc/Ddz moieties suitable for the solid phase peptide synthesis of thioxo peptides. Chem. Commun., 1809-1810. #13829
- 536. Wu, Y.D., & Wang, D.P. (1999) Theoretical study on side-chain control of the 14-helix and the 10/12-helix of beta-peptides. J. Amer. Chem. Soc., 121, 9352-9362. #14195
- 537. Wu, Y.T., Hsieh, H.P., Chen, S.T., & Wang, K.T. (1999) Direct solid phase synthesis of biologically active peptide alcohols. J. Chin. Chem. Soc., 46, 135-138. #13077
- 538. Xu, J.X., & Jin, S. (1999) Studies on synthesis, structure and antitumor activities of analogues of papaver somniferum pollen tridecapeptide [Chinese]. Chem. J. Chin. Univ., 20, 722-726. #13079
- 539. Yang, L., & Morriello, G. (1999) Solid phase synthesis of 'head-to-tail' cyclic peptides using a sulfonamide 'safety-catch' linker: The cleavage by cyclization approach. Tetrahedron Lett., 40, 8197-8200. #13839
- 540. Yao, S., & Chmielewski, J. (1999) A pH-tunable peptide ligase. Biopolymers (Pept. Sci.), 51, 370-375. #14273
- 541. Yao, Z.J., Gao, Y., Voigt, J.H., Ford, H., & Burke, T.R. (1999) Synthesis of Fmoc-protected 4-carboxydifluoromethyl-L-phenylalanine: A phosphotyrosyl mimetic of potential use for signal transduction studies. Tetrahedron, 55, 2865-2874. #12838
- 542. Yao, Z.J., King, C.R., Cao, T., Kelley, J., Milne, G.W.A., Voigt, J.H., & Burke, T.R. (1999) Potent inhibition of Grb2 SH2 domain binding by non-phosphate-containing ligands. J. Med. Chem., 42, 25-35. #12475
- 543. Yoshida, K., Ohmori, N., Mukai, Y., Niidome, T., Hatakeyama, T., & Aoyagi, H. (1999) Interaction of bundled Ser-rich amphiphilic peptides with phospholipid membranes. J. Peptide Sci., 5, 360-367. #14423
- 544. Yoshikawa, M., Fujisawa, T., & Izumi, J. (1999) Molecularly imprinted polymeric membranes having EFF derivatives as a chiral

- recognition site. Macromolecular Chemistry and Physics, 200, 1458-1465. #13088
- 545. Yu, C.X., & Taylor, J.W. (1999) Synthesis and study of peptides with semirigid i and i+7 side-chain bridges designed for alphahelix stabilization. Bioorg. Med. Chem., 7, 161-175. #12839
- 546. Zhang, H., Wang, Y., Thurmer, R., Al-Qawasmeh, R.A., & Voelter, W. (1999) Total synthesis of the nephritogenoside glycopeptide. Polish J. Chem., 73, 101-115. #12842
- 547. Zhang, H., Wang, Y.L., Thurmer, R., Parvez, K., Choudhary, I., Atta, u.R., & Voelter, W. (1999) Neighbouring group participation of C-6 substituents of glucose derivatives on the stereoselectivity of the N-glycosidic linkage of glycopeptides. Zeitschrift für Naturforschung Section B-A Journal of Chemical Sciences, 54, 692-698. #13089
- 548. Zhang, L.S., & Tam, J.P. (1999) Lactone and lactam library synthesis by silver ion-assisted orthogonal cyclization of unprotected peptides. J. Amer. Chem. Soc., 121, 3311-3320. #12897
- 549. Zheng, A., Shan, D., & Wang, B. (1999) A redox-sensitive resin linker for the solid phase synthesis of C-terminal modified peptides. J. Org. Chem., 64, 156-161. #12199
- 550. Zheng, A.L., Shan, D.X., Shi, X.L., & Wang, B.H. (1999) A novel resin linker for solid-phase peptide synthesis which can be cleaved using two sequential mild reactions. J. Org. Chem., 64, 7459-7466. #14205
- 551. Abdelmoaty-Khalil, I. (1998) Solid-state chemistry of organic compounds. Bradeis University, Waltham, MA, USA; 1 p. UMI, Order No.DA9829810.#11469
- 552. Albericio, F., Bofill, J.M., El-Faham, A., & Kates, S.A. (1998) Use of onium salt-based coupling reagents in peptide synthesis. J. Org. Chem., 63, 9678-9683. #12300
- 553. Albert, R., Smith-Jones, P., Stolz, B., Simeon, C., Knecht, H., Bruns, C., & Pless, J. (1998) Direct synthesis of [DOTA-DPhe(1)]-octreotide and [DOTA-DPhe(1),Tyr(3)]-octreotide (SMT487): Two conjugates for systemic delivery of radiotherapeutical nuclides to somatostatin receptor positive tumors in man. Bioorg. Med. Chem. Lett., 8, 1207-1210. #11096
- 554. Aldrian-Herrada, G., Rabie, A., Wintersteiger, R., & Brugidou, J. (1998) Solid-phase synthesis of peptide nucleic acid (PNA) monomers and their oligomerization using disulphide anchoring linkers. J. Peptide Sci., 4, 266-281. #10878
- 555. Andell-Jonsson, S., & Bartfai, T. (1998) Identification of the spinal degradation products and inhibition of adenylate cyclase by recombinant rat galanin message-associated peptide. Neuropeptides, 32, 191-196. #11101
- 556. Anderssen, E.L., Diep, D.B., Nes, I.F., Eijsink, V.G.H., & Nissenmeyer, J. (1998) Antagonistic activity of lactobacillus plantarum C11: Two new two-peptide bacteriocins, plantaricins EF and JK, and the induction factor plantaricin A. Appl. Environ. Microbiol., 64, 2269-2272. #11089
- 557. Angeletti, R.H., Bibbs, L., Bonewald, L.F., Fields, G.B., Kelly, J.W., McMurray, J.S., Moore, W.T. & Weintraub, S.T. (1998) A multicenter study of racemization during "standard" solid phase peptide synthesis. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 207-208). Mayflower Scientific Ltd., Kingswinford. #11501
- 558. Annis, D.A., Helluin, O., & Jacobsen, E.N. (1998) Stereochemistry as a diversity element: Solid-phase synthesis of cyclic RGD peptide derivatives by asymmetric catalysis. Angew. Chem. Int. Ed., 37, 1907-1909. #11251
- 559. Annis, I., Chen, L., & Barany, G. (1998) Novel solid-phase reagents for facile formation of intramolecular disulfide bridges in peptides under mild conditions. J. Amer. Chem. Soc., 120, 7226-7238. #12235
- Asche, G., Kunz, H., Nar, H., Koppen, H., Briem, H., Pook, K.H., Schiller, P.W., Chung, N.N., Lemieux, C., & Esser, F. (1998) Synthesis of cyclic dipeptide templates, their incorporation into peptides and studies on their conformational and biological properties. J. Pept. Res., 51, 323-336. #11140
- 561. Aubagnac, J.L., Enjalbal, C., Subra, G., Bray, A.M., Combarieu, R., & Martinez, J. (1998) Application of time-of-flight secondary ion mass spectrometry to in situ monitoring of solid-phase peptide synthesis on the Multipin(TM) system. J. Mass Spectrom., 33, 1094-1103. #12136
- 562. Augustyns, K., Kraas, W., & Jung, G. (1998) Investigation on the stability of the Dde protecting group used in peptide synthesis:

- Migration to an unprotected lysine. Int. J. Peptide Prot. Res., 51, 127-133. #10732
- 563. Avdagic, A., & Sunjic, V. (1998) Biocatalytic deracemization of 1,4-benzodiazepines in the synthesis of enantiomerically pure serine. Helv. Chim. Acta, 81, 85-92. #10684
- 564. Barlos, K., Gatos, D., & Koutsogianni, S. (1998) Fmoc/Trt amino acids: Comparison to Fmoc/tBu amino acids in peptide synthesis. Int. J. Peptide Prot. Res., 51, 194-200. #10710
- 565. Belhadj Jrad, B., & Bahraoui, E. (1998) Antigenicity of linear and cyclic peptides mimicking the disulfide loops in HIV-2 envelope glycoprotein: Synthesis, reoxidation and purification. J. Pept. Res., 51, 370-385. #11548
- 566. Beltran, M., Pedroso, E., & Grandas, A. (1998) A comparison of histidine protecting groups in the synthesis of peptideoligonucleotide conjugates. Tetrahedron Lett., 39, 4115-4118. #10841
- Bengtsson, M., Broddefalk, J., Dahmen, J., Henriksson, K., Kihlberg, J., Lonn, H., Srinivasa, B.R., & Stenvall, K. (1998)
   Convergent synthesis of neoglycopeptides by coupling of 2-bromoethyl glycosides to cysteine and homocysteine residues in T cell stimulating peptides. Glycoconjugate J., 15, 223-231. #10703
- 568. Bezay, N., Fuchs, G., Habermann, J., Liebe, B., Lohr, B., Seitz, O. & Kunz, H. (1998) Allylic anchoring groups in solid phase peptide synthesis. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 257-258). Mayflower Scientific Ltd., Kingswinford. #11502
- 569. Bianco, A., Zabel, C., Walden, P., & Jung, G. (1998) N-hydroxy-amide analogues of MHC-class I peptide ligands with nanomolar binding affinities. J. Peptide Sci., 4, 471-478. #12313
- 570. Bleicher, K.H., & Wareing, J.R. (1998) PhFl acetic acid: A new linker for solid phase organic synthesis. Tetrahedron Lett., 39, 4591-4594. #11164
- 571. Blondelle, S.E., Nefzi, A., Ostresh, J.M., & Houghten, R.A. (1998) Novel antifungal compounds derived from heterocyclic positional scanning. Combinatorial libraries. Pure Appl. Chem., 70, 2107-2145. #12907
- 572. Boeijen, A., & Liskamp, R.M.J. (1998) Sequencing of peptoid peptidomimetics by Edman degradation. Tetrahedron Lett., 39, 3589-3592. #10810
- 573. Bose, D.S., & Lakshminarayana, V. (1998) An efficient and highly selective cleavage of N-tert-butoxycarbonyl group under microwave irradiation. Tetrahedron Lett., 39, 5631-5634. #10955
- 574. Bosze, S., Kajtar, J., Szabo, R., Falus, A., & Hudecz, F. (1998) Synthesis, solution conformation and interleukin-6-related activities of interleukin-6 peptides. J. Pept. Res., 52, 216-228. #11826
- 575. Broddefalk, J., Bergquist, K.E., & Kihlberg, J. (1998) Use of acid-labile protective groups for carbohydrate moieties in synthesis of glycopeptides related to type II collagen. Tetrahedron, 54, 12047-12070. #11828
- 576. Bunin, B.A. (1998) The Combinatorial Index. Academic Press, San Diego. ISBN 0121413403 #10496
- 577. Burke, T.R., Yao, Z.J., Zhao, H., Milne, G.W.A., Wu, L., Zhang, Z.Y., & Voigt, J.H. (1998) Enantioselective synthesis of nonphosphorus-containing phosphotyrosyl mimetics and their use in the preparation of tyrosine phosphatase inhibitory peptides. Tetrahedron, 54, 9981-9994. #11262
- 578. Caciagli, V., Cardinali, F., Righetti, G. & Lombardi, P. (1998) Sterically hindered amino acids in SPPS: A study of their coupling reactions under dilute conditions. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 283-284). Mayflower Scientific Ltd., Kingswinford. #11555
- 579. Camarero, J.A., Ayers, B., & Muir, T.W. (1998) Studying receptor-ligand interactions using encoded amino acid scanning. Biochemistry, 37, 7487-7495. #11094
- 580. Camarero, J.A., Cotton, G.J., Adeva, A., & Muir, T.W. (1998) Chemical ligation of unprotected peptides directly from a solid support. J. Pept. Res., 51, 303-316. #11165
- 581. Campian, E., Peterson, M.L., Saneii, H.H. & Furka, A. (1998) Automated synthesis in a single process of nine first order sub-

- libraries of an amino acid occurrence library. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 285-286). Mayflower Scientific Ltd., Kingswinford. #11503
- 582. Campian, E., Chou, J., Peterson, M.L., Saneii, H.H. & Furka, A. (1998) Synthesis and aplicability of monomer tester mixtures. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 131-134). Mayflower Scientific Ltd., Kingswinford. #11494
- 583. Campian, E., Peterson, M.L., Saneii, H.H., & Furka, A. (1998) Deconvolution by omission libraries. Bioorg. Med. Chem. Lett., 8, 2357-2362. #11207
- 584. Carell, T., Schmid, H., & Reinhard, M. (1998) Efficient syntheses of a flavin and an 8-hydroxy-5-deazaflavin amino acid and their incorporation into oligopeptides. J. Org. Chem., 63, 8741-8747. #12325
- 585. Carrasco, M.R., Fitzgerald, M.C. & Kent, S.B.H. (1998) Real-time analysis of solid phase peptide synthesis reactions by matrix-assisted laser desorption/ionization mass spectrometry. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 293-294). Mayflower Scientific Ltd., Kingswinford. #11504
- 586. Cavallaro, V., Thompson, P., & Hearn, M. (1998) Solid phase synthesis of cyclic peptides: Model studies involving i-(i+4) side chain-to-side chain cyclization. J. Peptide Sci., 4, 335-343. #11264
- 587. Cervini, L.A., Donaldson, C.J., Koerber, S.C., Vale, W.W., & Rivier, J.E. (1998) Human growth hormone-releasing hormone hGHRH(1-29)-NH2: Systematic structure-activity relationship studies. J. Med. Chem., 41, 717-727. #10702
- 588. Chargelegue, D., Obeid, O.E., Hsu, S.C., Shaw, M.D., Denbury, A.N., Taylor, G., & Steward, M.W. (1998) A peptide mimic of a protective epitope of respiratory syncytial virus selected from a combinatorial library induces virus-neutralizing antibodies and reduces viral load in vivo. J. Virol., 72, 2040-2046. #10580
- 589. Chen, B. & Turner, A.P.F. (1998) Generation of a peptide library in searching for a specific binding ligand for glycosylated haemoglobin HbA(1C). In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 305-306). Mayflower Scientific Ltd., Kingswinford. #11505
- 590. Chen, H., Pyluck, A.L., Janik, M., & Sampson, N.S. (1998) Peptides corresponding to the epidermal growth factor-like domain of mouse fertilin: Synthesis and biological activity. Biopolymers, 47, 299-307. #12329
- 591. Chhabra, S.R., Hothi, B., Evans, D.J., White, P.D., Bycroft, B.W., & Chan, W.C. (1998) An appraisal of new variants of the Dde amine protecting group for solid phase peptide synthesis. Tetrahedron Lett., 39, 1603-1606. #10494
- 592. Chhabra, S.R., Khan, A.N., & Bycroft, B.W. (1998) Versatile Dde-based primary amine linkers for solid phase synthesis. Tetrahedron Lett., 39, 3585-3588. #10809
- 593. Chillemi, F., Francescato, P., Fraccari, A., & Galatulas, I. (1998) Synthesis and cytotoxic activity of new peptides containing basic amino acid residues. Anticancer Res., 18, 757-758. #11547
- 594. Chin, J., Fell, J.B., Jarosinski, M., Shapiro, M.J., & Wareing, J.R. (1998) HPLC/NMR in combinatorial chemistry. J. Org. Chem., 63, 386-390. #10682
- 595. Cho, C.Y., Youngquist, R.S., Paikoff, S.J., Beresini, M.H., Hebert, A.R., Berleau, L.T., Liu, C.W., Wemmer, D.E., Keough, T., & Schultz, P.G. (1998) Synthesis and screening of linear and cyclic oligocarbamate libraries: Discovery of high affinity ligands for GPIIb/IIIa. J. Amer. Chem. Soc., 120, 7706-7718. #11266
- Choma, C.T., Robillard, G.T., & Englebretsen, D.R. (1998) Synthesis of hydrophobic peptides: An Fmoc "solubilising tail" method. Tetrahedron Lett., 39, 2417-2420. #10596
- 597. Dallaire, C., & Arya, P. (1998) Synthesis of new building blocks: Towards the analogs of peptide nucleic acids (PNAs). Tetrahedron Lett., 39, 5129-5132. #10892
- 598. Daniels, S.B., Hantman, S.F., Sole, N.A., Gibney, B.R., Rabanal, F. & Kates, S.A. (1998) Pioneer(TM): A continuous-flow peptide synthesis system. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 323-324). Mayflower Scientific Ltd., Kingswinford. #11506

- 599. De Filippis, V., Quarzago, D., Vindigni, A., Dicera, E., & Fontana, A. (1998) Synthesis and characterization of more potent analogues of hirudin fragment 1-47 containing non-natural amino acids. Biochemistry, 37, 13507-13515. #11838
- 600. Dendrinos, K.G., & Kalivretenos, A.G. (1998) Synthesis of N-hydroxysuccinimide esters using polymer bound HOBT.

  Tetrahedron Lett., 39, 1321-1324. #10512
- 601. Dettin, M., Scarinci, C., Zanotto, C., Roncon, R., De Rossi, A., & Di Bello, C. (1998) Biological and conformational studies on analogues of a synthetic peptide enhancing HIV-1 infection. J. Peptide Sci., 4, 436-448. #12342
- 602. Dettin, M., Scarinci, C., Zanotto, C., Cabrelle, A., De Rossi, A., & Di Bello, C. (1998) Design, synthesis and CD4 binding studies of a fluorescent analogue of a peptide that enhances HIV-1 infectivity. Int. J. Peptide Prot. Res., 51, 110-115. #10731
- 603. Di Fenza, A., Tancredi, M., Galoppini, C., & Rovero, P. (1998) Racemization studies of Fmoc-Ser(tBu)-OH during stepwise continuous-flow solid-phase peptide synthesis. Tetrahedron Lett., 39, 8529-8532. #11945
- 604. Di Modugno, F., Rosano, L., Castelli, M., & Chersi, A. (1998) Isolation, characterization and comparison of antipeptide and antiprotein rabbit antibodies to the pi-isoform of glutathione S-transferase. Zeitschrift für Naturforschung Section C-A Journal of Biosciences, 53, 902-910. #12343
- 605. Dick, F., Fritschi, U., Haas, G., Hassler, O., Nyfeler, R. & Rapp, E. (1998) Solid phase synthesis of cyclic peptides using Fmoc strategy. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 339-340). Mayflower Scientific Ltd., Kingswinford. #11556
- 606. Dittrich, F., Tegge, W., & Frank, R. (1998) "Cut and combine": An easy membrane-supported combinatorial synthesis technique. Bioorg. Med. Chem. Lett., 8, 2351-2356. #11206
- 607. Dominguez, E., O'Donnell, M.J., & Scott, W.L. (1998) Solid-phase synthesis of substituted glutamic acid derivatives via Michael addition reactions. Tetrahedron Lett., 39, 2167-2170. #10601
- 608. Dooley, C.T., & Houghten, R.A. (1998) Synthesis and screening of positional scanning combinatorial libraries. Methods Mol. Biol., 87, 13-24. #10564
- Dooley, C.T., Ny, P., Bidlack, J.M., & Houghten, R.A. (1998) Selective ligands for the mu, delta, and kappa opioid receptors identified from a single mixture based tetrapeptide positional scanning combinatorial library. Journal of Biological Chemistry, 273, 18848-18856. #11274
- 610. Dorff, P.H., Chiu, G., Goldstein, S.W., & Morgan, B.P. (1998) Solid phase synthesis of phosphinopeptoids as transition state analog inhibitors. Tetrahedron Lett., 39, 3375-3378. #10803
- 611. Drakopoulou, E., Vizzavona, J., Neyton, J., Aniort, V., Bouet, F., Virelizier, H., Menez, A., & Vita, C. (1998) Consequence of the removal of evolutionary conserved disulfide bridges on the structure and function of charybdotoxin and evidence that particular cysteine spacings govern specific disulfide bond formation. Biochemistry, 37, 1292-1301. #10586
- 612. Efimov, V.A., Choob, M.V., Buryakova, A.A., Kalinkina, A.L., & Chakhmakhcheva, O.G. (1998) Synthesis and evaluation of some properties of chimeric oligomers containing PNA and phosphono-PNA residues. Nucl. Acid. Res., 26, 566-575. #10736
- 613. Eggenweiler, H.M., Clausen, N. & Bayer, E. (1998) A new base-labile linker for Boc solid phase peptide synthesis. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 359-360). Mayflower Scientific Ltd., Kingswinford. #11510
- 614. El-Faham, A. (1998) Bis(tetramethylene)fluoroformamidinium hexafluorophosphate(BTFFK): A convenient coupling reagent for solid phase peptide synthesis. Chem. Lett., 671-672. #11276
- 615. Englebretsen, D.R., Choma, C.T., & Robillard, G.T. (1998) Synthesis of a designed transmembrane protein by thioether ligation of solubilised segments: N-alpha-haloacetylated peptides survived resin cleavage using TFA with EDT as scavenger. Tetrahedron Lett., 39, 4929-4932. #10936
- 616. Erlandsson, M. & Und n, A. (1998) Reduction of aromatic nitro groups in peptides on solid phase. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 367-368). Mayflower Scientific Ltd., Kingswinford. #11557

- 617. Fauq, A.H., Hong, F., Cusack, B., Tyler, B.M., Ping-Pang, Y., & Richelson, E. (1998) Synthesis of (2S)-2-amino-3-(1H-4-indolyl) propanoic acid, a novel tryptophan analog for structural modification of bioactive peptides. Tetrahedron-Asymmetry, 9, 4127-4134. #12352
- 618. Fehrentz, J.A., Paris, M., Heitz, A., Velek, J., Winternitz, F. & Martinez, J. (1998) Solid phase synthesis of C-terminal peptide aldehydes: Study on two different linkers and on C-terminal residue epimerisation. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 375-376). Mayflower Scientific Ltd., Kingswinford. #11558
- 619. Feiertag, S., Wiesmuller, K.H. & Jung, G. (1998) Combinatorial head-to-tail cyclized peptide libraries: Evolution of synthestic and analytical methods based on 517 cyclohexapeptides. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 157-160). Mayflower Scientific Ltd., Kingswinford. #11497
- 620. Fernandez-Garcia, C., Prager, K., McKervey, M.A., Walker, B., & Williams, C.H. (1998) Solid phase synthesis of N-carboxy alkyl-containing peptides derived from enantiopure alpha-keto-beta-aminoacids. Bioorg. Med. Chem. Lett., 8, 433-436. #10692
- 621. Fields, G.B., Lauer, J.L., Dori, Y., Forns, P., Yu, Y.C., & Tirrell, M. (1998) Proteinlike molecular architecture: Biomaterial applications for inducing cellular receptor binding and signal transduction. Biopolymers (Pept. Sci.), 47, 143-151. #10665
- 622. Fletcher, M.D., & Campbell, M.M. (1998) Partially modified retro-inverso peptides: Development, synthesis, and conformational behavior. Chem. Rev., 98, 763-795. #11003
- 623. Fretz, H. (1998) O-(Carboxydifluoromethyl)-L-tyrosine: Design and synthesis of a novel non phosphorous-containing phosphotyrosine isostere. Tetrahedron, 54, 4849-4858. #10998
- 624. Friedler, A., Zakai, N., Karni, O., Broder, Y.C., Baraz, L., Kotler, M., Loyter, A., & Gilon, C. (1998) Backbone cyclic peptide, which mimics the nuclear localization signal of human immunodeficiency virus type 1 matrix protein, inhibits nuclear import and virus production in nondividing cells. Biochemistry, 37, 5616-5622. #11102
- 625. Fu, J.M., & Castelhano, A.L. (1998) Design and synthesis of a pyridone-based phosphotyrosine mimetic. Bioorg. Med. Chem. Lett., 8, 2813-2816. #12355
- 626. Fukuda, H., Irie, K., Nakahara, A., Oie, K., Ohigashi, H., & Wender, P.A. (1998) Synthesis and binding studies of the 116-mer peptide containing the double cysteine-rich motifs of protein kinase C gamma. Tetrahedron Lett., 39, 7943-7946. #11805
- 627. Funakoshi, S., Murayama, E., Guo, L., Fujii, N., & Yajima, H. (1998) A modified benzhydrylamine as a handle reagent for the solid phase synthesis of peptide amides based on the fluorenylmethoxycarbonyl method. J. Chem. Soc. Chem. Commun., 382-384. #11957
- 628. Furet, P., Gay, B., Caravatti, G., Garcia-Echeverria, C., Rahuel, J., Schoepfer, J., & Fretz, H. (1998) Structure-based design and synthesis of nigh affinity tripeptide ligands of the Grb2-SH2 domain. J. Med. Chem., 41, 3442-3449. #11846
- 629. Furlan, R.L.E., & Mata, E.G. (1998) Transprotection in solid phase peptide synthesis: Easy conversion of Fmoc carbamates into Boc carbamates. Tetrahedron Lett., 39, 6421-6422. #11230
- 630. Futaki, S. (1998) Peptide synthesis aiming at elucidation and creation of protein functions [Japanese]. Yakugaku Zasshi Journal of the Pharmaceutical Society of Japan, 118, 493-510. #12356
- 631. Garzotti, M., Rovatti, L., & Hamdan, M. (1998) Investigation of bombolitin complexes with mercury by liquid chromatography tandem mass spectrometry. Rapid Commun. Mass Spectrom., 12, 61-68. #10186
- 632. Garzotti, M., & Hamdan, M. (1998) Liquid chromatography/tandem mass spectrometry of synthesis products associated with the viral protein US11. Rapid Commun. Mass Spectrom., 12, 843-848. #11009
- 633. Gennari, C., Gude, M., Potenza, D., & Piarulli, U. (1998) Hydrogen-bonding donor/acceptor scales in beta-sulfonamidopeptides. Chem. Eur. J., 4, 1924-1931. #11848
- 634. Gilon, C., Huenges, M., Matha, B., Gellerman, G., Hornik, V., Afargan, M., Amitay, O., Ziv, O., Feller, E., Gamliel, A., Shohat, D., Wanger, M., Arad, O., & Kessler, H. (1998) A backbone-cyclic, receptor 5-selective somatstatin analogue: Synthesis, bioactivity, and nuclear magnetic resonance conformational analysis. J. Med. Chem., 41, 919-929. #11541

- 635. Goodman, M., Bhumralkar, M., Jefferson, E.A., Kwak, J., & Locardi, E. (1998) Collagen mimetics. Biopolymers (Pept. Sci. ), 47, 127-142. #10701
- 636. Goodwin, T.E., Holland, R.D., Lay, J.O.Jr., & Raney, K.D. (1998) A simple procedure for solid-phase synthesis of peptide nucleic acids with N-terminal cysteine. Bioorg. Med. Chem. Lett., 8, 2231-2234. #11188
  - 637. Greenberg, W.A., Baird, E.E., & Dervan, P.B. (1998) A comparison of H-pin and hairpin polyamide motifs for the recognition of the minor groove of DNA. Chem. Eur. J., 4, 796-805. #10949
  - 638. Guibe, F. (1998) Allylic protecting groups and their use in a complex environment. Part II. Allylic protecting groups and their removal through catalytic palladium pi-allyl methodology [review]. Tetrahedron, 54, 2967-3042. #10655
  - 639. Guichard, G., Abele, S., & Seebach, D. (1998) Preparation of N-Fmoc-protected beta(2)- and beta(3)-amino acids and their use as building blocks for the solid-phase synthesis of beta-peptides. Helv. Chim. Acta, 81, 187-206. #10669
  - 640. Gulyas, E.C., Soos, K., Varga, J., Toth, G., & Penke, B. (1998) Synthesis of tritium-labelled beta-amyloid fragments. J. Label. Compound. Radiopharm., 41, 763-771. #11291
  - 641. Haas, K., Ponikwar, W., Noth, H., & Beck, W. (1998) Facile synthesis of cyclic tetrapeptides from nonactivated peptide esters on metal centers. Angew. Chem. Int. Ed., 37, 1086-1089. #10974
  - 642. Habermann, J., & Kunz, H. (1998) Fragment condensation on solid-phase in the synthesis of an amphiphilic glycopeptide from the homophilic recognition domain of epithelial cadherin 1. Tetrahedron Lett., 39, 4797-4800. #10934
  - 643. Habermann, J., & Kunz, H. (1998) Glycopeptide synthesis using O-pentafluorophenyluronium salts as novel condensing reagents. Journal fur Praktische Chemie-Chemiker-Zeitung, 340, 233-239. #11000
  - 644. Habermann, J., & Kunz, H. (1998) Solid-phase synthesis of a glycopeptide from the homophilic recognition domain of epithelial cadherin 1 using a O-pentafluorophenyluronium salt. Tetrahedron Lett., 39, 265-268. #9873
  - 645. Hackeng, T.M., Dawson, P.E., Kent, S.B.H., & Griffin, J.H. (1998) Chemical synthesis of human protein s thrombin-sensitive module and first epidermal growth factor module. Biopolymers, 46, 53-63. #11067
  - 646. Hall, B.J., & Sutherland, J.D. (1998) A practical method for the combinatorial synthesis of peptide aldehydes. Tetrahedron Lett., 39, 6593-6596. #11232
  - 647. Hawthorne, S.J., Pagano, M., Harriott, P., Halton, D.W., & Walker, B. (1998) The synthesis and utilization of 2,4-dinitrolphenyl-labeled irreversible peptidyl diazomethyl ketone inhibitors. Anal. Biochem., 261, 131-138. #11852
  - 648. Hemmer, B., Vergelli, M., Pinilla, C., Houghten, R., & Martin, R. (1998) Probing degeneracy in T-cell recognition using peptide combinatorial libraries. Immunol. Today, 19, 163-168. #11151
  - 649. Hemmer, B., Vergelli, M., Gran, B., Ling, N., Conlon, P., Pinilla, C., Houghten, R., McFarland, H.F., & Martin, R. (1998) Cutting edge: Predictable TCR antigen recognition based on peptide scans leads to the identification of agonist ligands with no sequence homology. Journal of Immunology, 160, 3631-3636. #11692
  - 650. Henkel, B., & Bayer, E. (1998) Monitoring of solid phase peptide synthesis by FT-IR spectroscopy. J. Peptide Sci., 4, 461-470. #12369
  - 651. Hirschmann, R. (1998) Peptide related research: A means to further biological and chemical understanding. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 3-17). Mayflower Scientific Ltd., Kingswinford. #11476
  - 652. Hirschmann, R., Yao, W., Arison, B., Maechler, L., Rosegay, A., Sprengeler, P.A., & Smith, A.B.I. (1998) Synthesis of the first tricyclic homodetic peptide: Use of coordinated orthogonal deprotection to achieve directed ring closure. Tetrahedron, 54, 7179-7202. #10948
  - 653. Hlavacek, J., Tykva, R., Bennettova, B., & Barth, T. (1998) The C-terminus shortened analogs of the insect peptide oostatic hormone with accelerated activity. Bioorg. Chem., 26, 131-140. #12370

- 654. Holm, J., Linse, S., & Kihlberg, J. (1998) Synthesis of an N-linked glycopeptide from vitamin K-dependent protein S. Tetrahedron, 54, 11995-12006. #11855
- 655. Hsieh, H.P., Wu, Y.T., Chen, S.T., & Wang, K.T. (1998) Dihydropyran-2-carboxylic acid, a novel bifunctional linker for the solid-phase synthesis of peptides containing a C-terminal alcohol. Chem. Commun., 649-650. #11030
- 656. Inazu, T., Haneda, K., & Mizuno, M. (1998) Synthetic study on N-glycopeptides. J. Syn. Org. Chem. Jpn., 56, 210-220. #11048
- 657. Jacobs, J.W., Patel, D.V., Yuan, Z., Holmes, C.P., Schullek, J., Antonenko, V.V., Grove, J.R., Kulikov, N., Maclean, D., Navre, M., Nguyen, C., Shi, L., Sundaram, A. & Sundberg, S.A. (1998) Light-directed chemical synthesis of positionally encoded peptide arrays. In E.M. Gordon & J.F.J. Kerwin (Eds.), Combinatorial Chemistry and Molecular Diversity in Drug Discovery. (pp. 111-131). John Wiley & Sons, New York. #11710
- 658. Jaskiewicz, A., Lesner, A., Rozycki, J., Kupryszewski, G., & Rolka, K. (1998) Analogue of Cucurbita maxima trypsin inhibitor III (CMTI-III) with all L-cysteine residues substituted by L-penicillamine (Pen). Synthesis and evaluation of trypsin inhibitory activity. Polish J. Chem., 72, 2537-2540. #12379
- 659. Jensen, K.J., Alsina, J., Songster, M.F., Vagner, J., Albericio, F., & Barany, G. (1998) Backbone amide linker (BAL) strategy for solid-phase synthesis of C-terminal-modified and cyclic peptides. J. Amer. Chem. Soc., 120, 5441-5452. #10931
- 660. Jezek, J., Velek, J., Trnka, T., Pisacka, M. & Marecek, F. (1998) Solid phase synthesis of glycopeptide dendrimers with Tn antigenic structure and their biological activities. In R. Ramage & R. Epton (Eds.), Peptides 1996. (pp. 503-504). Mayflower, London. #14411
- 661. Jiang, L., Davison, A., Tennant, G., & Ramage, R. (1998) Synthesis and application of a novel coupling reagent, ethyl 1-hydroxy-1H-1,2,3-triazole-4-carboxylate. Tetrahedron, 54, 14233-14254. #12380
- 662. Jones, D.S., Gamino, C.A., & Randow, M.E. (1998) Synthesis of a cyclic-thioether peptide which binds anti-cardiolipin antibodies. Tetrahedron Lett., 39, 6107-6110. #11023
- 663. Jonsson, D., Molin, H., & Und n, A. (1998) Solid phase synthesis of tropane derivatives. Tetrahedron Lett., 39, 1059-1062. #10678
- 664. Kaplan, B.E., Hefta, L.J., Blake, R.C., II, Swiderek, K.M., & Shively, J.E. (1998) Solid-phase synthesis and characterization of carcinoembryonic antigen (CEA) domains. J. Pept. Res., 52, 249-260. #12165
- Karawajczyk, B., Wysocki, J., Kunikowska, D., Mackiewicz, Z., Glosnicka, R., Korzeniowski, A., Gorski, J., & Kupryszewski, G. (1998) Synthesis of human heat shock protein 70 (29-42) fragment and evaluation of its immunogenicity. Polish J. Chem., 72, 1017-1020. #10944
- 666. Karawajczyk, B., Mackiewicz, Z., Kunikowska, D., Dziadziuszko, H., Deratomaszewska, B., Glosnicka, R., & Kupryszewski, G. (1998) Synthesis of hepatitis c virus protein fragments and evaluation of their immunogenicity. Polish J. Chem., 72, 84-88. #10497
- 667. Karigiannis, G., Mamos, P., Balayiannis, G., Katsoulis, I., & Papaioannou, D. (1998) Simple fragment syntheses of all four isomers of the spermine alkaloid kukoamine. Tetrahedron Lett., 39, 5117-5120. #10923
- Karlsson, K.F., Walse, B., Drakenberg, T., Roy, S., Bergquist, K.E., Pinkner, J.S., Hultgren, S.J., & Kihlberg, J. (1998) Binding of peptides in solution by the Escherichia coli chaperone PapD as revealed using an inhibition ELISA and NMR spectroscopy. Bioorg. Med. Chem., 6, 2085-2101. #12381
- 669. Kassim, S.Y., Restrepo, I.M., & Kalivretenos, A.G. (1998) Synthesis and purification of hydrophobic peptides for use in biomimetic ion channels. J. Chromatogr., A., 11-20. #11299
- 670. Kawakami, T., Yoshimura, S., & Aimoto, S. (1998) Synthesis of reaper, a cysteine-containing polypeptide, using a peptide thioester in the presence of silver chloride as an activator. Tetrahedron Lett., 39, 7901-7904. #11804
- 671. Kellam, B., Drouillat, B., Dekany, G., Starr, M.S., & Toth, I. (1998) Synthesis and in vitro evaluation of lipoamino acid and carbohydrate-modified enkephalins as potential antinociceptive agents. Int. J. Pharm., 161, 55-64. #10699
- 672. Kent, S.B.H., Bark, S.J., Canne, L.E., Dawson, P.E., Fitzgerald, M.C., Hackeng, T., Lu, W. & Muir, T. (1998) Total chemical synthesis of proteins as a probe of structure and function. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the

- Twenty-Fourth European Peptide Symposium. (pp. 187-192). Mayflower Scientific Ltd., Kingswinford. #11554
- 673. Klugmann, K., Kunikowska, D., Glosnicka, R., & Mackiewicz, Z. (1998) Synthesis of flagellin fragments and studies of their interactions with antibodies. Part I. Polish J. Chem., 72, 2093-2097. #11868
- 674. Ko, D.H., Kim, D.J., Lyu, C.S., Min, I.K., & Moon, H.S. (1998) New cleavage approaches to combinatorial synthesis of homoserine lactones. Tetrahedron Lett., 39, 297-300. #9875
- 675. Koerber, S.C., Gulyas, J., Lahrichi, S.L., Corrigan, A., Craig, A.G., Rivier, C., Vale, W., & Rivier, J. (1998) Constrained corticotropin-releasing factor (CRF) agonists and antagonists with i-(i+3) Glu-Xaa-DXbb-Lys bridges. J. Med. Chem., 41, 5002-5011. #12385
- 676. Kohmura, M., & Ariyoshi, Y. (1998) Chemical synthesis and characterization of the sweet protein mabinlin II. Biopolymers, 46, 215-223. #11545
- 677. Konno, K., Hisada, M., Itagaki, Y., Naoki, H., Kawai, N., Miwa, A., Yasuhara, T., & Takayama, H. (1998) Isolation and structure of pompilidotoxins, novel peptide neurotoxins in solitary wasp venoms. Biochem. Biophys. Res. Commun., 250, 612-616. #11869
- 678. Kosaganov, Y.N., Stetsenko, D.A., Lubyako, E.N., Kvitko, N.P., & Lazurkin, Y.S. (1998) Stability of DNA complexes with peptide-nucleic acid. Mol. Biol., 32, 105-108. #11174
- 679. Krambovitis, E., Hatzidakis, G., & Barlos, K. (1998) Preparation of MUC-1 oligomers using an improved convergent solid-phase peptide synthesis. Journal of Biological Chemistry, 273, 10874-10879. #11119
- 680. Kruijtzer, J.A.W., Hofmeyer, L.J.F., Heerma, W., Versluis, C., & Liskamp, R.M.J. (1998) Solid-phase syntheses of peptoids using Fmoc-protected N-substituted glycines: The synthesis of (retro)peptoids of Leu-enkephalin and substance P. Chem. Eur. J., 4, 1570-1580. #11305
- 681. Kudryavtseva, E.V., Sidorova, M.V., & Evstigneeva, R.P. (1998) Peculiarities of synthesis of the cysteine-containing peptides. Uspekhi Khimii, 67, 611-630. #11873
- 682. Kundu, B., Bauser, M., Betschinger, J., Kraas, W., & Jung, G. (1998) Identification of a potent analogue of nazumamide a through iteration of combinatorial tetrapeptide libraries. Bioorg. Med. Chem. Lett., 8, 1669-1672. #11183
- 683. Lam, K.S., Sroka, T., Chen, M.L., Zhao, Y., Lou, Q., Wu, J.Z., & Zhao, Z.G. (1998) Application of one-bead one-compound combinatorial library methods in signal transduction research. Life Sci., 62, 1577-1583. #11160
- 684. Lam, K.S. (1998) Determination of peptide substrate motifs for protein kinases using a "one-bead one-compound" combinatorial library approach. Methods Mol. Biol., 87, 83-86. #11535
- 685. Lam, K.S., & Lebl, M. (1998) Synthesis of a one-bead one-compound combinatorial peptide library. Methods Mol. Biol., 87, 1-6. #11470
- 686. Lang, I., Donze, N., Garrouste, P., Dumy, P., & Mutter, M. (1998) Chemoselectively addressable HCan building blocks in peptide synthesis: L-homocanaline derivatives. J. Peptide Sci., 4, 72-80. #11144
- 687. Lange, M., Cuthbertson, A.S., Towart, R., & Fischer, P.M. (1998) Synthesis and activity of dimeric bradykinin antagonists containing diaminodicarboxylic acid bridge residues. J. Peptide Sci., 4, 289-293. #11309
- 688. Larsen, B.D., & Holm, A. (1998) Sequence-assisted peptide synthesis (SAPS). J. Pept. Res., 52, 470-476. #12394
- 689. Larsen, B.D. & Holm, A. (1998) Sequence assisted peptide synthesis (SAPP). In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 567-568). Mayflower Scientific Ltd., Kingswinford. #11559
- 690. Lebl, M. (1998) Solid-phase synthesis on planar supports. Biopolymers (Pept. Sci.), 47, 397-404. #12265
- 691. Leger, R., Yen, R., She, M.W., Lee, V.J., & Hecker, S.J. (1998) N-Linked solid phase peptide synthesis. Tetrahedron Lett., 39, 4171-4174. #10853
- 692. Lehmann, J., Linden, A., & Heimgartner, H. (1998) Synthesis of the endothiopeptide Boc-Trp-lle-Ala-Aib-Ile-Val-psi[CSNH]Aib-

- Leu-Aib-Pro-OMe by a variation of the 'azirine/oxazolone method'. Tetrahedron, 54, 8721-8736. #11311
- 693. Lelievre, D., Chabane, H., & Delmas, A. (1998) Simple and efficient solid-phase synthesis of unprotected peptide aldehyde for peptide segment ligation. Tetrahedron Lett., 39, 9675-9678. #12055
- 694. Lescrinier, T., Hendrix, C., Kerremans, L., Rozenski, J., Link, A., Samyn, P., Vanaerschot, A., Lescrinier, E., Eritja, R., Vanbeeumen, J., & Herdewijn, P. (1998) DNA-binding ligands from peptide libraries containing unnatural amino acids. Chem. Eur. J., 4, 425-433. #11019
- 695. Li, X., Kawakami, T., & Aimoto, S. (1998) Direct preparation of peptide thioesters using an Fmoc solid-phase method. Tetrahedron Lett., 39, 8669-8672. #11961
- 696. Limal, D., Quesnel, A., & Briand, J.P. (1998) Direct synthesis of N-protected beta-amino dimethylhydroxamates: Application to the solid-phase synthesis of a peptide incorporating a new amide bond surrogate psi[CH2CH2NH]. Tetrahedron Lett., 39, 4239-4242. #10857
- 697. Limal, D., Briand, J.P., Dalbon, P., & Jolivet, M. (1998) Solid-phase synthesis and on-resin cyclization of a disulfide bond peptide and lactam analogues corresponding to the major antigenic site of HIV gp41 protein. J. Pept. Res., 52, 121-129. #11313
- 698. Lin, P., & Ganesan, A. (1998) Solid-phase synthesis of peptidomimetic oligomers with a phosphodiester backbone. Bioorg. Med. Chem. Lett., 8, 511-514. #10693
- 699. Link, A., Lescrinier, T., Kerremans, L., Rozenski, J., & Herdewijn, P. (1998) Side reactions in solid phase peptide synthesis caused by sulfonyl-ornithine building blocks. Pharmazie, 53, 200-201. #11161
- Lohse, P.A., & Felber, R. (1998) Incorporation of a phosphonic acid isostere of aspartic acid into peptides using Fmoc-solid phase synthesis. Tetrahedron Lett., 39, 2067-2070. #10599
- Loughnan, M., Bond, T., Atkins, A., Cuevas, J., Adams, D.J., Broxton, N.M., Livett, B.G., Down, J.G., Jones, A., Alewood, P.F., & Lewis, R.J. (1998) Alpha-conotoxin Epl, a novel sulfated peptide from conus episcopatus that selectively targets neuronal nicotinic acetylcholine receptors. Journal of Biological Chemistry, 273, 15667-15674. #11069
- Lowik, D.W.P.M., Weingarten, M.D., Broekema, M., Brouwer, A.J., Still, W.C., & Liskamp, R.M.J. (1998) Tweezers with different bite: Increasing the affinity of synthetic receptors by varying the hinge part. Angew. Chem. Int. Ed., 37, 1846-1850. #11316
- 703. Lozano, J.M., Espejo, F., Diaz, D., Salazar, L.M., Rodriguez, J., Pinzon, C., Calvo, J.C., Guzman, F., & Patarroyo, M.E. (1998) Reduced amide pseudopeptide analogues of a malaria peptide possess secondary structural elements responsible for induction of functional antibodies which react with native proteins expressed in Plasmodium falciparum erythrocyte stages. J. Pept. Res., 52, 457-469. #12400
- Lu, W.Y., Starovasnik, M.A., & Kent, S.B.H. (1998) Total chemical synthesis of bovine pancreatic trypsin inhibitor by native chemical ligation. FEBS Lett., 429, 31-35. #11071
- Lumma, W.C.Jr., Witherup, K.M., Tucker, T.J., Brady, S.F., Sisko, J.T., Naylor-Olsen, A.M., Lewis, S.D., Lucas, B.J., & Vacca, J.P. (1998) Design of novel, potent, noncovalent inhibitors of thrombin with nonbasic P-1 substructures: Rapid structure-activity studies by solid-phase synthesis. J. Med. Chem., 41, 1011-1013. #10591
- 706. Lutsiak, C.M.E., Sosnowski, D.L., Wishart, D.S., Kwon, G.S., & Samuel, J. (1998) Use of a liposome antigen delivery system to alter immune responses in vivo. J. Pharm. Sci., 87, 1428-1432. #12401
- Maeda, M., Kawasaki, K., Mu, Y., Kamada, H., Tsutsumi, Y., Smith, T.J., & Mayumi, T. (1998) Amino acids and peptides.
   XXXIII. A bifunctional poly(ethylene glycol) hybrid of laminin-related peptides. Biochem. Biophys. Res. Commun., 248, 485-489. #11317
- 708. Maeda, M., Izuno, Y., Kawasaki, K., Kaneda, Y., Mu, Y., Tsutsumi, Y., Nakagawa, S., & Mayumi, T. (1998) Amino acids and peptides. XXXI. Preparation of analogs of the laminin-related peptide YIGSR and their inhibitory effect on experimental metastasis. Chem. Pharm. Bull. Tokyo, 46, 347-350. #10706
- 709. Magerlein, M., Hock, D., Adermann, K., Neidlein, R., Forssmann, W.G., & Strein, K. (1998) Production of sequence specific polyclonal antibodies to human parathyroid hormone 1-37 by immunization with multiple antigenic peptides. Arzneim. -Forsch.,

- 48, 783-787. #11007
- Mah, E., Vossen, P., Van den Hooven, H.W., Le-Nguyen, D., Vervoort, J., & De Wit, P.J.G.M. (1998) Solid-phase synthesis, conformational analysis, and biological activity of AVR9 elcitor peptides of the fungal tomato pathogen Cladosporium fulvum. J. Pept. Res., 52, 482-494. #12173
- 711. Marcaurelle, L.A., & Bertozzi, C.R. (1998) Direct incorporation of unprotected ketone groups into peptides during solid-phase synthesis: Application to the one-step modification of peptides with two different biophysical probes for FRET. Tetrahedron Lett., 39, 7279-7282. #11570
- 712. Martin, L., Cornille, F., Coric, P., Roques, B.P., & Fournie-Zaluski, M.C. (1998) Beta-amino-thiols inhibit the zinc metallopeptidase activity of tetanus toxin light chain. J. Med. Chem., 41, 3450-3460. #11879
- 713. Martin, S.F., Dwyer, M.P., & Lynch, C.L. (1998) Application of AlMe3-mediated amidation reactions to solution phase peptide synthesis. Tetrahedron Lett., 39, 1517-1520. #10520
- 714. Mathe, C., Perigaud, C., Gosselin, G., & Imbach, J.L. (1998) Phosphopeptide prodrug bearing an S-acyl-2-thioethyl enzyme-labile phosphate protection. J. Org. Chem., 63, 8547-8550. #12406
- 715. Matsuura, K., Ikoma, S., Sugiyama, M., Funauchi, M., & Sinohara, H. (1998) Amidolytic and peptidolytic activities of immunoglobulin G present in sera from patients with rheumatoid arthritis, Sjogren's-syndrome and systemic lupus erythematosus. Immunology, 95, 26-30. #11880
- 716. Matysiak, S., Boldicke, T., Tegge, W., & Frank, R. (1998) Evaluation of monomethoxytrityl and dimethoxytrityl as orthogonal amino protecting groups in Fmoc solid phase peptide synthesis. Tetrahedron Lett., 39, 1733-1734. #10607
- 717. Maurer, M.C., Peng, J.L., An, S.S., Trosset, J.Y., Henschen-Edman, A., & Scheraga, H.A. (1998) Structural examination of the influence of phosphorylation on the binding of fibrinopeptide a to bovine thrombin. Biochemistry, 37, 5888-5902. #11106
- 718. Mayer, J.P., Zhang, J., Groeger, S., Liu, C.F., & Jarosinski, M.A. (1998) Lanthionide macrocyclization by in situ activation of serine. J. Pept. Res., 51, 432-436. #10813
- 719. Mayerfligge, P., Volz, J., Kruger, U., Sturm, E., Gernandt, W., Schafer, K.P., & Przybylski, M. (1998) Synthesis and structural characterization of human-identical lung surfactant SP-C protein. J. Peptide Sci., 4, 355-363. #11322
- Meisenbach, M., Echner, H. & Voelter, W. (1998) New 9-substituted xanthenylamine linkers for the solid phase synthesis of protected peptide amides. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 633-634). Mayflower Scientific Ltd., Kingswinford. #11516
- 721. Meldal, M., Renil, M., Juliano, M.A., Jansson, A.M., Meinjohanns, E., Buchardt, J. & Schleyer, A. (1998) Novel PEG-based resins and synthetic methods: Azido acids in SPPS and direct solid phase peptide glycosylations. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 141-152). Mayflower Scientific Ltd., Kingswinford. #11496
- 722. Mellor, S.L., Rose, F., McGuire, C., Spencer, A.E., Smith, E.E. & Chan, W.C. (1998) Synthesis and evaluation of atypical antibacterial hexapeptides: Application of facile solid phase strategies. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 635-636). Mayflower Scientific Ltd., Kingswinford. #11517
- 723. Melnyk, O., Bossus, M., David, D., Rommens, C., & Gras-Masse, H. (1998) Synthesis of lipopeptides using hydrazone chemical ligation. J. Pept. Res., 52, 180-184. #11882
- 724. Melnyk, O., Chaurand, P., Rommens, C., Drobecq, H., Wieruszeski, J.M., Spengler, B., & Gras-Masse, H. (1998) Identification of a sequence-dependent reversible acylation of tosylarginine in a peptidyl-resin reacted with isonicotinyl p-nitrophenylcarbonate. J. Pept. Res., 51, 188-193. #12257
- 725. Melnyk, O., Klinguer, C., Loing, E., Boutillon, C. & Gras-Masse, H. (1998) A versatile method for peptide modifications through hydrazone chemical ligation. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 637-638). Mayflower Scientific Ltd., Kingswinford. #11518
- 726. Mezo, G., Mihala, N., Koczan, G., & Hudecz, F. (1998) Cyclohexyloxycarbonyl based orthogonal solid phase peptide synthesis in Boc chemistry. Tetrahedron, 54, 6757-6766. #10959

- 727. Miller, S.C., & Scanlan, T.S. (1998) oNBS-SPPS: A new method for solid-phase peptide synthesis. J. Amer. Chem. Soc., 120, 2690-2691. #11033
- 728. Millington, C.R., Quarrell, R., & Lowe, G. (1998) Aryl hydrazides as linkers for solid phase synthesis which are cleavable under mild oxidative conditions. Tetrahedron Lett., 39, 7201-7204. #11223
- 729. Miranda, L.P., Jones, A., Meutermans, W.D.F., & Alewood, P.F. (1998) p-Cresol as a reversible acylium ion scavenger in solid-phase peptide synthesis. J. Amer. Chem. Soc., 120, 1410-1420. #10659
- 730. Miyashita, M., Nakamori, T., Murai, T., Miyagawa, H., Akamatsu, M., & Ueno, T. (1998) Facile syntheses of AM-toxins and analogs as cyclic depsipeptides by the solid-phase method. Biosci. Biotechnol. Biochem., 62, 1799-1801. #11883
- 731. Mizuno, M., Muramoto, I., Kawakami, T., Seike, M., Aimoto, S., Haneda, K., & Inazu, T. (1998) A synthesis of a glycopeptide analogue of eel calcitonin. Tetrahedron Lett., 39, 55-58. #9880
- 732. Moyesherman, D., Jin, S., Ham, I., Lim, D.Y., Scholtz, J.M., & Burgess, K. (1998) Conformational preferences of RNAse a C-peptide derivatives containing a highly constrained analogue of phenylalanine. J. Amer. Chem. Soc., 120, 9435-9443. #11884
- 733. Nakahara, Y., Ito, Y., & Ogawa, T. (1998) Solid-phase synthesis of the B-chain of human alpha-2HS glycoprotein. Carbohyd. Res., 309, 287-296. #11328
- 734. Nefzi, A., Dooley, C., Ostresh, J.M., & Houghten, R.A. (1998) Combinatorial chemistry: From peptides and peptidomimetics to small organic and heterocyclic compounds. Bioorg. Med. Chem. Lett., 8, 2273-2278. #11193
- 735. Noda, M. (1998) Facile synthesis of 3-methoxy-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-one: Application to the acid-labile peptide amide linker containing the 10,11-dihydro-5H-dibenzo[a,d]cycloheptene moiety. Chem. Pharm. Bull. Tokyo, 46, 1157-1159. #11330
- 736. Nunami, K., Yamada, M., & Shimizu, R. (1998) Design of novel tripeptides with macrophage migration-enhancing activity. Bioorg. Med. Chem. Lett., 8, 2517-2520. #11888
- 737. Ojima, I., Wang, H., Wang, T., & Ng, E.W. (1998) New approaches to the asymmetric synthesis of dipeptide isosteres via beta-Lactam Synthon Method. Tetrahedron Lett., 39, 923-926. #10486
- 738. Orosz, G., & Kiss, L.P. (1998) Simple and efficient synthesis of 2-chlorotritylchloride resin. Tetrahedron Lett., 39, 3241-3242. #10800
- 739. Ostresh, J.M., Dorner, B., & Houghten, R.A. (1998) Peralkylation. "Libraries from libraries": Chemical transformation of synthetic combinatorial libraries. Methods Mol. Biol., 87, 41-49. #10563
- 740. Ota, M., Shimizu, Y., Tonosaki, K., & Ariyoshi, Y. (1998) Synthesis, characterization, and sweetness: Suppressing activities of gurmarin analogues missing one disulfide bond. Biopolymers, 46, 65-73. #11068
- 741. Palkeeva, M.E., Sidorova, M.V., Molokoedov, A.S., Kuznetsova, T.V., Tishchenko, V.A., Kobylyanskii, A.G., Bespalova, Z.D., Nasonov, E.L., & Evstigneeva, R.P. (1998) The synthesis and biological properties of artificial antigens on the basis of the 280-289 fragment of beta(2)-Glycoprotein I. Bioorganicheskaia Khimiia. 24, 502-508. #11891
- 742. Panke, G., & Frank, R. (1998) Improved preparation of safety-catch linker for the solid phase synthesis of peptide acids finally released into aqueous buffers. Tetrahedron Lett., 39, 17-18. #9879
- 743. Paris, M., Heitz, A., Guerlavais, V., Cristau, M., Fehrentz, J.A., & Martinez, J. (1998) Synthesis of peptide aldehydes on solid support using ozonolysis. Tetrahedron Lett., 39, 7287-7290. #11571
- 744. Paris, M., Pothion, C., Heitz, A., Martinez, J., & Fehrentz, J.A. (1998) Synthesis of N- and side chain protected aspartyl and glutamyl aldehyde derivatives. Reinvestigation of the reduction of Weinreb amides. Tetrahedron Lett., 39, 1341-1344. #10513
- 745. Pascal, R., & Sola, R. (1998) Preservation of the Fmoc protective group under alkaline conditions by using CaCl2. Applications in peptide synthesis. Tetrahedron Lett., 39, 5031-5034. #10921
- 746. Patek, M., Bildstein, S., & Flegelova, Z. (1998) Monitoring solid phase reactions with Ion-selective electrode. Tetrahedron Lett.,

- 39, 753-756. #10049
- 747. Perez, J.A., Canto, J., Reig, F., Perez, J.J., & Haro, I. (1998) Conformational behavior of the HAV-VP3(110-121) peptidic sequence and synthetic analogs in membrane environments studied by CD and computational methods. Biopolymers, 45, 479-492. #11141
- 748. Peukert, S., & Giese, B. (1998) The pivaloylglycol anchor group: A new platform for a photolabile linker in solid-phase synthesis. J. Org. Chem., 63, 9045-9051. #12182
- 749. Peyrottes, S., Mestre, B., Burlina, F., & Gait, M.J. (1998) The synthesis of peptide-oligonucleotide conjugates by a fragment coupling approach. Tetrahedron, 54, 12513-12522. #11896
- 750. Pfeifer, M.E., & Robinson, J.A. (1998) Stabilisation of beta-hairpin conformations in a protein surface mimetic using a bicyclic template derived from (2S,3R,4R)-diaminoproline. Chem. Commun., 1977-1978. #11897
- 751. Phoon, C.W., Oliver, S.F., & Abell, C. (1998) The use of trichloroacetimidate-activated resin for ester formation. Tetrahedron Lett., 39, 7959-7962. #11806
- 752. Pietrzynski, G., Slon-Usakiewicz, J.J. & Konishi, Y. (1998) Peptide synthesis database. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 723-724). Mayflower Scientific Ltd., Kingswinford. #11521
- 753. Polaskova, M.E., Ede, N.J., & Lambert, J.N. (1998) Synthesis of nanotubule-forming cyclic octapeptides via an Fmoc strategy. Aust. J. Chem., 51, 535-540. #11335
- 754. Polverino de Laureto, P., Scaramella, E., De Filippis, V., Marin, O., Doni, M.G., & Fontana, A. (1998) Chemical synthesis and structural characterization of the RGD-protein decorsin: A potent inhibitor of platelet aggregation. Protein Sci., 7, 433-444. #10718
- 755. Quesnel, A., & Briand, J.P. (1998) Incomplete trifluoroacetic acid deprotection of asparagine-trityl-protecting group in the vicinity of a reduced peptide bond. J. Pept. Res., 52, 107-111. #11339
- 756. Quetard, C., Bourgerie, S., Normandsdiqui, N., Mayer, R., Strecker, G., Midoux, P., Roche, A.C., & Monsigny, M. (1998) Novel glycosynthons for glycoconjugate preparation: Oligosaccharylpyroglutamylanilide derivatives. Bioconjug. Chem., 9, 268-276. #11167
- 757. Rabenstein, D.L., Russell, J., & Gu, J. (1998) Characterization of the thiol/disulfide chemistry of peptides corresponding to the 603-609 disulfide loop of the human immunodeficiency virus (HIV) envelope glycoprotein gp41. J. Pept. Res., 51, 437-443. #10814
- 758. Rademaker, G.J., Pergantis, S.A., Bloktip, L., Langridge, J.I., Kleen, A., & Thomasoates, J.E. (1998) Mass spectrometric determination of the sites of O-glycan attachment with low picomolar sensitivity. Anal. Biochem., 257, 149-160. #11169
- 759. Ramage, R., & Epton, R. (1998) Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. Mayflower Scientific Ltd., Kingswinford. ISBN 095270112X #11474
- 760. Ramage, R., Swenson, H.R., & Shaw, K.T. (1998) A novel purification protocol for soluble combinatorial peptide libraries. Tetrahedron Lett., 39, 8715-9718. #11964
- 761. Raman, P., Stokes, S.S., Angell, Y.M., Flentke, G.R., & Rich, D.H. (1998) Methods to circumvent a difficult coupling in the solid-phase synthesis of cyclosporine analogues. J. Org. Chem., 63, 5734-5735. #11902
- 762. Rau, H.K., & Haehnel, W. (1998) Design, synthesis, and properties of a novel cytochrome B model. J. Amer. Chem. Soc., 120, 468-476. #10687
- 763. Reichwein, J.F., & Liskamp, R.M.J. (1998) Site-specific N-alkylation of peptides on the solid phase. Tetrahedron Lett., 39, 1243-1246. #10506
- Reis, C.A., Sorensen, T., Mandel, U., David, L., Mirgorodskaya, E., Roepstorff, P., Kihlberg, J., Hansen, J.E.S., & Clausen, H. (1998) Development and characterization of an antibody directed to an alpha-N-acetyl-D-galactosamine glycosylated MUC2 peptide. Glycoconjugate J., 15, 51-62. #10717

- Rene, L., Yaouancq, L., & Badet, B. (1998) Orthogonally protected imidazolidine-2-carboxylic acid, a new proline surrogate suitable for SPPS. Tetrahedron Lett., 39, 2569-2570. #10618
- 766. Renil, M., Ferreras, M., Delaisse, J.M., Foged, N.T., & Meldal, M. (1998) PEGA supports for combinatorial peptide synthesis and solid-phase enzymatic library assays. J. Peptide Sci., 4, 195-210. #10862
- 767. Renil, M., Meldal, M., Delaisse, J.M. & Foged, N.T. (1998) Fluorescent quenched peptide libaries as a tool for identification of enzyme substrates for matrix metalloproteinase (MMP)-9 from osteoclasts. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 753-754). Mayflower Scientific Ltd., Kingswinford. #11523
- 768. Revesz, L., Bonne, F., Manning, U., & Zuber, J.F. (1998) Solid phase synthesis of a biased mini tetrapeptoid-library for the discovery of monodentate itam mimics as ZAP-70 inhibitors. Bioorg. Med. Chem. Lett., 8, 405-408. #10691
- 769. Ridge, S., & Hettiarachchi, K. (1998) Peptide purity and counter ion determination of bradykinin by high performance liquid chromatography and capillary electrophoresis. J. Chromatogr., 817, 215-222. #11903
- Rivier, J., Lahrichi, S.L., Gulyas, J., Erchegyi, J., Koerber, S.C., Craig, A.G., Corrigan, A., Rivier, C., & Vale, W. (1998) Minimal-size, constrained corticotropin-releasing factor agonists with i-(i+3) Glu-Lys and Lys-Gln bridges. J. Med. Chem., 41, 2614-2620.
   #11342
- 771. Rivier, J., Gulyas, J., Corrigan, A., Martinez, V., Craig, A.G., Tache, Y., Vale, W., & Rivier, C. (1998) Astressin analogues (corticotropin-releasing factor antagonists) with extended duration of action in the rat. J. Med. Chem., 41, 5012-5019. #12424
- 772. Roberge, J.Y., Beebe, X., & Danishefsky, S.J. (1998) Convergent synthesis of N-linked glycopeptides on a solid support. J. Amer. Chem. Soc., 120, 3915-3927. #10988
- 773. Roberts, K.D., Lambert, J.N., Ede, N.J., & Bray, A.M. (1998) Efficient synthesis of thioether-based cyclic peptide libraries. Tetrahedron Lett., 39, 8357-8360. #12017
- 774. Rodriguez, J.G., Martinvillamil, R., & Ramos, S. (1998) Solid phase synthesis of anhydrides and amides: Polymer-assisted reaction in the synthesis of dipeptides. New J. Chem., 22, 865-868. #11906
- 775. Rohwedder, B., Mutti, Y., Dumy, P., & Mutter, M. (1998) Hydrazinolysis of Dde:Complete orthogonality with Aloc protecting groups. Tetrahedron Lett., 39, 1175-1178. #10505
- 776. Romanovskis, P., & Spatola, A.F. (1998) Preparation of head-to-tail cyclic peptides via side-chain attachment: Implications for library synthesis. J. Pept. Res., 52, 356-374. #12184
- 777. Root-Bernstein, R.S., & Holsworth, D.D. (1998) Antisense peptides: A critical mini-review [review]. J. Theor. Biol., 190, 107-119. #10705
- 778. Royo, M., Contreras, M.A., Giralt, E., Albericio, F., & Pons, M. (1998) An easy entry to a new high-symmetry, large molecular framework for molecular recognition studies and de novo protein design: Solvent modulation of the spontaneous formation of a cyclic monomer, dimer, or trimer from a bis-cysteine peptide. J. Amer. Chem. Soc., 120, 6639-6650. #11343
- 779. Saha, S., Waugh, D.J.J., Zhao, P., Abel, P.W., & Smith, D.D. (1998) Role of conformational constrains of position 7 of the disulphide bridge of h-a-CGRP derivatives in their agonist versus antagonist properties. J. Pept. Res., 52, 112-120. #11693
- 780. Satyanarayana, J., Gururaja, T.L., Naganagowda, G.A., Ramasubbu, N., & Levine, M.J. (1998) A concise methodology for the stereoselective synthesis of O-glycosylated amino acid building blocks: Complete H-1 NMR assignments and their application in solid-phase glycopeptide synthesis. J. Pept. Res., 52, 165-179. #11912
- 781. Sauter, G., & Leumann, C. (1998) Distamycin-NA: A DNA analog with an aromatic heterocyclic polyamide backbone. Part 2. Solid-phase synthesis of distamycin-NAs containing the nucleobase uracil: Unexpected solvent participation in the coupling step. Helv. Chim. Acta, 81, 916-931. #10972
- 782. Scheibler, L., Dumy, P., Stamou, D., Duschl, C., Vogel, H., & Mutter, M. (1998) Self-assembling functionalized templates in biosensor technology. Polym. Bull., 40, 151-157. #10653
- 783. Schmidt, B., & Kuhn, C. (1998) Racemic, yet diastereomerically pure azido acids as both gamma-turn and inverse gamma-turn

- mimetics for solid-phase peptide synthesis. Synlett, 1240. #12434
- 784. Sebestyen, F., Szendrei, G., Mak, M., Doda, M., Illyes, E., Szokan, G., Kindla, K., Rapp, W., Szego, P., Campian, E., & Furka, A. (1998) Coloured peptides: Synthesis, properties and use in preparation of peptide sub-library kits. J. Peptide Sci., 4, 294-299. #10880
- 785. Seebach, D., Abele, S., Sifferlen, T., Hanggi, M., Gruner, S., & Seiler, P. (1998) Preparation and structure of beta-peptides consisting of geminally disubstituted beta(2,2)- and beta(3,3)-amino acids: A turn motif for beta-peptides. Helv. Chim. Acta, 81, 2218-2243. #12436
- 786. Seligmann, B., Lebl, M. & Lam, K.S. (1998) Solid-phase peptide synthesis, lead generation, and optimization. In E.M. Gordon & J.F.J. Kerwin (Eds.), Combinatorial Chemistry and Molecular Diversity in Drug Discovery. (pp. 39-109). John Wiley & Sons, New York. #11709
- 787. Servent, D., Mourier, G., Antil, S., & Menez, A. (1998) How do snake curaremimetic toxins discriminate between nicotinic acetylcholine receptor subtypes. Toxicology Letters, 103, 199-203. #12811
- 788. Sheibani, N., & Frazier, W.A. (1998) Direct use of synthetic peptides for antiserum production. BioTechniques, 25, 28-32. #10871
- 789. Shogren-Knaak, M.A., & Imperiali, B. (1998) A reversible affinity tag for the purification of N-glycol capped peptides. Tetrahedron Lett., 39, 8241-8244. #12012
- Sim, M.M., Lee, C.L., & Ganesan, A. (1998) Solid-phase C-acylation of active methylene compounds. Tetrahedron Lett., 39, 2195-2198. #10602
- Sintihaki, E., Minakaki, P., Tzougraki, C., Loukas, S., & Stassinopoulou, C.I. (1998) Solid-phase synthesis and structural studies of the N-terminal 25 peptides of ribosomal S2 proteins from Th.thermophilus and E.coli. J. Protein Chem., 17, 551-551. #11350
- 792. Sivanandaiah, K.M., Babu, V.V.S., & Shankaramma, S.C. (1998) Solid phase synthesis of O-glycoopioid peptides related to dermorphin. Ind. J. Chem. B-Org. Chem. Med. Chem., 37, 760-767. #12444
- 793. Spetzler, J.C., Westphal, V., Winther, J.B., & Meldal, M. (1998) Preparation of fluorescence quenched libraries containing interchain disulphide bonds for studies of protein disulphide isomerases. J. Peptide Sci., 4, 128-137. #10628
- 794. St.Hilaire, P.M., Lowary, T., Meldal, M. & Bock, K. (1998) Synthesis and analysis of glycopeptide libraries and their application in solid phase assays of carbohydrate binding-proteins. In R. Ramage & R. Epton (Eds.), Peptides 1996: Proceedings of the Twenty-Fourth European Peptide Symposium. (pp. 817-818). Mayflower Scientific Ltd., Kingswinford. #11525
- 795. Suarez, E., De, E., Molle, G., Lazaro, R., & Viallefont, P. (1998) Synthesis and characterization of a new biotinylated gramicidin. J. Peptide Sci., 4, 371-377. #11560
- 796. Svensson, A., Bergquist, K.E., Fex, T., & Kihlberg, J. (1998) Fluorinated linkers for monitoring solid-phase synthesis using gelphase 19F NMR spectroscopy. Tetrahedron Lett., 39, 7193-7196. #11222
- Sweeney, S.M., Guy, C.A., Fields, G.B., & San Antonio, J.D. (1998) Defining the domains of type I collagen involved in heparinbinding and endothelial tube formation. Proc. Natl. Acad. Sci. USA, 95, 7275-7280. #11057
- 798. Szabo, L., Smith, B.L., Mcreynolds, K.D., Parrill, A.L., Morris, E.R., & Gervay, J. (1998) Solid phase synthesis and secondary structural studies of (1->5) amide-linked sialooligomers. J. Org. Chem., 63, 1074-1078. #10661
- 799. Tam, J.P., & Lu, Y.A. (1998) A biomimetic strategy in the synthesis and fragmentation of cyclic protein. Protein Sci., 7, 1583-1592. #11358
- 800. Tamamura, H., Arakaki, R., Funakoshi, H., Imai, M., Otaka, A., Ibuka, T., Nakashima, H., Murakami, T., Waki, M., Matsumoto, A., Yamamoto, N., & Fujii, N. (1998) Effective lowly cytotoxic analogs of an HIV-cell fusion inhibitor, T22 ([Tyr(5,12),Lys(7)]-polyphemusin II). Bioorg. Med. Chem., 6, 231-238. #11154
- 801. Taylor, C.M. (1998) Glycopeptides and glycoproteins: Focus on the glycosidic linkage. Tetrahedron, 54, 11317-11362. #11920
- 802. Tedebark, U., Meldal, M., Panza, L., & Bock, K. (1998) C-Linked glycosyl azido acid in novel solid-phase C-glycopeptide

- synthesis. Tetrahedron Lett., 39, 1815-1818. #10609
- 803. Tetzlaff, C.N., Schwope, I., Bleczinski, C.F., Steinberg, J.A., & Richert, C. (1998) A convenient synthesis of 5'-amino-5'-deoxythymidine and preparation of peptide-DNA hybrids. Tetrahedron Lett., 39, 4215-4218. #10856
- 804. Toniolo, C., Crisma, M., & Formaggio, F. (1998) TOAC, a nitroxide spin-labeled, achiral C(alpha)-tetrasubstituted alpha-amino acid, is an excellent tool in material science and biochemistry. Biopolymers (Pept. Sci.), 47, 153-158. #10666
- 805. Toth, K., Kovacs, M., Zarandi, M., Halmos, G., Groot, K., Nagy, A., Kele, Z., & Schally, A.V. (1998) New analogs of human growth hormone-releasing hormone (1-29) with high and prolonged antagonistic activity. Int. J. Peptide Prot. Res., 51, 134-141. #10733
- 806. Tu, J., Yu, Z., & Chu, Y.H. (1998) Combinatorial search for diagnostic agents: Lyme antibody H9724 as an example. Clin. Chem., 44, 232-238. #10579
- 807. Tuchscherer, G., Lehmann, C., & Mathieu, M. (1998) New protein mimetics: The zinc finger motif as a locked-in tertiary fold. Angew. Chem., 2990-2993. #12457
- 808. Ueki, M., Goto, R., Okumura, J., & Ishii, K. (1998) N,N'-Dialkyldiamide-type phosphate protecting groups for Fmoc synthesis of phosphotyrosine-containing peptides: Optimization of the alkyl group. Bull. Chem. Soc. Jpn., 71, 1887-1898. #11922
- 809. Varkey, J.T., & Pillai, V.N. (1998) Synthesis of thioredoxin partial sequences on 1,6-hexanediol diacrylate (HDODA)-cross-linked polystyrene resin. Int. J. Peptide Prot. Res., 51, 49-54. #10397
- 810. Vilaseca, M., Nicolas, E., Capdevila, F., & Giralt, E. (1998) Reduction of methionine sulfoxide with NH4I/TFA: Compatibility with peptides containing cysteine and aromatic amino acids. Tetrahedron, 54, 15273-15286. #12198
- Virgilio, A.A. & Ellman, J.A. (1998) Conformationally restricted peptide and peptidomimetic libraries. In E.M. Gordon & J.F.J. Kerwin (Eds.), Combinatorial Chemistry and Molecular Diversity in Drug Discovery. (pp. 133-149). John Wiley & Sons, New York. #3765
- 812. Vyle, J.S., Williams, N.H., & Grasby, J.A. (1998) A novel solid support for synthesis of 2',3'-cyclic phosphate terminated oligonucleotides. Tetrahedron Lett., 39, 7975-7978. #11807
- 813. Wang, P., Layfield, R., Landon, M., Mayer, R.J., & Ramage, R. (1998) Transfer active ester condensation: A novel technique for peptide segment coupling. Tetrahedron Lett., 39, 8711-8714. #11963
- 814. Wang, P., Shaw, K.T., Whigham, B., & Ramage, R. (1998) Synthesis of peptide C-terminal derivatives using the transfer active ester condensation technique. Tetrahedron Lett., 39, 8719-8720. #11965
- 815. Wang, R., Hu, X.Y., Ma, Y.P., & Li, D. (1998) Studies on the solution conformation of angiotensin II [Chinese]. Chem. J. Chin. Univ., 19, 1758-1762. #12463
- 816. Wang, X., Liang, S., & Luo, Z. (1998) Chemical synthesis and physiological activity analysis of K3A-HWTX-I: A mutant of huwentoxin-I. Shengming Kexue Yanjiu, 2, 87-92. #11467
- 817. Welch, C.J., Bhat, G., & Protopopova, M.N. (1998) Silica-based solid phase synthesis of chiral stationary phases. Enantiomer, 3, 463-469. #12466
- 818. Wen, J.J., & Crews, C.M. (1998) Towards the semi-synthesis of didemnin M. Solution and solid phase syntheses of the pseudotetrapeptide:pGlu-Gln-psi-[COO]Ala-Pro-OH. Tetrahedron Lett., 39, 779-782. #10051
- 819. Wen, J.J., & Crews, C.M. (1998) Synthesis of 9-fluorenylmethoxycarbonyl-protected amino aldehydes. Tetrahedron-Asymmetry, 9, 1855-1858. #10906
- 820. Wilken, J., & Kent, S. (1998) Chemical protein synthesis. Curr. Opin. Biotechnol., 9, 412-426. #12915
- 821. Williams, L., Kather, K., & Kemp, D.S. (1998) High helicities of Lys-containing, Ala-rich peptides are primarily attributable to a large, context-dependent Lys stabilization. J. Amer. Chem. Soc., 120, 11033-11043. #12468

- 822. Wirkus-Romanowska, I., Ciurak, M., Miecznikowska, H., Rolka, K., Janusz, M., Szymaniec, S., Krukowska, K., Lisowski, J., & Kupryszewski, G. (1998) Cyclic analogues of proline-rich protein fragments. Part I. Synthesis and evaluation of immunotropic activity. Polish J. Chem., 72, 2394-2398. #12470
- 823. Witte, K., Seitz, O., & Wong, C.H. (1998) Solution- and solid-phase synthesis of N-protected glycopeptide esters of the benzyl type as substrates for subtilisin-catalyzed glycopeptide couplings. J. Amer. Chem. Soc., 120, 1979-1989. #11052
- 824. Wojciechowska, I., Kochanska, B., Stelmanska, E., Knap, N., Mackiewicz, Z., & Kupryszewski, G. (1998) Statherin and its shortened analogues. Polish J. Chem., 72, 2098-2102. #11931
- 825. Wu, J.J., Phan, H., & Lam, K.S. (1998) Comparison of the intrinsic kinase activity and substrate specificity of c-Abl and Brc-Abl. Bioorg. Med. Chem. Lett., 8, 2279-2284. #11194
- 826. Yamada, M., Miyajima, T., & Horikawa, H. (1998) Solid-phase synthesis of dehydroalanine derivatives. Tetrahedron Lett., 39, 289-292. #9874
- 827. Yao, S., Ghosh, I., Zutshi, R., & Chmielewski, J. (1998) A self-replicating peptide under ionic control. Angew. Chem. Int. Ed., 37, 478-481. #10652
- 828. Yao, Z.J., Ye, B., Miyoshi, K., Otaka, A., & Burke, T.R. (1998) Preparation of phospho-L-azatyrosine suitably protected for the synthesis of signal transduction related peptides: A correction. Synlett, 428. #10985
- 829. Yokote, K., Hellman, U., Ekman, S., Saito, Y., Ronnstrand, L., Heldin, C.H., & Mori, S. (1998) Identification of Tyr-762 in the platelet-derived growth factor alpha-receptor as the binding site for Crk proteins. Oncogene, 16, 1229-1239. #10695
- 830. Yoshikawa, M., Fujisawa, T., Izumi, J., Kitao, T., & Sakamoto, S. (1998) Molecularly imprinted polymeric membranes involving tetrapeptide EQKL derivatives as chiral-recognition sites toward amino acids. Anal. Chim. Acta, 365, 59-67. #10925
- 831. Yu, L., Lai, Y., Wade, J.V., & Coutts, S.M. (1998) A simple and efficient method for the synthesis of thioether cyclic peptides. Tetrahedron Lett., 39, 6633-6636. #11228
- 832. Yu, Y.C., Tirrell, M., & Fields, G.B. (1998) Minimal lipidation stabilizes protein-like molecular architecture. J. Amer. Chem. Soc., 120, 9979-9987. #11937
- 833. Yu, Z., Yu, X.C., & Chu, Y.H. (1998) MALDI-MS determination of cyclic peptidomimetic sequences on single beads directed toward the generation of libraries. Tetrahedron Lett., 39, 1-4. #9878
- 834. Zhang, A.J., Russell, D.H., Zhu, J., & Burgess, K. (1998) A method of removal of N-Boc protecting groups from substrates on TFA-sensitive resins. Tetrahedron Lett., 39, 7439-7442. #11561
- 835. Zuberi, S., Glen, A., Hider, R.C., & Bansal, S.S. (1998) Synthesis of asymmetric cystines. Tetrahedron Lett., 39, 7567-7570. #11939
- 836. Albericio, F., Lloyd-Williams, P., & Giralt, E. (1997) Convergent solid-phase peptide synthesis. Methods Enzymol., 289, 313-336. #9931
- 837. Albericio, F., & Carpino, L.A. (1997) Coupling reagents and activation. Methods Enzymol., 289, 104-126. #9993
- 838. Albericio, F., Cases, M., Alsina, J., Triolo, S.A., Carpino, L.A., & Kates, S.A. (1997) On the use of PyAOP, a phosphonium salt derived from HOAt, in solid-phase peptide synthesis. Tetrahedron Lett., 38, 4853-4856. #9043
- 839. Alewood, P., Alewood, D., Miranda, L., Love, S., Meutermans, W., & Wilson, D. (1997) Rapid in situ neutralization protocols for Boc and Fmoc solid-phase chemistries. Methods Enzymol., 289, 14-29. #9990
- 840. Alsina, J., Chiva, C., Ortiz, M., Rabanal, F., Giralt, E., & Albericio, F. (1997) Active carbonate resins for solid-phase synthesis through the anchoring of a hydroxyl function. Synthesis of cyclic and alcohol peptides. Tetrahedron Lett., 38, 883-886. #8231
- 841. Ammosova, T.N., Ouporov, I.V., Rubtsova, M.Y., Ignatenko, O.V., Egorov, A.M., Kolesanova, E.F., & Archakov, A.I. (1997) Epitope mapping of horseradish peroxidase (isoenzyme C). Biochemistry-Engl. Tr., 62, 440-447. #9217

- 842. Andrews, M.J.I., & Tabor, A.B. (1997) Synthesis of an orthogonally-protected bifunctional amino acid for conformationally constrained peptides. Tetrahedron Lett., 38, 3063-3066. #8790
- 843. Angeletti, R.H., Bonewald, L.F., & Fields, G.B. (1997) Six-year study of peptide synthesis. Methods Enzymol., 289, 697-717.
- 844. Annis, I., Hargittai, B., & Barany, G. (1997) Disulfide bond formation in peptides. Methods Enzymol., 289, 198-221. #9994
- 845. Arano, Y., Akizawa, H., Uezono, T., Akaji, K., Ono, M., Funakoshi, S., Koizumi, M., Yokoyama, A., Kiso, Y., & Saji, H. (1997) Conventional and high-yield synthesis of DTPA-conjugated peptides: Application of a monoreactive dtpa to DTPA-D-Phe(1)-octreotide synthesis. Bioconjug. Chem., 8, 442-446. #9208
- 846. Arnold, D., Keilholz, W., Schild, H., Dumrese, T., Stevanovic, S., & Rammensee, H.G. (1997) Substrate specificity of cathepsins D and E determined by N-terminal and C-terminal sequencing of peptide pools. Eur. J. Biochem., 249, 171-179. #10103
- 847. Bagno, A., Bicciato, S., Dettin, M., & Di Bello, C. (1997) A novel algorithm for the coupling control in solid-phase peptide synthesis. J. Pept. Res., 50, 231-237. #9634
- 848. Barany, G., Albericio, F., Kates, S.A., & Kempe, M. (1997) Poly(ethylene glycol)-containing supports for solid-phase synthesis of peptides and combinatorial organic libraries. ACS Symp. Ser., 680, 239-264. #9612
- 849. Barany, G. & Kempe, M. (1997) The contex of solid-phase synthesis. In A.W. Czarnik & S.H. DeWitt (Eds.), A Practical Guide to Combinatorial Chemistry. (pp. 51-97). American Chemical Society, Washington, DC. #11696
- 850. Barany, G., Albericio, F., Kates, S.A., & Kempe, M. (1997) Polyethylene glycol-containing supports for solid phase synthesis of peptides and combinatorial organic libraries. Polym. Preprint., 38, 587-588. #8520
- 851. Barbar, E., Gross, C.M., Woodward, C., & Barany, G. (1997) Chemical synthesis and nuclear magnetic resonance characterization of partially folded proteins. Methods Enzymol., 289, 587-611. #10006
- 852. Bay, S., Loman, R., Osinaga, E., Nakada, H., Leclerc, C., & Cantacuzene, D. (1997) Preparation of a multiple antigen glycopeptide (MAG) carrying the Tn antigen: A possible approach to a synthetic carbohydrate vaccine. J. Pept. Res., 49, 620-625. #9438
- 853. Bechinger, B. (1997) Structure and dynamics of the M13 coat signal sequence in membranes by multidimensional high-resolution and solid-state NMR spectroscopy. Proteins, 27, 481-492. #9243
- 854. Beekman, N.J., Schaaper, W.M., Tesser, G.I., Dalsgaard, K., Kamstrup, S., Langeveld, J.P., Boshuizen, R.S., & Meloen, R.H. (1997) Synthetic peptide vaccines: Palmitoylation of peptide antigens by a thioester bond increases immunogenicity. Int. J. Peptide Prot. Res., 50, 357-364. #10114
- 855. Behrends, H.W., Beck-Sickinger, A.G., & Folkers, G. (1997) A new approach to secondary structure evaluation: Secondary structure prediction of porcine adenylate kinase and yeast guanylate kinase by CD spectroscopy og overlapping synthetic peptide segments. Biopolymers, 41, 213-231. #9959
- 856. Beltran, M., Maseda, M., Perez, Y., Robles, J., Pedroso, E., & Grandas, A. (1997) Stepwise solid-phase synthesis of serine-tyrosine- and homoserine-nucleopeptides. Nucleosides Nucleotides, 16, 1487-1488. #10115
- Benkovic, S., Miller, G.P., Zong, W. & Smiley, J. (1997) Combinatorial libraries of peptides, proteins, and antibodies using biological systems. In S.R. Wilson & A.W. Czarnik (Eds.), Combinatorial Chemistry. Synthesis and Application. (pp. 241-260). John Wiley & Sons, Inc., New York. #8361
- 858. Bennett, F.A., Barlow, D.J., Dodoo, A.N.O., Hider, R.C., Lansley, A.B., Lawrence, M.J., Marriott, C., & Bansal, S.S. (1997) L-(6,7-Dimethoxy-4-coumaryl)alanine: An intrinsic probe for the labelling of peptides. Tetrahedron Lett., 38, 7449-7452. #9574
- 859. Beranova-Giorgianni, S., & Desiderio, D.M. (1997) Fast atom bombardment mass spectrometry of synthetic peptides. Methods Enzymol., 289, 478-499. #10003
- Bernhardt, A., Drewello, M., & Schutkowski, M. (1997) The solid-phase synthesis of side-chain-phosphorylated peptide-4nitroanilides. J. Pept. Res., 50, 143-152. #9410

- 861. Bitan, G., Sukhotinsky, I., Mashriki, Y., Hanani, M., Selinger, Z., & Gilon, C. (1997) Synthesis and biological activity of novel backbone-bicyclic substance-P analogs containing lactam and disulfide bridges. J. Pept. Res., 49, 421-426. #9198
- 862. Blackburn, C., & Kates, S.A. (1997) Solid-phase synthesis of cyclic homodetic peptides. Methods Enzymol., 289, 175-198. #9857
- 863. Blackburn, C., Pingali, A., Kehoe, T., Herman, L.W., Wang, H.Q., & Kates, S.A. (1997) Libraries of angiotensin converting enzyme inhibitors: Solid-phase synthesis and affinity selection. Bioorg. Med. Chem. Lett., 7, 823-826. #9255
- 864. Blondelle, S.E., Forood, B., Houghten, R.A., & Perez-Paya, E. (1997) Polyaniline-based peptides as models for self-associated beta-pleated-sheet complexes. Biochemistry, 36, 8393-8400. #9477
- 865. Bodi, J., Suli-Vargha, H., Ludanyi, K., Vekey, K., & Orosz, G. (1997) New startegy for the synthesis of large peptides as applied to the C-terminal cysteine-rich 41 amino acid fragment of the mouse agouti protein. Tetrahedron Lett., 38, 3293-3296. #8799
- 866. Boumrah, D., Campbell, M.M., Fenner, S., & Kinsman, R.G. (1997) Spacer molecules in peptide sequences: Incorporation into analogues of atrial natriuretic factor. Tetrahedron, 53, 6977-6992. #9106
- 867. Braisted, A.C., Judice, J.K., & Wells, J.A. (1997) Synthesis of proteins by subtiligase. Methods Enzymol., 289, 298-313. #9997
- 868. Bray, A.M. (1997) Multiple peptide synthesis with commercially available Multipin kits. In I. Lefkovits (Ed.), Immunology Methods Manual. (pp. 809-816). Academic Press, San Diego. #9738
- 869. Breslav, M., Becker, J., & Naider, F. (1997) Dithioketal formation during synthesis of Bpa containing peptides. Tetrahedron Lett., 38, 2219-2222. #8774
- 870. Bruninghoff, A., Schlosser, S., Groger, G., Ortigao, J.F.R., & Seliger, H. (1997) Solid phase synthesis, cloning and expression of a synthetic gene encoding a collagen-like peptide. Nucleosides Nucleotides, 16, 875-882. #9423
- 871. Burdick, D.J., & Stults, J.T. (1997) Analysis of peptide synthesis products by electrospray ionization mass spectrometry. Methods Enzymol., 289, 499-519. #10004
- 872. Burgess, K., Lim, D., Bois-Choussy, M., & Zhu, J. (1997) Rapid and efficient solid phase syntheses of cyclic peptides with endocyclic biaryl ether bonds. Tetrahedron Lett., 38, 3345-3348. #8804
- 873. Burgess, K., Li, W., Lim, D.Y., & Moyesherman, D. (1997) Comparisons of the conformational biases imposed by trans-2,3-methanomethionine and alpha-methylmethionine. Biopolymers, 42, 439-453. #10132
- 874. Burgess, K., Li, W., Linthicum, D.S., Ni, Q., Pledger, D., Rothman, R.B., & Shitangkoon, A. (1997) Libraries of opiate and antiopiate peptidomimetics containing 2,3-methanoleucine. Bioorg. Med. Chem., 5, 1867-1871. #10131
- 875. Caciagli, V., Lombardi, P., Tuchalski, G., Hansicke, A., & Cardinali, F. (1997) Large-scale production of peptides using the solid-phase continuous flow method. Preparative synthesis of the novel tachykinin antagonist MEN 10628. J. Peptide Sci., 3, 224-230. #9936
- 876. Campbell, R.M., Felix, A.M., Stricker, P.R., Miller, R.W., Lee, Y., Lambros, T.J., Eisenbeis, H.G., Ahmad, M., & Heimer, E.P. (1997) Pegylates peptides. V.Carboxy-terminal PEGylated analogs of growth hormone-releasing factor (GRF) display enhanced duration of biological activity in vivo. J. Pept. Res., 49, 527-537. #9944
- 877. Canne, L.E., Winston, R.L., & Kent, S.B.H. (1997) Synthesis of a versatile purification handle for use with Boc chemistry solid phase peptide synthesis. Tetrahedron Lett., 38, 3361-3364. #8809
- 878. Cao, Y.J., Kojro, E., Gimpl, G., Jasionowski, M., Kasprzykowski, F., Lankiewicz, L., & Fahrenholz, F. (1997) Photoaffinity labeling analysis of the interaction of pituitary adenylate-cyclase-activating polypeptide (PACAP) with the PACAP type I receptor. Eur. J. Biochem., 244, 400-406. #9291
- 879. Carey, R.I., Bordas, L.W., Slaughter, R.A., Meadows, B.C., Wadsworth, J.L., Huang, H.H., Smith, J.J., & Furusjo, E. (1997) Preparation and properties of N-alpha-Bpoc-amino acid pentafluorophenyl esters. J. Pept. Res., 49, 570-581. #9437
- 880. Carrasco, M.R., Fitzgerald, M.C., Oda, Y., & Kent, S.B.H. (1997) Direct monitoring of organic reactions on polymeric supports. Tetrahedron Lett., 38, 6331-6334. #9318

- 881. Chandy, M.C., & Pillai, V.N.R. (1997) Gel phase synthesis of an octapeptide, H-Lys-Gly-Asp-Glu-Glu-Ser-Leu-Ala-OH on a flexible triethyleneglycol dimethacrylate-crosslinked polystyrene support. Ind. J. Chem. B-Org. Chem. Med. Chem., 36, 303-307. #9539
- 882. Chen, J.J., & Spatola, A.F. (1997) Solid phase synthesis of peptide hydroxamic acids. Tetrahedron Lett., 38, 1511-1514. #8374
- 883. Chen, L., Zoulikova, I., Slaninova, J., & Barany, G. (1997) Synthesis and pharmacology of novel analogues of oxytocin and deaminooxytocin: Directed methods for the construction of disulfide and trisulfide bridges in peptides. J. Med. Chem., 40, 864-876. #9286
- 884. Chen, S.T., Tseng, P.H., Yu, H.M., Wu, C.Y., Hsiao, K.F., Wu, S.H., & Wang, K.T. (1997) The studies of microwave effects on the chemical reactions. J. Chin. Chem. Soc., 44, 169-182. #9531
- 885. Chillemi, F., Galatulas, I., Fraccari, A., Bossa, R., & Francescato, P. (1997) Enhancement of cytotoxic activity by synthesis of peptide multimeric forms. Anticancer Res., 17, 3609-3611. #9924
- 886. Chirinos-Rojas, C.L., Steward, M.W., & Partidos, C.D. (1997) Of a solid-phase random peptide library to identify inhibitors of TNF-alpha mediated cytotoxicity in vitro. Cytokine, 9, 226-232. #9258
- 887. Cho, Y.R., Entress, R.M.H., & Williams, D.H. (1997) Synthesis of cell-wall analogues of vancomycin-resistant enterococci using solid phase peptide synthesis. Tetrahedron Lett., 38, 5229-5232. #9036
- 888. Choi, H., Murray, T.F., DeLander, G.E., Schmidt, W.K., & Aldrich, J.V. (1997) Synthesis and opioid activity of [D-Pro(10)] dynorphin A-(1-11) analogues with N-terminal alkyl substitution. J. Med. Chem., 40, 2733-2739. #9401
- 889. Cilli, E.M., Marchetto, R., Schreier, S., & Nakaie, C.R. (1997) Use of spin label EPR spectra to monitor peptide chain aggregation inside resin beads. Tetrahedron Lett., 38, 517-520. #12510
- 890. Cody, W.L., He, J.X., Reily, M.D., Haleen, S.J., Walker, D.M., Reyner, E.L., Stewart, B.H., & Doherty, A.M. (1997) Design of a potent combined pseudopeptide endothelin-A/endothelin-B receptor antagonist, AC-DBHG(16)-Leu-Asp-Ile-[NME]Ile-Trp(21) (PD 156252): Examination of its pharmacokinetic and spectral properties. J. Med. Chem., 40, 2228-2240. #9483
- 891. Corey, D.R. (1997) Peptide nucleic acids: Expanding the scope of nucleic acid recognition. Trends Biotech., 15, 224-229. #9192
- 892. Cormack, P.A., Moore, B.D., & Sherrington, D.C. (1997) Monodisperse liquid crystalline peptides. J. Material. Chem., 7, 1977-1983. #10147
- Cross, T.A. (1997) Solid-state nuclear magnetic resonance characterization of gramicidin channel structure. Methods Enzymol., 289, 672-696. #9930
- 894. Dahmani, B., Krebs, D., el Antri, S., Troalen, F., & Fermandjian, S. (1997) Synthesis and FTIR conformational studies of peptides from the basic region of c-Jun: A critical analysis on the basis of CD and NMR data. J. Biomol. Struct. Dyn., 14, 429-439. #9287
- 895. Davies, M., & Bradley, M. (1997) C-terminally modified peptides and peptide libraries: Another end to peptide synthesis. Angew. Chem. Int. Ed., 36, 1097-1099. #12487
- 896. Dawson, P.E., Fitzgerald, M.C., Muir, T.W., & Kent, S.B. (1997) Methods for the chemical synthesis and readout of self-encoded arrays of polypeptide analogues. J. Amer. Chem. Soc., 119, 7917-7927. #10155
- 897. Del Nery, E., Juliano, M.A., Meldal, M., Svendsen, I., Scharfstein, J., Walmsley, A., & Juliano, L. (1997) Characterization of the substrate specificity of the major cysteine protease (cruzipain) from Trypanosoma cruzi using a portion-mixing combinatorial library and fluorogenic peptides. Biochem. J., 323, 427-433. #9248
- 898. Dettin, M., Pegoraro, S., Rovero, P., Bicciato, S., Bagno, A., & Di Bello, C. (1997) SPPS of difficult sequences: A comparison of chemical conditions, synthetic strategies and on-line monitoring. J. Pept. Res., 49, 103-111. #9225
- 899. Dettin, M., Roncon, R., Simonetti, M., Tormene, S., Falcigno, L., Paolillo, L., & Di Bello, C. (1997) Synthesis, characterization and conformational analysis of gp120-derived synthetic peptides that specifically enhance HIV-1 infectivity. J. Peptide Sci., 3, 15-30. #8579

- 900. Dhalluin, C., Boutillon, C., Tartar, A., & Lippens, G. (1997) Magic angle spinning nuclear magnetic resonance in solid-phase peptide synthesis. J. Amer. Chem. Soc., 119, 10494-10500. #9602
- 901. Dong, C.Z., Derocquigny, H., Remy, E., Mellac, S., Fournie-Zaluski, M.C., & Roques, B.P. (1997) Synthesis and biological activities of fluorescent acridine-containing HIV-1 nucleocapsid proteins for investigation of nucleic acid-ncp7 interactions. Int. J. Peptide Prot. Res., 50, 269-278. #10161
- 902. Dorner, B., Ostresh, J.M., Blondelle, S.E., Dooley, C.T. & Houghten, R.A. (1997) Peptidomimetic synthetic combinatorial libraries. In Anonymous, Advances in Amino Acid Mimetics and Peptidomimetics. (pp. 109-125). JAI Press, Inc., #11629
- 903. Drouot, C., Enjalbal, C., Fulcrand, P., Martinez, J., & Aubagnac, J.L. (1997) Tof-SIMS analysis of polymer bound Fmoc-protected peptides. Tetrahedron Lett., 38, 2455-2458. #8779
- 904. Eggleston, I. M., Peggion, C., Svendsen, J. S., and Mutter, M. Practical synthesis and application of a beta-turn mimetic for SPPS. 1997. Poster presented at the Fifth International Symposium on Solid Phase Synthesis & Combinatorial Chemical Libraries, London, England, UK on 2nd-6th September 1997. #9699
- 905. Egner, B.J., & Bradley, M. (1997) Monitoring the solid phase synthesis of analogues of lysobactin and the katanosins using in situ Maldi-Tof MS. Tetrahedron, 53, 14021-14030. #10166
- 906. Eichhorn, U., Jakubke, H.D., Schaaf, R., & Beck-Piotraschke, K. (1997) Solid-phase acyldonor as a substrate pool in kinetically controlled protease-catalysed peptide synthesis. J. Peptide Sci., 3, 261-266. #9983
- 907. Eichler, J., & Houghten, R.A. (1997) Synthesis of cyclic disulfide peptides: Comparison of oxidation methods. Protein Peptide Lett., 4, 157-164. #11603
- 908. Elgendy, S., Patel, G., Green, D., Goodwin, C.A., Scully, M.F., Husman, W., Skordalakes, E., Kakkar, V.V., & Deadman, J.J. (1997) Design of a novel class of bifunctional thrombin inhibitors, synthesised by the first application of peptide boronates in solid phase chemistry. Tetrahedron Lett., 38, 3305-3308. #8802
- 909. Elofsson, M., Salvador, L.A., & Kihlberg, J. (1997) Preparation of Tn and sialyl Tn building blocks used in Fmoc solid-phase synthesis of glycopeptide fragments from HIV gp120. Tetrahedron, 53, 369-390. #12511
- 910. Epton, R. (1997) Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries. Mayflower Scientific Limited, Birmingham. ISBN 0952701111 #8773
- 911. Evans, D.M., & Herman, L.W. (1997) Identification of peptide ligands specific for the sugar-binding site of concanavalin A by screening a synthetic peptide combinatorial library. J. Biomol. Screen., 2, 225-233. #10168
- 912. Fehrentz, J.A., Paris, M., Heitz, A., Velek, J., Winternitz, F., & Martinez, J. (1997) Solid phase synthesis of C-terminal peptide aldehydes. J. Org. Chem., 62, 6792-6796. #10172
- 913. Feng, Y.B., Melacini, G., & Goodman, M. (1997) Collagen-based structures containing the peptoid residue N-isobutylglycine (Nleu): Synthesis and biophysical studies of Gly-Nleu-Pro sequences by circular dichroism and optical rotation. Biochemistry, 36, 8716-8724. #9449
- 914. Fields, G.B., & Colowick, S.P. (1997) Solid-Phase Peptide Synthesis. Academic Press, San Diego. ISBN 0121821900 #9830
- 915. Flouzat, C., Marguerite, F., Croizet, F., Percebois, M., Monteil, A., & Combourieu, M. (1997) Solid-phase synthesis of "head-to-side chain" cyclic tripeptides using allyl deprotection. Tetrahedron Lett., 38, 1191-1194. #8390
- 916. Frank, R. (1997) Multiple peptide synthesis with SPOT technique. In I. Lefkovits (Ed.), Immunology Methods Manual. (pp. 783-795). Academic Press, San Diego. #9739
- 917. Frank, R. (1997) Principles of simultaneous synthesis of multiple peptides. In I. Lefkovits (Ed.), Immunology Methods Manual. (pp. 777-781). Academic Press, San Diego. #9740
- 918. Franzyk, H., Christensen, M.K., Jorgensen, R.M., Meldal, M., Cordes, H., Mouritsen, S., & Bock, K. (1997) Constrained glycopeptide ligands for MPRs. Limitations of unprotected phosphorylated building blocks. Bioorg. Med. Chem., 5, 21-40. #9947

- 919. Fujii, M., Yoshida, K., & Hidaka, J. (1997) Nucleic acid analog peptide (NAAP). 2. Syntheses and properties of novel DNA analog peptides containing nucleobase linked beta-aminoalanine. Bioorg. Med. Chem. Lett., 7, 637-640. #10023
- 920. Fujii, M., Yamamoto, K., Hidaka, J., & Ohtsu, T. (1997) Nucleic acid analog peptide (NAAP). Solid phase synthesis of a DNA analog peptide. Tetrahedron Lett., 38, 417-420. #8246
- 921. Furet, P., Gay, B., Garcia-Echeverria, C., Rahuel, J., Fretz, H., Schoepfer, J., & Caravatti, G. (1997) Discovery of 3-aminobenzyloxycarbonyl as an N-terminal group conferring high affinity to the minimal phosphopeptide sequence recognized by the Grb2-SH2 domain. J. Med. Chem., 40, 3551-3556. #10178
- 922. Futaki, S., Sogawa, K., Maruyama, J., Asahara, T., Niwa, M., & Hojo, H. (1997) Preparation of peptide thioesters using Fmocsolid-phase peptide synthesis and its application to the construction of a template-assembled synthetic protein (TASP). Tetrahedron Lett., 38, 6237-6240. #9316
- 923. Galeotti, N., Plagnes, E., & Jouin, P. (1997) Synthesis of peptidyl aldehydes from thiazolidines. Tetrahedron Lett., 38, 2459-2462. #8780
- 924. Gariani, T., & Leatherbarrow, R.J. (1997) Stability of protease inhibitors based on the Bowman-Birk reactive site loop to hydrolysis by proteases. J. Pept. Res., 49, 467-475. #9435
- 925. Gasparri, A., Sidoli, A., Sanchez, L.P., Longhi, R., Siccardi, A.G., Marchisio, P.C., & Corti, A. (1997) Chromogranin a fragments modulate cell adhesion: Identification and characterization of a pro-adhesive domain. Journal of Biological Chemistry, 272, 20835-20843. #9402
- 926. Giblin, M.F., Jurisson, S.S., & Quinn, T.P. (1997) Synthesis and characterization of rhenium-complexed alpha- melanotropin analogs. Bioconjug. Chem., 8, 347-353. #9206
- 927. Gobbo, M., Biondi, L., Cavaggion, F., Filira, F., Piek, T., Mantel, P., & Rocchi, R. (1997) Synthesis and biological activities of head-to-tail cyclic bradykinin analogues of varying ring size. Int. J. Peptide Prot. Res., 50, 336-341. #10190
- 928. Goodnow, R.A.Jr., Richou, A.R., & Tam, S. (1997) Synthesis of thymine, cytosine, adenine, and guanine containing N-Fmoc protected amino acids: Building blocks for construction of novel oligonucleotide backbone analogs. Tetrahedron Lett., 38, 3195-3198. #8796
- 929. Goodnow, R.A.Jr., Tam, S., Pruess, D.L., & McComas, W.W. (1997) Oligomer synthesis of DNA / RNA recognition properties of a novel oligonucleotide backbone analog: Glucopyranosyl nucleic amide (GNA). Tetrahedron Lett., 38, 3199-3202. #8797
- 930. Grant, G.A., Crankshaw, M.W., & Gorka, J. (1997) Edman sequencing as a tool for characterization of synthetic peptides. Methods Enzymol., 289, 395-419. #9999
- 931. Griffith, D.L., O'Donnell, M.J., Pottorf, R.S., Scott, W.L., & Porco, J.A.Jr. (1997) Tandem UPS: Sequential mono- and dialkylation of resin-bound glycine via automated synthesis. Tetrahedron Lett., 38, 8821-8824. #9697
- 932. Grim, M.D. (1997) New technology for the preparation of peptides and peptidomimetics. Pharm. Manufact. Int., 49-49. #11184
- 933. Guibourdenche, C., Seebach, D., & Natt, F. (1997) Use of the Wolff rearrangement of diazo ketones from amino acids as a synthetic method for the formation of oligonucleo-peptides a novel approach to chimeric biomolecules. Helv. Chim. Acta, 80, 1-13. #8386
- 934. Guichard, G., & Seebach, D. (1997) Solid-phase synthesis of beta-oligopeptides. Chimia, 51, 315-318. #9538
- 935. Guo, Z.W., Nakahara, Y., & Ogawa, T. (1997) Solid-phase synthesis of CD52 glycopeptide and an efficient route to Asn-core pentasaccharide conjugate. Bioorg. Med. Chem., 5, 1917-1924. #10199
- 936. Guo, Z.W., Nakahara, Y., & Ogawa, T. (1997) Solid-phase synthesis of a glycosylated peptide fragment of the il-8 receptor containing two vicinal oligosaccharide chains. Carbohyd. Res., 303, 373-377. #10200
- 937. Guo, Z.W., Nakahara, Y., & Ogawa, T. (1997) Solid-phase synthesis of the CD52 glycopeptide carrying an N-linked core pentasaccharide structure. Angew. Chem. Int. Ed., 36, 1464-1466. #12512

- 938. Guy, C.A., & Fields, G.B. (1997) Trifluoroacetic acid cleavage and deprotection of resin-bound peptides following synthesis by Fmoc chemistry. Methods Enzymol., 289, 67-83. #9992
- 939. Hackeng, T.M., Mounier, C.M., Bon, C., Dawson, P.E., Griffin, J.H., & Kent, S.B.H. (1997) Total chemical synthesis of enzymatically active human type II secretory phospholipase A(2). Proc. Natl. Acad. Sci. USA, 94, 7845-7850. #9451
- 940. Hamy, F., Felder, E.R., Heizmann, G., Lazdins, J., Aboul-ela, F., Varani, G., Karn, J., & Klimkait, T. (1997) An inhibitor of the Tat/TAR RNA interaction that effectively suppresses HIV-1 replication. Proc. Natl. Acad. Sci. USA, 94, 3548-3553. #8591
- 941. Han, Y.X., Albericio, F., & Barany, G. (1997) Occurrence and minimization of cysteine racemization during stepwise solid-phase peptide synthesis. J. Org. Chem., 62, 4307-4312. #9506
- 942. Han, Y.X., & Barany, G. (1997) Novel s-xanthenyl protecting groups for cysteine and their applications for the n-alpha-9-fluorenylmethyloxycarbonyl (fmoc) strategy of peptide synthesis. J. Org. Chem., 62, 3841-3848. #9074
- 943. Harada, K., Martin, S.S., Tan, R.Y., & Frankel, A.D. (1997) Molding a peptide into an RNA site by in vivo peptide evolution. Proc. Natl. Acad. Sci. USA, 94, 11887-11892. #10206
- 944. Harthfritschy, E., & Cantacuzene, D. (1997) Esterification of 9-fluorenylmethoxycarbonyl-glycosylated serine and cysteine derivatives with an hydroxymethyl resin. Int. J. Peptide Prot. Res., 50, 415-420. #10208
- 945. Henkel, B., Zhang, L.S., & Bayer, E. (1997) Investigations on solid-phase peptide synthesis in N-to-C direction (inverse synthesis). Liebig. Ann., 2161-2168. #10211
- 946. Henkel, B., Zeng, W., & Bayer, E. (1997) 9-Hydroxy-9-(4-carboxyphenyl)xanthene: A new linker for the synthesis of peptide amides. Tetrahedron Lett., 38, 3511-3512. #8849
- 947. Hernandez, J.F., Bersch, B., Petillot, Y., Gagnon, J., & Arlaud, G.J. (1997) Chemical synthesis and characterization of the epidermal growth factor-like module of human complement protease C1r. J. Pept. Res., 49, 221-231. #9213
- 948. Hoffmann, R., Hoffmann, T., Tholey, A., Schulte, A.C., & Kalbitzer, H.R. (1997) Synthesis and NMR spectroscopy of peptides containing either phosphorylated or phosphorylated cis- or trans-4-hydroxy-l-proline. J. Pept. Res., 49, 163-173. #9227
- 949. Hojo, H., Akamatsu, Y., Yamauchi, K., Kinoshita, M., Miki, S., & Nakamura, Y. (1997) Synthesis and structural characterization of triple-helical peptides which mimic the ligand binding site of the human macrophage scavenger receptor. Tetrahedron, 53, 14263-14274. #10215
- 950. Holm, A., Ostergaard, S., Hodges, R.S., Husband, D. & Buus, S. (1997) Multiple peptide synthesis on resin. In I. Lefkovits (Ed.), Immunology Methods Manual. (pp. 847-851). Academic Press, San Diego. #9737
- 951. Holmes, D.L., Smith, E.M., & Nowick, J.S. (1997) Solid-phase synthesis of artificial beta-sheets. J. Amer. Chem. Soc., 119, 7665-7669. #10216
- 952. Hruby, V.J., Ahn, J.M., & Liao, S. (1997) Synthesis of oligopeptide and peptidomimetic libraries. Curr. Opin. Chem. Biol., 1, 114-119. #8770
- 953. Humphrey, J.M., & Chamberlin, A.R. (1997) Chemical synthesis of natural product peptides: Coupling methods for the incorporation of noncoded amino acids into peptides [review]. Chem. Rev., 97, 2243-2266. #10225
- 954. Husbyn, M., Orning, L., Sakariassen, K.S., & Fischer, P.M. (1997) Peptides corresponding to the second epidermal growth factor-like domain of human blood coagulation factor. VII. Synthesis, folding and biological activity. Int. J. Peptide Prot. Res., 50, 475-482. #10226
- 955. Ingenhoven, N., & Beck-Sickinger, A.G. (1997) Fluorescent labelled analogues of neuropeptide Y for the characterization of cells expressing NPY receptor subtypes. J. Recept. Signal Transduct. Res., 17, 407-418. #9953
- 956. Izdebski, J., Orlowska, A., Pachulska, M., & Witkowska, E. (1997) Peptide synthesis with the use of N,N'-dicyclopentylcarbodiimide. Polish J. Chem., 71, 903-907. #9542
- 957. Jaikaran, D.C.J., Woolley, G.A., Sansom, M.S., Wenschuh, H., & Biggin, P.C. (1997) Structure-function relationships in helix-

- bundle channels probed via total chemical synthesis of alamethicin dimers: Effects of a Gln7 to Asn7 mutation. Biochemistry, 36, 13873-13881. #9937
- 958. Janevik-Ivanovska, E., Gruaz-Guyon, A., Barbet, J., Rostene, W., Tartar, A., Milhaud, G., Cohen, M., Hillairet de Boisferon, M., & Gautherot, E. (1997) Bivalent hapten-bearing peptides designed for iodine-131 pretargeted radioimmunotherapy. Bioconjug. Chem., 8, 526-533. #9434
- 959. Jelinek, R., Valente, A.P., Valentine, K.G., & Opella, S.J. (1997) Two-dimensional NMR spectroscopy of peptides on beads. J. Magn. Resonance, 125, 185-187. #9150
- 960. Johansson, E., Lange, S., & Lonnroth, I. (1997) Identification of an active site in the antisecretory factor protein. BBA-Mol. Basis Dis., 1362, 177-182. #11170
- 961. Jones, D.M., Sueiras-Diaz, J., Szelke, M., Leckie, B.J., Beattie, S.R., Morton, J., Neidle, S., & Kuroda, R. (1997) New renin inhibitors containing novel analogues of statine. J. Pept. Res., 50, 109-121. #9409
- 962. Kaiser, T., & Voelter, W. (1997) Solid phase peptide synthetic approaches to mouse macrophage migration inhibitory factor (mMIF): Comparison of different strategies producing the C-terminal difficult sequence peptide mMIF(93-115) [german]. Zeitschrift fur Naturforschung Section B-A Journal of Chemical Sciences, 52, 281-295. #9170
- 963. Kaiser, T., & Voelter, W. (1997) Solid phase peptide synthetic approaches to mouse macrophage migration inhibitory factor (mMIF): Synthesis of peptide fragments for a convergent solid phase peptide synthetic strategy [german]. Journal fur Praktische Chemie-Chemiker-Zeitung, 339, 371-380. #9072
- 964. Karginov, V.A., Mamaev, S.V., An, H.Y., Vancleve, M.D., Hecht, S.M., Komatsoulis, G.A., & Abelson, J.N. (1997) Probing the role of an active site aspartic acid in dihydrofolate reductase. J. Amer. Chem. Soc., 119, 8166-8176. #10235
- 965. Karlstrom, A. H. and Und n, A. E. Development of a new class of protecting groups for use in solid phase peptide synthesis. 1997. Poster presented at the 15th APS, June 14-19,1997, Nashville, Tennessee. #11796
- 966. Kellam, B., Chan, W.C., Chhabra, S.R., & Bycroft, B.W. (1997) Transient affinity tags based on the Dde protection/deprotection strategy: Synthesis and application of 2-biotinyl- and 2-hexanoyldimedone. Tetrahedron Lett., 38, 5391-5394. #9054
- 967. Kennedy, K.J., Lundquist, J.T.IV., Simandan, T.L., Beeson, C.C., & Dix, T.A. (1997) Asymmetric synthesis of non-natural homologues of lysine. Bioorg. Med. Chem. Lett., 7, 1937-1940. #9426
- 968. Kihlberg, J., Elofsson, M., & Salvador, L.A. (1997) Direct synthesis of glycosylated amino acids from carbohydrate peracetates and Fmoc amino acids: Solid-phase synthesis of biomedicinally interesting glycopeptides. Methods Enzymol., 289, 221-245. #9934
- 969. Kihlberg, J., & Elofsson, M. (1997) Solid-phase synthesis of glycopeptides: Immunological studies with T cell stimulating glycopeptides. Curr. Med. Chem., 4, 85-116. #12516
- 970. Kimura, T., Fukui, T., Tanaka, S., Akaji, K., & Kiso, Y. (1997) A reductive acidolysis final deprotection strategy in solid phase peptide synthesis based on safety-catch protection. Chem. Pharm. Bull. Tokyo, 45, 18-26. #8234
- 971. King, A.D., Huh, N.W., Pedersen, L.G., & Hiskey, R.G. (1997) Studies on the peptide corresponding to residues 34-47 of bovine factor X. J. Pept. Res., 50, 34-38. #9416
- 972. Kitagawa, K., Aida, C., Fujiwara, H., Yagami, T., & Futaki, S. (1997) Efficient solid-phase synthesis of sulfated tyrosine-containing peptides using 2-chlorotrityl resin: Facile synthesis of gastrin/cholecystokinin peptides. Tetrahedron Lett., 38, 599-602. #8237
- 973. Klein, S.I., Czekaj, M., Molino, B.F., & Chu, V. (1997) Constrained beta-alanine based GpIIb/IIa antagonists. Bioorg. Med. Chem. Lett., 7, 1773-1778. #9473
- 974. Klodt, J., Adermann, K., Forssmann, W.G., Rosch, P., Martin, S., Marx, U.C., & Kuhn, M. (1997) Synthesis, biological activity and isomerism of guanylate cyclase C-activating peptides guanylin and uroguanylin. J. Pept. Res., 50, 222-230. #9925
- 975. Ko, S.Y., & Wenger, R.M. (1997) 53. Solid-phase total synthesis of cyclosporine analogues. Helv. Chim. Acta, 80, 695-705. #9087

- 976. Koch, T., Borre, M.B., Naesby, M., Batz, H.G., & Orum, H. (1997) PNA: Synthesis and diagnostic application. Nucleosides Nucleotides, 16, 1771-1774. #9741
- 977. Koppitz, M., Huenges, M., Gratias, R., Kessler, H., Goodman, S.L., & Jonczyk, A. (1997) Synthesis of unnatural lipophilic N-(9H-fluoren-9-yl-methoxy)carbonyl-substituted alpha-amino acids and their incorporation into cyclic RGD-peptides: A structure activity study. Helv. Chim. Acta, 80, 1280-1300. #9461
- 978. Kowalczyk, C., & O'Shea, M. (1997) Solid-phase synthesis of neuropeptides by Fmoc strategies. Methods Mol. Biol., 73, 41-48. #9952
- 979. Lal, D., Mahale, S.D., Nandedkar, T.D., & Iyer, K.S. (1997) Identification of bioneutralization epitopes of human follicle stimulating hormone in the regions 31-52 and 66-75 of its beta- subunit. J. Reprod. Immunol., 33, 1-14. #9202
- 980. Lebl, M., & Krchnak, V. (1997) Synthetic peptide libraries. Methods Enzymol., 289, 336-392. #9998
- 981. Lelievre, D., Hsu, S.C., Daubos, P., Favard, C., Vigny, P., Trudelle, Y., Steward, M.W., & Delmas, A. (1997) Structural properties of chimeric peptides containing a T-cell epitope linked to a fusion peptide and their importance for in vivo induction of cytotoxic T-cell responses. Eur. J. Biochem., 249, 895-904. #10264
- 982. Leonard, D.M., Shuler, K.R., Poulter, C.J., Eaton, S.R., Sawyer, T.K., Hodges, J.C., Su, T.Z., Scholten, J.D., Gowan, R.C., Sebolt-Leopold, J.S., & Doherty, A.M. (1997) Structure-activity relationships of cysteine-lacking pentapeptide derivatives that inhibit ras farnesyltransferase. J. Med. Chem., 40, 192-200. #8243
- 983. Liakopouloukyriakides, M., Pachatouridis, C., Ekateriniadou, L., & Papageorgiou, V.P. (1997) A new synthesis of the tripeptide Gly-His-Lys with antimicrobial activity. Amino Acids, 13, 155-161. #10266
- 984. Liebe, B., & Kunz, H. (1997) Solid-phase synthesis of a tumor-associated sialyl-TN antigen glycopeptide with a partial sequence of the "tandem repeat" of the MUC-1 mucin. Angew. Chem. Int. Ed., 36, 618-621. #12518
- 985. Liu, C.F., & Tam, J.P. (1997) Synthesis of a symmetric branched peptide. Assembly of a cyclic peptide on a small tetraacetate template. Chem. Commun., 1619-1620. #11687
- 986. Liu, W.Q., Roques, B.P., & Garbay, C. (1997) Synthesis of l-2,3,5,6-tetrafluoro-4-(phosphonomethyl) phenylalanine, a novel non-hydrolyzable phosphotyrosine mimetic and l-4-(phosphonodifluoromethyl)phenylalanine. Tetrahedron Lett., 38, 1389-1392. #8379
- 987. Lloyd-Williams, P., Albericio, F., & Giralt, E. (1997) Chemical Approaches to the Synthesis of Peptides and Proteins. CRC Press, Boca Raton. ISBN 0849391423 #9305
- 988. Lohse, J., Nielsen, P.E., Harrit, N., & Dahl, O. (1997) Fluorescein-conjugated lysine monomers for solid phase synthesis of fluorescent peptides and PNA oligomers. Bioconjug. Chem., 8, 503-509. #9433
- 989. Lokey, R.S., Kwok, Y., Guelev, V., Pursell, C.J., Hurley, L.H., & Iverson, B.L. (1997) A new class of polyintercalating molecules. J. Amer. Chem. Soc., 119, 7202-7210. #9419
- 990. Lou, Q., Leftwich, M.E., Mckay, R.T., Salmon, S.E., Rychetsky, L., & Lam, K.S. (1997) Potent pseudosubstrate-based peptide inhibitors for p60(c-src) protein tyrosine kinase. Cancer Res., 57, 1877-1881. #9070
- 991. Lozanov, V., Guarnaccia, C., Patthy, A., Foti, S., & Pongor, S. (1997) Synthesis and cysteine/cysteine-catalyzed oxidative folding of the amaranth alpha-amylase inhibitor. J. Pept. Res., 50, 65-72. #9417
- 992. Lu, H.S.M., Volk, M., Kholodenko, Y., Gooding, E., Hochstrasser, R.M., & DeGrado, W.F. (1997) Aminothiotyrosine disulfide, an optical trigger for initiation of protein folding. J. Amer. Chem. Soc., 119, 7173-7180. #9418
- 993. Lucietto, P., Mascagni, P., Giuliani, P., Ball, H.L., & Fossati, G. (1997) Mycobacterium tuberculosis chaperonin 10 and N-truncated fragments. Their synthesis and purification by the isoelectric focusing technique carried out in solution. J. Pept. Res., 49, 308-323. #9950
- 994. Lynas, J.F., & Walker, B. (1997) Peptide argininol "inverse substrates" of anisic acid: Novel inhibitors of the trypsin-like serine proteinases. Bioorg. Med. Chem. Lett., 7, 1133-1138. #10029

- 995. Ma, J.B., Xie, Z.D., Wang, Y.J., Wang, Y.N., Fan, Y.G., & He, B.L. (1997) Polymeric adsorbents with peptide pendants as artificial receptors for beta-lactam antibiotics by mimicking the binding sites of beta-lactamases. Sci. China Ser. B, 40, 405-411. #9522
- 996. Maeda, M., Izuno, Y., Kawasaki, K., Kaneda, Y., Mu, Y., Tsutsumi, Y., Nakagawa, S., & Mayumi, T. (1997) Amino acids and peptides. XXX.Preparation of Arg-Gly-Asp (RGD) hybrids with poly(ethylene glycol)analogs and their antimetastatic effect. Chem. Pharm. Bull. Tokyo, 45, 1788-1792. #10274
- Maeda, M., Izuno, Y., Kawasaki, K., Kaneda, Y., Mu, Y., Tsutsumi, Y., Lem, K.W., & Mayumi, T. (1997) Amino acids and peptides. XXXII.A bifunctional poly(ethylene glycol)hybrid of fibronectin-related peptides. Biochem. Biophys. Res. Commun., 241, 595-598. #10273
- 998. Maletinska, L., Neugebauer, W., Perodin, J., Lefebvre, M., & Escher, E. (1997) Angiotensin analogues palmitoylated in positions land 4. J. Med. Chem., 40, 3271-3279. #9940
- 999. Manning, M., Cheng, L.L., Stoev, S., Klis, W.A., Nawrocka, E., Olma, A., Sawyer, W.H., Wo, N.C., & Chan, W.Y. (1997)
  Position three in vasopressin antagonist tolerates conformationally restricted and aromatic amino acid substitutions: A striking contrast with vasopressin agonists. J. Peptide Sci., 3, 31-46. #8580
- 1000. Mant, C.T., Kondejewski, L.H., Cachia, P.J., Monera, O.D., & Hodges, R.S. (1997) Analysis of synthetic peptides by high-performance liquid chromatography. Methods Enzymol., 289, 426-469. #10001
- 1001. Marsh, I.R., & Bradley, M. (1997) Solid phase synthesis of polyamine conjugates for the study of trypanothione reductase. Tetrahedron, 53, 17317-17334. #10278
- 1002. Marsh, I.R., Bradley, M., & Teague, S.J. (1997) Solid-phase total synthesis of oscillamide Y and analogues. J. Org. Chem., 62, 6199-6203. #10279
- 1003. Marti, R.E., Bleicher, K.H., & Bair, K.W. (1997) Solid phase synthesis of beta-peptides via Arndt-Eistert homologation of Fmoc-protected amino acid diazoketones. Tetrahedron Lett., 38, 6145-6148. #9313
- 1004. Mayo, K.H. (1997) Solution nuclear magnetic resonance characterization of peptide folding. Methods Enzymol., 289, 646-672. #10009
- 1005. Meisenbach, M., Echner, H., & Voelter, W. (1997) New methoxy-substituted 9-phenylxanthen-9-ylamine linkers for the solid phase synthesis of protected peptide amides. Chem. Commun., 849-850. #9124
- 1006. Meisenbach, M., & Voelter, W. (1997) New methoxy-substituted tritylamine linkers for the solid phase synthesis of protected peptide amides. Chem. Lett., 1265-1266. #10289
- 1007. Meldal, M. (1997) Properties of solid supports. Methods Enzymol., 289, 83-104. #9929
- 1008. Meldal, M., Juliano, M.A., & Jansson, A.M. (1997) Azido acids in a novel method of solid-phase peptide synthesis. Tetrahedron Lett., 38, 2531-2534. #8781
- 1009. Merchan, F.L., Merino, P., & Tejero, T. (1997) A new entry to glycosylamines. Glycoconjugate J., 14, 497-499. #9480
- 1010. Merrifield, B. (1997) Concept and early development of solid-phase peptide synthesis. Methods Enzymol., 289, 3-13. #9932
- 1011. Mihara, H., Hayashida, J., Hasegawa, H., Ogawa, H.I., Fujimoto, T., & Nishino, N. (1997) A pair of pyrene groups as a conformational probe for antiparallel beta-sheet structure, formed in cyclic peptides. J. Chem. Soc. Perkin Trans. 2, 517-522. #9174
- 1012. Miller, S.C., & Scanlan, T.S. (1997) Site-selective N-methylation of peptides on solid support. J. Amer. Chem. Soc., 119, 2301-2302. #9552
- 1013. Milton, S. C. F., de L.Milton, R. C., Kates, S. A., and Glabe, C. Synthesis of Alzheimer's (1-42)A beta-amyloid peptide with preformed Fmoc-aminoacyl fluorides. In: D.R. Marshak. (Eds.) 1997; Academic Press, San Diego, CA p.865 Tech. Protein Chem. VIII (Symp. Protein Soc.) 10th. #9669

- 1014. Mizukoshi, T., Ariyoshi, Y., Suzuki, E., & Kohmura, M. (1997) Structure and dynamic studies by NMR of the potent sweet protein monellin and a non-sweet analog. Evidence on the importance of residue AspB7 for sweet taste. FEBS Lett., 413, 409-416. #9943
- 1015. Mokotoff, M., Swanson, D.P., Jonnalagadda, S.S., Epperly, M.W., & Brown, M.L. (1997) Evaluation of laminin peptide fragments labeled with indium-111 for the potential imaging of malignant tumors. J. Pept. Res., 49, 510-516. #9436
- 1016. Moore, W.T. (1997) Laser desorption mass spectrometry. Methods Enzymol., 289, 520-542. #10005
- 1017. Moos, W.H. (1997) Introduction: Combinatorial chemistry approaches the next millenium. In A.W. Czarnik & S.H. DeWitt (Eds.), A Practical Guide to Combinatorial Chemistry. (pp. 1-16). American Chemical Society, Washington, DC. #11694
- 1018. Muir, T.W., Dawson, P.E., & Kent, S.B.H. (1997) Protein synthesis by chemical ligation of unprotected peptides in aqueous solution. Methods Enzymol., 289, 266-298. #9996
- 1019. Nakahara, Y., Ito, Y., & Ogawa, T. (1997) Total synthesis of B-chain of human alpha2HS glycoprotein. Tetrahedron Lett., 38, 7211-7214. #9562
- 1020. Nastri, F., Lombardi, A., Morelli, G., Maglio, O., Dauria, G., Pedone, C., & Pavone, V. (1997) Hemoprotein models based on a covalent helix-heme-helix sandwich. 1.Design, synthesis, and characterization. Chem. Eur. J., 3, 340-349. #9136
- 1021. Nicolas, E., Pujades, M., Bacardit, J., Giralt, E., & Albericio, F. (1997) A new approach to Hmb-backbone protection of peptides: Synthesis and reactivity of N(alpha)-Fmoc-N(alpha)-(Hmb)amino acids. Tetrahedron Lett., 38, 2317-2320. #8777
- 1022. Nicolas, E., Clemente, J., Ferrer, T., Albericio, F., & Giralt, E. (1997) The use of the Nbb-resin for the solid-phase synthesis of peptide alkylesters and alkylamides: Synthesis of leuprolide. Tetrahedron, 53, 3179-3194. #8369
- 1023. Nilsson, K.G.I., Ljunger, G., & Melin, P.M. (1997) Glycosidase-catalysed synthesis of glycosylated amino acids: Synthesis of Galnac-alpha-Ser and Glcnac-beta-Ser derivatives. Biotechnol. Lett., 19, 889-892. #10304
- 1024. Niu, J.K., & Lawrence, S. (1997) Nonphosphorylatable tyrosine surrogates Implications for protein kinase inhibitor design. Journal of Biological Chemistry, 272, 1493-1499. #8242
- 1025. Noda, M., & Kiffe, M. (1997) New mild acid-labile protecting groups for the guanidino function of N-alpha-fluorenylmethoxycarbonyl-l-arginine in solid-phase peptide synthesis: 10,11-Dihydro-5h-dibenzo[a,d]cyclohepten-5-yl, 2-methoxy-10,11-dihydoro-5h-dibenzo[a,d]cyclopenten-5-yl and 5h-dibenzo[a,d]cyclohepten-5-yl groups. Int. J. Peptide Prot. Res., 50, 329-335. #10305
- 1026. Nokihara, K., Hayakawa, T., Naruse, S., Ando, E., & Wray, V. (1997) Synthesis, solution structure, binding activity, and cGMP activation of human guanylin and its disulfide isomer. Regul. Peptides, 70, 111-120. #9945
- 1027. Norinder, U., Rivera, C., & Unden, A. (1997) A quantitative structure-activity relationship study of some substance P-related peptides: A multivariate approach using PLS and variable selection. J. Pept. Res., 49, 155-162. #9226
- 1028. O'Donnell, M.J., Lugar, C.W., Pottorf, R.S., Zhou, C., Scott, W.L., & Cwi, C.L. (1997) Solid-phase synthesis of unnatural amino acids using unactived alkyl halides. Tetrahedron Lett., 38, 7163-7166. #9560
- 1029. Offer, J., Quibell, M., and Johnson, T. A new generation of backbone-amide protection. 1997. Poster presented at the Fifth International Symposium on Solid Phase Synthesis & Combinatorial Chemical Libraries, London, England, UK on 2nd-6th September 1997. #9607
- 1030. Ohta, N., Mochizuki, T., Hoshino, M., Jun, L., Kobayashi, H., & Yanaihara, N. (1997) Adrenocorticotropic hormone-releasing activity of urotensin i and its fragments in vitro. Int. J. Peptide Prot. Res., 50, 178-183. #10309
- 1031. Okada, Y., Wang, J., Yamamoto, T., Yokoi, T., & Mu, Y. (1997) Amido acids and peptides. L.Development of a novel N-pi-protecting group for histidine, N-pi-2-adamantyloxymethylhistidine, and its application to peptide synthesis. Chem. Pharm. Bull. Tokyo, 45, 452-456. #8521
- 1032. Paquet, A., Blackwell, B., Johns, M., & Nikiforuk, J. (1997) Synthesis of phosphotyrosine-containing peptides using bis-(2,2,2-trichloro)ethyl groups for phosphate protection. Int. J. Peptide Prot. Res., 50, 262-268. #10311

- 1033. Pascal, R., & Sola, R. (1997) Calcium-promoted hydrolysis of N-acylureas allows mild release of peptides anchored with the Dpr (Phoc) linker to hydrophilic resins. Tetrahedron Lett., 38, 4549-4552. #8873
- 1034. Paulitz, C., & Steglich, W. (1997) Stereoselective modification of a cyclopentapeptide via an alpha-(ethylthio)glycine residue. J. Org. Chem., 62, 8474-8478. #10318
- 1035. Paulsen, H., Schleyer, A., Mathieux, N., Meldal, M., & Bock, K. (1997) New solid-phase oligosaccharide synthesis on glycopeptides bound to a solid phase. J. Chem. Soc. Perkin Trans. 1, 281-293. #12522
- 1036. Pawlak, D., Chung, N.N., Schiller, P.W., & Izdebski, J. (1997) Synthesis of a novel side-chain to side-chain cyclized enkephalin analogue containing a carbonyl bridge. J. Peptide Sci., 3, 277-281. #9384
- 1037. Pehk, T., & Uri, A. (1997) Synthesis and structural characterization of conjugates of adenosine and tetra-aspartate, novel analogs of ATP. Bioorg. Med. Chem. Lett., 7, 2159-2164. #10321
- 1038. Peluso, S., Dumy, P., Eggleston, I.M., Garrouste, P., & Mutter, M. (1997) Protein mimetics (TASP) by sequential condensation of peptide loops to an immobilised topological template. Tetrahedron, 53, 7231-7236. #9101
- 1039. Perich, J.W. (1997) Synthesis of phosphopeptides using modern chemical approaches. Methods Enzymol., 289, 245-266. #9995
- 1040. Pernerstorfer, J., Schuster, M., & Blechert, S. (1997) Cyclisation/cleavage of macrocycles by ring-closing metathesis on solid support: Conformational studies. Chem. Commun., 1949-1950. #12523
- 1041. Pfeifer, M.E., Linden, A., & Robinson, J.A. (1997) Synthesis of a novel tricyclic dipeptide template and its incorporation into a cyclic peptide mimetic containing an NPNA motif. Helv. Chim. Acta, 80, 1513-1527. #9400
- 1042. Pichette, A., Voyer, N., Larouche, R., & Meillon, J.C. (1997) A useful method for the preparation of fully protected peptide acids and esters. Tetrahedron Lett., 38, 1279-1282. #8857
- 1043. Pinilla, C., Appel, J.R. & Houghten, R.A. (1997) The generation of peptide combinatorial libraries using tea bag synthesis: Identification of B-cell epitopes. In I. Lefkovits (Ed.), Immunology Methods Manual. (pp. 837-845). Academic Press Limited, London. #8690
- 1044. Pothion, C., Paris, M., Heitz, A., Rocheblave, L., Rouch, F., Fehrentz, J.A., & Martinez, J. (1997) Use of ozonolysis in the synthesis of C-terminal peptide aldehydes on solid support. Tetrahedron Lett., 38, 7749-7752. #9679
- 1045. Pyun, J.C., Cheong, M.Y., Park, S.H., Kim, H.Y., & Park, J.S. (1997) Modification of short peptides using epsilon-aminocaproic acid for improved coating efficiency in indirect enzyme-linked immunosorbent assays (ELISA). J. Immunol. Method, 208, 141-149. #10334
- 1046. Qabar, M.N., Urban, J., & Kahn, M. (1997) A facile solution and solid phase synthesis of phosphotyrosine mimetic L-4-[diethylphosphono(difluoromethyl)]-phenylalanine (F(2)Pmp(Eto)(2)) derivatives. Tetrahedron, 53, 11171-11178. #9413
- 1047. Rano, T.A., Timkey, T., Peterson, E.P., Rotonda, J., Nicholson, D.W., Becker, J.W., Chapman, K.T., & Thornberry, N.A. (1997) A combinatorial approach for determining protease specificities: Application to interleukin-1beta converting enzyme (ICE). Chem. Biol., 4, 149-155. #9284
- 1048. Richter, L.S., & Desai, M.C. (1997) A TFA-cleavable linkage for solid-phase synthesis of hydroxamic acids. Tetrahedron Lett., 38, 321-322. #8244
- 1049. Rodionova, L.N., Zagranichny, V.E., Rodionov, I.L., Lipkin, V.M., & Ivanov, V.T. (1997) The total solid phase synthesis of the gamma subunit of cGMP phosphodiesterase from bovine retina and some physicochemical properties of the synthetic protein [russian]. Bioorganicheskaia Khimiia, 23, 933-948. #10498
- 1050. Rodriguez, E.C., Winans, K.A., King, D.S., & Bertozzi, C.R. (1997) A strategy for the chemoselective synthesis of O-linked glycopeptides with native sugar-peptide linkages. J. Amer. Chem. Soc., 119, 9905-9906. #10342
- 1051. Romoff, T.T., & Goodman, M. (1997) Urethane-protected N-carboxyanhydrides (UNCAs) as unique reactants for the study of intrinsic racemization tendencies in peptide synthesis. J. Pept. Res., 49, 282-292. #9951

- 1052. Rosenthal, K., Karlstrom, A., & Und n, A. (1997) The 2,4-dimethylpent-3-yloxycarbonyl (Doc) group as a new, nucleophile-resistant protecting group for tyrosine in solid phase peptide synthesis. Tetrahedron Lett., 38, 1075-1078. #8410
- 1053. Rosse, G., Sequin, U., Mett, H., Furet, P., Traxler, P., & Fretz, H. (1997) Synthesis of modified tripeptides and tetrapeptides as potential bisubstrate inhibitors of the epidermal growth factor receptor protein tyrosine kinase. Helv. Chim. Acta, 80, 653-670. #9086
- 1054. Rovatti, L., Masin, B., Catinella, S., & Hamdan, M. (1997) Crudes of synthesis of neuropeptide Y and beta-amyloid peptide examined by liquid chromatography electrospray tandem mass spectrometry. Rapid Commun. Mass Spectrom., 11, 1223-1229. #9525
- 1055. Sabirov, A.N., Kim, Y.D., Kim, H.J., & Samukov, V.V. (1997) FMOC- and NSC-groups as a base labile N(alpha)-amino protection: A comparative study in the automated SPPS. Protein Peptide Lett., 4, 307-312. #9742
- 1056. Safavy, A., Buchsbaum, D.J., Qin, H., & Khazaeli, M.B. (1997) Synthesis of bombesin analogues for radiolabeling with rhenium-188. Cancer, 80, 2354-2359. #9926
- 1057. Samson, I., Rozenski, J., Samyn, B., van Aerschot, A., van Beeumen, J., & Herdewijn, P. (1997) Screening a random pentapeptide library, composed of 14 D-amino acids, against the COOH-terminal sequence of fructose-1,6-bisphosphate aldolase from Trypanosoma brucei. Journal of Biological Chemistry, 272, 11378-11383. #9247
- 1058. Sanchez, A., & Smith, A.J. (1997) Capillary electrophoresis. Methods Enzymol., 289, 469-478. #10002
- 1059. Sarantakis, D., & Bicksler, J.J. (1997) Solid phase synthesis of sec-amides and removal from the polymeric support under mild conditions. Tetrahedron Lett., 38, 7325-7328. #9570
- 1060. Sasaki, Y., & Abe, J. (1997) Protection of Psi(CH2NH) peptide bond with 2,4-dimethoxybenzyl group in solid-phase peptide synthesis. Chem. Pharm. Bull. Tokyo, 45, 13-17, #8233
- 1061. Satoh, T., Aramini, J.M., Li, S., Friedman, T.M., Gao, J., Edling, A.E., Townsend, R., Koch, U., Choksi, S., Germann, M.W., Korngold, R., & Huang, Z. (1997) Bioactive peptide design based on protein surface epitopes. A cyclic heptapeptide mimics CD4 domain 1 CC' loop and inhibits CD4 biological function. Journal of Biological Chemistry, 272, 12175-12180. #11665
- 1062. Scarfi, S., Gasparini, A., Damonte, G., & Benatti, U. (1997) Synthesis, uptake, and intracellular metabolism of a hydrophobic tetrapeptide-peptide nucleic acid (PNA)-biotin molecule. Biochem. Biophys. Res. Commun., 236, 323-326. #9366
- 1063. Schleyer, A., Meldal, M., Manat, R., Paulsen, H., & Bock, K. (1997) Direct solid-phase glycosylations of peptide templates on a novel PEG-based resin. Angew. Chem. Int. Ed., 36, 1976-1978. #9631
- 1064. Schneider, C.H. (1997) Peptides in Immunolgy. John Wiley & Sons, New York. ISBN 0471965901 #11586
- 1065. Scholten, J.D., Zimmerman, K.K., Oxender, M.G., Leonard, D., Sebolt-Leopold, J.S., Gowan, R., & Hupe, D.J. (1997) Synergy between anions and farnesyldiphosphate competitive inhibitors of farnesyl-protein transferase. Journal of Biological Chemistry, 272, 18077-18081. #9456
- 1066. Scott, W.L., Zhou, C., Fang, Z., & O'Donnell, M.J. (1997) The solid phase synthesis of alpha, alpha-disubstituted unnatural amino acids and peptides (di-UPS). Tetrahedron Lett., 38, 3695-3698. #8853
- 1067. Sebastian, D., Heuser, A., Schulze, S., & Waldmann, H. (1997) Selective enzymatic removal of protecting groups from phosphopeptides: Chemoenzymatic synthesis of a characteristic phosphopeptide fragment of the RAF-1 kinase. Synthesis-Stuttgart, 1098-1108. #10356
- 1068. Seitz, O., & Wong, C.H. (1997) Chemoenzymatic solution- and solid-phase synthesis of O-glycopeptides of the mucin domain of madcam-1: A general route to O-lacnac, O-sialyl-lacnac, and O-sialyl-lewis-x peptides [review]. J. Amer. Chem. Soc., 119, 8766-8776. #10357
- 1069. Seitz, O., & Kunz, H. (1997) HYCRON, an allylic anchor for high-efficiency solid phase synthesis of protected peptides and glycopeptides. J. Org. Chem., 62, 813-826. #8389
- 1070. Shapiro, G., Buchler, D., Dalvit, C., Frey, P., del Carmen Fernandez, M., Gomez-Lor, B., Pombo-Villar, E., Stauss, U., Swoboda,

- R., & Waridel, C. (1997) Combined Fmoc-Alloc strategy for a general SPPS of phosphoserine peptides: Preparation of phosphorylation-dependent tau antisera. Bioorg. Med. Chem., 5, 147-156. #9948
- 1071. Sharma, S., & Pasha, S. (1997) Rapid anchoring of cesium salts of Boc-protected amino acids on chloromethyl polystyrene resin using dibenzo-18-crown-6 as a catalyst. Bioorg. Med. Chem. Lett., 7, 2077-2080. #10359
- 1072. Shin, J.A. (1997) Specific DNA binding peptide-derivatized solid support. Bioorg. Med. Chem. Lett., 7, 2367-2372. #9987
- 1073. Shin, S.Y., Lee, M.K., Kim, S.Y., Jang, S.Y., & Hahm, K.S. (1997) The use of multiple antigenic peptide (MAP) In the immunodiagnosis of human immunodeficiency virus infection. Biochem. Mol. Biol. Int., 43, 713-721. #11139
- 1074. Sidorova, M.V., Molokoedov, A.S., Ovchinnikov, M.V., Bespalova, Z.D., & Bushuev, V.N. (1997) Solid phase synthesis of beta-amyloid(1-42) [russian]. Bioorganicheskaia Khimiia, 23, 46-55. #9279
- 1075. Singh, P., Hurrell, C.R., Findlay, J.B.C., & Fishwick, C.W.G. (1997) Two novel amino acid derivatives containing side-chain thioamides for the synthesis of photoactivatable peptides. Bioorg. Med. Chem. Lett., 7, 715-718. #10026
- 1076. Smith, A.J. (1997) Amino acid analysis. Methods Enzymol., 289, 419-426. #10000
- 1077. Soderstrom, K., Choi, H.Y., Berman, F.W., Aldrich, J.V., & Murray, T.F. (1997) N-alkylated derivatives of [D-Pro(10)]dynorphin a-(1-11) are high affinity partial agonists at the cloned rat kappa-opioid receptor. Eur. J. Pharmacol., 338, 191-197. #10369
- 1078. Song, A.I., & Rana, T.M. (1997) Synthesis of an amino acid analogue to incorporate p-aminobenzyl-EDTA in peptides. Bioconjug. Chem., 8, 249-252. #9276
- 1079. Songster, M.F., & Barany, G. (1997) Handles for solid-phase peptide synthesis. Methods Enzymol., 289, 126-174. #9935
- 1080. Stewart, J.M. (1997) Cleavage methods following Boc-based solid-phase peptide synthesis. Methods Enzymol., 289, 29-44. #9933
- 1081. Sueiras-Diaz, J., Jones, D.M., Szelke, M., Leckie, B.J., Beattie, S.R., Beattie, C., & Morton, J.J. (1997) Potent in vivo inhibitors of rat renin analogues of human and rat angiotensinogen sequences containing different classes of pseudodipeptides at the scissile site. Int. J. Peptide Prot. Res., 50, 239-247. #10379
- 1082. Sugimoto, N., & Nakano, S. (1997) Sandwiching interaction of peptides with a porphyrin. Chem. Lett., 939-940. #10380
- 1083. Suhara, Y., Izumi, M., Ichikawa, M., Penno, M.B., & Ichikawa, Y. (1997) Peptide-sugar hybrids: Like peptide, like oligosaccharide. Tetrahedron Lett., 38, 7167-7170. #9561
- 1084. Tam, J.P., & Spetzler, J.C. (1997) Multiple antigen peptide system. Methods Enzymol., 289, 612-637. #10007
- 1085. Tandon, R., Singh, M.P., Haq, W., Dikshit, M., Patnaik, G.K., & Dhar, M.M. (1997) Synthesis of novel tetrapeptides as potential ACE inhibitors. Ind. J. Chem. B-Org. Chem. Med. Chem., 36, 386-390. #9540
- 1086. Tegge, W., & Frank, R. (1997) Peptide synthesis on Sepharose(TM) beads. J. Pept. Res., 49, 355-362. #8702
- 1087. Thieriet, N., Alsina, J., Giralt, E., Guibe, F., & Albericio, F. (1997) Use of Alloc-amino acids in solid-phase peptide synthesis. Tandem deprotection-coupling reactions using neutral conditions. Tetrahedron Lett., 38, 7275-7278. #9564
- 1088. Tholey, A., & Hoffmann, B. (1997) Solid-phase synthesis of tyrosyl H-phosphonopeptides and methylphosphonopeptides. J. Peptide Sci., 3, 186-192. #9890
- 1089. Thompson, P.E., & Hearn, M.T.W. (1997) Fmoc-protected tropane-based amino acids for peptide structure-function studies. Tetrahedron Lett., 38, 2907-2910. #8788
- 1090. Torres, J.L., & Clapes, P. (1997) Neoglycopeptide synthesis and purification in multi-gram scale: Preparation of O-(2,3,4,6-tetra-O-acetyl-beta-D-galactopyranosyl)-N-alpha-fluoren-9-yl-methoxycarbonyl-hydroxyproline and its use in the pilot-scale synthesis of the potent analgesic glycopeptide 01.5-beta-D-galactopyranosyl[DMet2,Hyp5]enkephalinamide. J. Peptide Sci., 3, 99-109. #9946

- 1091. Uray, K. (1997) Analysis of the epitope structure of mucin glycoproteins with synthetic peptides and mucin specific monoclonal antibodies [hungarian]. Magyar Kemiai Folyoirat, 103, 237-240. #9076
- 1092. Valentijn, A.R.P.M., van der Marel, G.A., Sliedregt, L.A.J.M., van Berkel, T.J.C., Biessen, E.A.L., & van Boom, J.H. (1997) Solid-phase synthesis of lysine-based cluster galactosides with high affinity for the asialoglycoprotein receptor. Tetrahedron, 53, 759-770. #8267
- 1093. Vandijk, A.A., Vanwijk, L.L., van Vliet, A., Haris, P., Vanswieten, E., Tesser, G.I., & Robillard, G.T. (1997) Structure characterization of the central repetitive domain of high molecular weight gluten proteins. 1. Model studies using cyclic and linear peptides. Protein Sci., 6, 637-648. #9289
- 1094. Varanda, L.M., & Miranda, M.T.M. (1997) Solid-phase peptide synthesis at elevated temperatures: A search for an optimized synthesis condition of unsulfated cholecystokinin-12. J. Pept. Res., 50, 102-108. #9408
- 1095. Vunnam, S., Juvvadi, P., & Merrifield, R.B. (1997) Synthesis and antibacterial action of cecropin and proline- arginine-rich peptides from pig intestine. J. Pept. Res., 49, 59-66. #9224
- 1096. Wade, J.D., & Tregear, G.W. (1997) Relaxin. Methods Enzymol., 289, 637-646. #10008
- 1097. Walker, M.A. (1997) Protein synthesis by chemical ligation of unprotected peptides in aqueous solution. Angew. Chem. Int. Ed., 36, 1069-1071. #9519
- 1098. Wang, Q.M., Johnson, R.B., Cohen, J.D., Voy, G.T., Richardson, J.M., & Jungheim, L.N. (1997) Development of a continuous fluorescence assay for rhinovirus 14 3C protease using synthetic peptides. Antivir. Chem. Chemother., 8, 303-310. #9458
- 1099. Wang, R., Li, X.X., Hu, X.Y., Ni, J.M., Li, C.L., & Lu, J.F. (1997) The synthesis, bioactivity and paramagnetic resonance of angiotensin II and its spin labelled derivative. Chin. Sci. Bull., 42, 1843-1846. #10679
- 1100. Wang, Y.J.J., Yurttas, L., Dale, B.E., Russell, D.H., Kinsel, G., Prestonschaffter, L.M., Johnson, V., & Hayes, T.K. (1997) MALDI-MS as a monitor of the purification and folding of synthetic eclosion hormone. Peptides, 18, 337-346. #9239
- 1101. Weber, P.J.A., & Beck-Sickinger, A.G. (1997) Comparison of the photochemical behavior of four different photoactivatable probes. J. Pept. Res., 49, 375-383. #9197
- 1102. Webster, K.L., Rutherford, T.J., & Gani, D. (1997) Comparison of solution-phase and solid-phase syntheses of a restrained proline-containing analogue of the nodularin macrocycle. Tetrahedron Lett., 38, 5713-5716. #9317
- 1103. Wellings, D.A., & Atherton, E. (1997) Standard Fmoc protocols. Methods Enzymol., 289, 44-67. #9991
- 1104. Wilson, C.J., Husain, S.S., Stimson, E.R., Dangott, L.J., Miller, K.W., & Maggio, J.E. (1997) p-(4-Hydroxybenzoyl)phenylalanine: A photoreactive amino acid analog amenable to radioiodination for elucidation of peptide-protein interaction application to substance P receptor. Biochemistry, 36, 4542-4551. #9257
- 1105. Wipf, P., & Henninger, T.C. (1997) Solid-phase synthesis of peptide mimetics with (E)-alkene amide bond replacements derived from alkenylaziridines. J. Org. Chem., 62, 1586-1587. #9168
- 1106. Xu, J.X., & Jin, S. (1997) Studies on synthesis and antitumor activities of analogues and segments of papaver somniferum pollen heptadecapeptide [chinese]. Chem. J. Chin. Univ., 18, 1098-1102. #9533
- 1107. Yanai, Y., Irie, K., Ohigashi, H., & Wender, P.A. (1997) Synthesis and characterization of the first cysteine-rich domain of novel protein kinase C. Bioorg. Med. Chem. Lett., 7, 117-122. #8658
- 1108. Yao, S., Ghosh, I., Zutshi, R., & Chmielewski, J. (1997) A pH-modulated, self-replicating peptide. J. Amer. Chem. Soc., 119, 10559-10560. #10428
- 1109. Ye, Y.H., Fan, C.X., Zhang, D.Y., Xie, H.B., Hao, X.L., & Tian, G.L. (1997) Application of four novel organophosphorus compounds as coupling reagents for synthesis of bioactive peptides [chinese]. Chem. J. Chin. Univ., 18, 1086-1092. #9532
- 1110. Yokum, T.S., Tungaturthi, P.K., & McLaughlin, M.L. (1997) Benz[f]tryptophan, a bathochromic analog of tryptophan, synthesis of its N-alpha-t-Boc derivative. Tetrahedron Lett., 38, 5111-5114. #9033

- ·)
- 1111. Yokum, T.S., Bursavich, M.G., Piha-Paul, S.A., Hall, D.A., & McLaughlin, M.L. (1997) Synthesis of a series of polar, orthogonally protected, alpha,alpha-disubstituted amino acids. Tetrahedron Lett., 38, 4013-4016. #9048
- 1112. Yoshikawa, M., Izumi, J., & Kitao, T. (1997) Enantioselective electrodialysis of amino acids with charged polar side chains through molecularly imprinted polymeric membranes containing dide derivatives. Polym. J., 29, 205-210. #9148
- 1113. Zenouaki, I., Kharrat, R., Sabatier, J.M., Devaux, C., Karoui, H., Van Rietschoten, J., Elayeb, M., & Rochat, H. (1997) In vivo protection against androctonus australis hector scorpion toxin and venom by immunization with a synthetic analog of toxin ii. Vaccine, 15, 187-194. #8370
- 1114. Zhang, L., & Tam, J.P. (1997) Orthogonal coupling of unprotected peptide segments through histidyl amino terminus. Tetrahedron Lett., 38, 3-6. #8607
- 1115. Zhang, L.S., & Tam, J.P. (1997) Synthesis and application of unprotected cyclic peptides as building blocks for peptide dendrimers. J. Amer. Chem. Soc., 119, 2363-2370. #9178
- 1116. Zhang, L.S., & Tam, J.P. (1997) Metal ion-assisted peptide cyclization. Tetrahedron Lett., 38, 4375-4378. #9515
- 1117. Zhao, Z.G. & Lam, K.S. (1997) Synthetic peptide libraries. In W.H. Moos, M.R. Pavia, B.K. Kay & A.D. Ellington (Eds.), Annual Reports in Combinatorial Chemistry and Molecular Diversity. (pp. 192-209). ESCOM, Leiden. #8624
- 1118. Zhong, H.M., Greco, M.N., & Maryanoff, B.E. (1997) Solid-phase synthesis of arginine-containing peptides by guanidine attachment to a sulfonyl linker. J. Org. Chem., 62, 9326-9330. #9734

[WebCounter] This page has been seen 22619 times since Sun Mar 15 22:43:13 1998

Last Modified: Sunday, March 18, 2001
For more information contact michal@5z.com

Translate this page to: French

rench

Translate

**PSD** Home Page 1 of 2



# Peptide Synthesis Database













Solid phase peptide synthesis is, in most cases, a straightforward procedure using Boc or Fmoc chemistry. A variety of Bocand Fmoc- derivatives are commercially available and, in addition, many organic acids may be incorporated as N-terminal residues, providing a large structural diversity of synthetic peptides and peptidomimetics. However, it is difficult to overview this diversity because the compounds may be listed in different catalogs. The chemical structure of the compounds, which is useful to visualize the diversity, is also missing in many catalogs. It is with this in mind that we have prepared the Peptide Synthesis Database.

The Biotechnology Research Institute, located in Montreal, is a branch of the National Research Council of Canada. It is the leader in Canadian biotechnology R&D, carrying out state-of-the-art research in molecular biology and biochemical engineering with specific applications in the pharmaceutical and resource industries. The Peptide Chemistry group, responsible for the development of the PSD, plays an important role in pharmaceutical R&D here at the BRI.

#### Get the Database

Obtain your own copy of the PSD by either downloading a copy from this site or ordering the database on CD.

### Register Online

You can now register online at no charge to receive notifications of major updates to the database.

#### Contact Us

If you have any questions or comments regarding the PSD, feel free to write to one of our experts here at the NRC's Biotechnology Research Institute.

# New Database "Drug Library"!

Drug Library is a similar database as PSD except it contains information of drugs which are available as chemicals. Some drugs are available as a kit in 96-well microplates for high throughput screening.

## New in 6.2!

PSD v. 6.2 adds over 300 compounds. Click here to find out more.

Visit the Mirror site of this site at Shimadzu in Japan. SHIMADZU





Home | Search | Help | Suppliers | Links